

ACKNOWLEDGMENTS

I would like to thank Drs. Theeraphap Chareonviriyaphap of Department of Entomology, Faculty of Agriculture, Kasetsart University and Pornpimol Rongnoparut of Department of Biochemistry, Faculty of Science, Mahidol University, Bangkok, Thailand for their generous supports, encouragement, valuable comments and expert guidance on this thesis. I am grateful to Drs. Pongthep Akwatanakul and Lertluk Ngernsiri for providing critical comments on thesis improvement. My sincere thanks go to Dr. Micheal J. Bangs of U.S. Naval Medical Research Unit (NAMRU) for providing mosquito specimens from Indonesia and Timor-Leste. Special thanks go to Ms. Atchariya Prabaripai of Faculty of Liberal Art and Sciences, Kasetsart University at Kamphaengsean Campus, Nakhonpathom Province for her helpful on statistic analysis. Appreciation is also extended to my colleagues who supported to thesis achievement. This research would not have been possible without the mosquito collection team from Department of Entomology, Faculty of Agriculture, Kasetsart University and the assistance of molecular laboratory team from Department of Biochemistry, Faculty of Science, Mahidol University, Bangkok, Thailand.

This thesis is supported by the Center for Agricultural Biotechnology (CAB) through the fund from Subproject Graduate Study and Research in Agricultural Biotechnology under Higher Education Development Project, the Commission on Higher Education, Ministry of Education, the Kasetsart University Research and Development Institute (KURDI), Kasetsart University and the Thailand Research Fund (TRF) Organization, Bangkok, Thailand.

Vithee Muenworn

March 2006