



# **THESIS**

**CHARACTERIZATION OF ENZYMES AND  
PHENOLICS INVOLVED IN THE PERICARP  
HARDENING OF MANGOSTEEN FRUIT STORED  
AT LOW TEMPERATURE**

**SIRIWAN DANGCHAM**

**GRADUATE SCHOOL, KASETSART UNIVERSITY**

**2008**



## THESIS APPROVAL

GRADUATE SCHOOL, KASETSART UNIVERSITY

Doctor of Philosophy (Horticulture)

DEGREE

Horticulture

FIELD

Horticulture

DEPARTMENT

TITLE: Characterization of Enzymes and Phenolics Involved in the Pericarp  
Hardening of Mangosteen Fruit Stored at Low Temperature

NAME: Ms. Siriwan Dangcham

THIS THESIS HAS BEEN ACCEPTED BY

THESIS ADVISOR

( Professor Saichol Ketsa, Ph.D. )

COMMITTEE MEMBER

( Professor Jingtair Siriphanich, Ph.D. )

COMMITTEE MEMBER

( Mrs. Parichart Burns, Ph.D. )

COMMITTEE MEMBER

( Professor Ian B. Ferguson, Ph.D. )

DEPARTMENT HEAD

( Assistant Professor Poonpipope Kasemsap, Ph.D. )

APPROVED BY THE GRADUATE SCHOOL ON \_\_\_\_\_

DEAN

( Associate Professor Vinai Artkongharn, M.A. )

**THESIS**

**CHARACTERIZATION OF ENZYMES AND PHENOLICS INVOLVED IN  
THE PERICARP HARDENING OF MANGOSTEEN FRUIT STORED  
AT LOW TEMPERATURE**

**SIRIWAN DANGCHAM**

**A Thesis Submitted in Partial Fulfillment of  
the Requirements for the Degree of  
Doctor of Philosophy (Horticulture)  
Graduate School, Kasetsart University  
2008**

Siriwan Dangcham 2007: Characterization of Enzymes and Phenolics Involved in the Pericarp Hardening of Mangosteen Fruit Stored at Low Temperature. Doctor of Philosophy (Horticulture), Major Field: Horticulture, Department of Horticulture. Thesis Advisor: Professor Saichol Ketsa, Ph.D. 141 pages.

We investigated chilling injury symptom (pericarp hardening) in mangosteen fruit at two maturities, reddish brown and reddish purple, stored at 6°C (87.0%RH) and 12°C (83.5%RH) for 15 days. Fruit stored at 6°C had greater pericarp firmness than those stored at 12°C and reddish purple fruit had greater pericarp firmness than reddish brown fruit. Fruit stored at 6°C for 9 days and transferred to room temperature (29.5°C, 74.0%RH) for 3 days showed more prominent symptom of pericarp hardening. When pericarp hardening occurred, pericarp firmness and lignin contents increased whilst levels of total phenolics decreased. The main phenolic acids associated with pericarp hardening were identified by HPLC as *p*-coumaric and sinapic acids. *p*-Coumaric acid levels of fruit stored at 6°C decreased whereas those of fruit stored at 12°C did not significantly change. On the contrary, sinapic acid levels of fruit stored at both temperatures increased throughout the storage time. A histochemical study of lignified cells in mangosteen pericarp using safranin O and toluidine blue stains showed that lignin accumulation increased with pericarp hardening. Reddish purple fruit held in 0.25% O<sub>2</sub> during storage at 6°C did not reduce pericarp hardening and showed no significant differences in firmness, lignin and total free phenolics levels when compared with fruit stored in normal air conditions. Fruit still increased in firmness and lignin contents, while total free phenolics decreased under low O<sub>2</sub>. Enzymes in the lignin biosynthetic pathway, namely phenylalanine ammonia lyase (PAL), cinnamyl alcohol dehydrogenase (CAD) and peroxidase (POD), were also determined. PAL and POD activities in pericarp of fruit stored at 6°C increased with increasing storage time. Upon transfer of fruit to room temperature, both PAL and POD activities in the pericarp were higher than those in fruit stored at 6°C, and activities declined at the end of storage. CAD activity in fruit pericarp did not change. The key enzymes of lignification in pericarp hardening of mangosteen fruit stored at low temperature are likely to be PAL and POD, and expression of PAL and lignin peroxidase (LgPOD) genes in fruit pericarp was also determined using semi-quantitative RT-PCR, real-time PCR and northern blot techniques. Expression of PAL and LgPOD genes increased with storage time at low temperature and their enzyme activities. However, low O<sub>2</sub> treatment slightly affected on the expression of PAL and LgPOD genes.

These results suggested that the increase in pericarp firmness of mangosteen fruit resulted from induction of lignin synthesis associated with an increase in PAL and POD activities and gene expression.

---

Student's signature

---

Thesis Advisor's signature

## **ACKNOWLEDGEMENTS**

I wish to express my sincere gratitude to my supervisor, Professor Dr. Saichol Ketsa for valuable advice, critically reviewing the manuscript and his encouragement during my study course. I also wish to express my sincere appreciation to Professor Dr. Ian B. Ferguson, Professor Dr. Jingtair Siriphanich and Dr. Parichart Burns, and the representative of the Graduate School, Associate Professor Dr. Supat Attathom for their kind suggestions on my thesis studies. I greatly appreciated their advice, insight and perspectives that only real experts can convey. The staffs of Postharvest Research Unit, Central Laboratory and Greenhouse Complex, Kasetsart University, Kamphaeng Sean Campus, Nakhon Pathom, Thailand are thanked especially for their hospitality and facilitating the instruments and materials during the research. I am also very grateful to Ms. Judith Bowen, Dr. Sean Bulley and the staffs of HortResearch, Mt Albert Research Centre, Auckland, New Zealand, for their kind assistance.

I am indebted to the Thailand Research Fund grant number PHD/00089/2545 for awarding me a scholarship to undertake this study and for the financial support by the Postgraduate Education and Research Development Project in Postharvest Technology.

Finally, I express my deep appreciation to my family, particularly my parents whose provide constant encouragement and support contributed enormously to my success throughout every stage of my study.

Siriwan Dangcham

February 2008

**TABLE OF CONTENTS**

	<b>Page</b>
TABLE OF CONTENTS	i
LIST OF TABLES	ii
LIST OF FIGURES	vi
LIST OF ABBREVIATIONS	xii
INTRODUCTION	1
OBJECTIVES	3
LITERATURE REVIEW	4
MATERIALS AND METHODS	22
THE EXPERIMENTAL TIME AND PLACES	40
RESULTS	41
DISCUSSION	88
CONCLUSION	98
LITERATURE CITED	100
APPENDIX	124

## LIST OF TABLES

<b>Table</b>		<b>Page</b>
1	Upstream (forward) and downstream (reverse) primers for PCR reactions	30
2	Specific forward and reverse primers for PCR DIG labeling using PCR reactions	34
3	The components of PCR reactions with DIG labeling	34
4	The sequence of specific primers used for reverse transcription followed by semi-quantitative RT-PCR and real-time PCR assays	36
5	The components of reactions for semi-quantitative RT-PCR	38
6	The sizes of amplified cDNA fragments and the sequences of genes involved in lignin biosynthesis in mangosteen pericarp	69

### Appendix Table

1	Firmness, lignin contents and total free phenolics contents of reddish brown and reddish purple mangosteen pericarp stored at 6 and 12°C for 15 days	125
2	Pericarp firmness of reddish brown mangosteen fruit stored at 6 and 12°C and measured immediately and after transfer to room temperature (RT) for 3 days	126
3	Pericarp firmness of reddish purple mangosteen fruit stored at 6 and 12°C and measured immediately and after transfer to room temperature for 3 days	127
4	Lignin contents of reddish brown mangosteen pericarp stored at 6 and 12°C and measured immediately and after transfer to room temperature (RT) for 3 days	128

**LIST OF TABLES (Continued)**

<b>Appendix Table</b>		<b>Page</b>
5	Lignin contents of reddish purple mangosteen pericarp stored at 6 and 12°C and measured immediately and after transfer to room temperature (RT) for 3 days	129
6	Total free phenolics contents of reddish brown mangosteen pericarp stored at 6 and 12°C and measured immediately and after transfer to room temperature(RT) for 3 days	130
7	Total free phenolics contents of reddish purple mangosteen pericarp stored at 6 and 12°C and measured immediately and after transfer to room temperature (RT) for 3 days	131
8	Changes in <i>p</i> -coumaric and sinapic acids of reddish purple mangosteen pericarp stored at 6 and 12°C for 12 days	132
9	Pericarp firmness of reddish purple mangosteen fruit stored at 6°C for 0, 6 and 12 days and transferred to room temperature (RT) for 1, 2 and 3 days	133
10	Lignin contents of reddish purple mangosteen pericarp stored at 6°C for 0, 6 and 12 days and transferred to room temperature (RT) for 1, 2 and 3 days	133
11	Total free phenolics contents of reddish purple mangosteen pericarp stored at 6°C for 0, 6 and 12 days and transferred to room temperature (RT) for 1, 2 and 3 days	134
12	Pericarp firmness of reddish purple mangosteen fruit stored at 6°C in normal air and low O <sub>2</sub> conditions and measured immediately and after transfer to room temperature (RT) for 3 days	135
13	Lignin contents of reddish purple mangosteen pericarp stored at 6°C in normal air and low O <sub>2</sub> conditions and measured immediately and after transfer to room temperature (RT) for 3 days	135



**LIST OF TABLES (Continued)**

<b>Appendix Table</b>		<b>Page</b>
14	Total free phenolics contents of reddish purple mangosteen pericarp stored at 6°C in normal air and low O <sub>2</sub> conditions and measured immediately and after transfer to room temperature (RT) for 3 days	136
15	Phenylalanine ammonia lyase (PAL) activity of reddish purple mangosteen pericarp stored at 6°C in normal air and low O <sub>2</sub> conditions and measured immediately and after transfer to room temperature (RT) for 3 days	136
16	Cinnamyl alcohol dehydrogenase (CAD) activity of reddish purple mangosteen pericarp stored at 6°C in normal air and low O <sub>2</sub> conditions and measured immediately and after transfer to room temperature (RT) for 3 days	137
17	Peroxidase (POD) activity of reddish purple mangosteen pericarp stored at 6°C in normal air and low O <sub>2</sub> conditions and measured immediately and after transfer to room temperature (RT) for 3 days	137
18	Firmness, lignin contents, total free phenolics contents and enzyme activities of reddish purple mangosteen pericarp stored at 6°C in normal air and low O <sub>2</sub> conditions and transferred to room temperature (RT) for 3 days	138
19	Pericarp firmness of reddish purple mangosteen fruit stored at 6°C for 9 days and transferred to room temperature (RT) for 3 days in normal air and low O <sub>2</sub> conditions	139
20	Lignin contents of reddish purple mangosteen pericarp stored at 6°C for 9 days and transferred to room temperature (RT) for 3 days in normal air and low O <sub>2</sub> conditions	139

**LIST OF TABLES (Continued)**

<b>Appendix Table</b>		<b>Page</b>
21	Total free phenolics contents of reddish purple mangosteen pericarp stored at 6°C for 9 days and transferred to room temperature (RT) for 3 days in normal air and low O <sub>2</sub> conditions	140
22	Phenylalanine ammonia lyase (PAL) activity of reddish purple mangosteen pericarp stored at 6°C for 9 days and transferred to room temperature (RT) for 3 days in normal air and low O <sub>2</sub> conditions	140
23	Cinnamyl alcohol dehydrogenase (CAD) activity of reddish purple mangosteen pericarp stored at 6°C for 9 days and transferred to room temperature (RT) for 3 days in normal air and low O <sub>2</sub> conditions	141
24	Peroxidase (POD) activity of reddish purple mangosteen pericarp stored at 6°C for 9 days and transferred to room temperature (RT) for 3 days in normal air and low O <sub>2</sub> conditions	141

## LIST OF FIGURES

Figure		Page
1	The pathway of lignin synthesis [modified from Whetten and Sederoff (1995); Campbell and Sederoff (1996); Boudet (2000); Meyermans <i>et al.</i> (2000)].	18
2	Mangosteen fruit at reddish brown (A) and reddish purple (B) stages were used in this study.	23
3	Pericarp firmness of reddish brown (A and B) and reddish purple (C and D) mangosteen fruit stored at 6°C (○,●) and 12°C (□,■) and measured immediately (○,□) and after transfer to room temperature for 3 days (●,■). Data are means ± S.D. of three replications.	44
4	Lignin contents of reddish brown (A and B) and reddish purple (C and D) mangosteen pericarp stored at 6°C (○,●) and 12°C (□,■) and measured immediately (○,□) and after transfer to room temperature for 3 days (●,■). Data are means ± S.D. of three replications.	45
5	Total free phenolics contents of reddish brown (A and B) and reddish purple (C and D) mangosteen pericarp stored at 6°C (○,●) and 12°C (□,■) and measured immediately (○,□) and after transfer to room temperature for 3 days (●,■). Data are means ± S.D. of three replications.	46
6	Changes in (A) <i>p</i> -coumaric and (B) sinapic acids of reddish purple mangosteen fruit stored at 6°C and 12°C for 0 (□), 6 (▣) and 12 (■) days. Data are means ± S.D. of three replications.	47
7	Non-chilled and chilled pericarp (hardening) of reddish purple fruit stained with (A) safranin O and (B) toluidine blue.	48

## LIST OF FIGURES (Continued)

Figure		Page
8	Pericarp firmness of reddish purple mangosteen fruit stored at 6°C for 0 day (●), 6 days (◆), and 12 days (▲) and transferred to room temperature for 3 days. Data are means ± S.D. of three replications.	50
9	Lignin contents of reddish purple mangosteen pericarp stored at 6°C for 0 day (●), 6 days (◆), and 12 days (▲) and transferred to room temperature for 3 days. Data are means ± S.D. of three replications.	51
10	Total free phenolics contents of reddish purple mangosteen pericarp stored at 6°C for 0 day (●), 6 days (◆), and 12 days (▲) and transferred to room temperature for 3 days. Data are means ± S.D. of three replications.	52
11	Pericarp firmness of reddish purple mangosteen fruit stored at 6°C in (A) normal air (○,●) and (B) low O <sub>2</sub> (□,■) conditions and measured immediately (○,□) and after transfer to room temperature for 3 days (●,■). Data are means ± S.D. of three replications.	55
12	Lignin contents of reddish purple mangosteen pericarp stored at 6°C in (A) normal air (○,●) and (B) low O <sub>2</sub> (□,■) conditions and measured immediately (○,□) and after transfer to room temperature for 3 days (●,■). Data are means ± S.D. of three replications.	56
13	Total free phenolics contents of reddish purple mangosteen pericarp stored at 6°C in (A) normal air (○,●) and (B) low O <sub>2</sub> (□,■) conditions and measured immediately (○,□) and after transfer to room temperature for 3 days (●,■). Data are means ± S.D. of three replications.	57

## LIST OF FIGURES (Continued)

<b>Figure</b>	<b>Page</b>
14 Phenylalanine ammonia lyase (PAL) activity of reddish purple mangosteen pericarp stored at 6°C in (A) normal air (○,●) and (B) low O <sub>2</sub> (□,■) conditions and measured immediately (○,□) and after transfer to room temperature for 3 days (●,■). Data are means ± S.D. of three replications.	58
15 Cinnamyl alcohol dehydrogenase (CAD) activity of reddish purple mangosteen pericarp stored at 6°C in (A) normal air (○,●) and (B) low O <sub>2</sub> (□,■) conditions and measured immediately (○,□) and after transfer to room temperature for 3 days (●,■). Data are means ± S.D. of three replications.	59
16 Peroxidase (POD) activity of reddish purple mangosteen pericarp stored at 6°C in (A) normal air (○,●) and (B) low O <sub>2</sub> (□,■) conditions and measured immediately (○,□) and after transfer to room temperature for 3 days (●,■). Data are means ± S.D. of three replications.	60
17 Pericarp firmness of reddish purple mangosteen fruit stored at 6°C for 9 days (◆) and transferred to room temperature for 3 days in normal air (●) and low O <sub>2</sub> (■) conditions. Data are means ± S.D. of three replications.	63
18 Lignin contents of reddish purple mangosteen pericarp stored at 6°C for 9 days (◆) and transferred to room temperature for 3 days in normal air (●) and low O <sub>2</sub> (■) conditions. Data are means ± S.D. of three replications.	64
19 Total free phenolics contents of reddish purple mangosteen pericarp stored at 6°C for 9 days (◆) and transferred to room temperature for 3 days in normal air (●) and low O <sub>2</sub> (■) conditions. Data are means ± S.D. of three replications.	65

## LIST OF FIGURES (Continued)

Figure		Page
20	Phenylalanine ammonia lyase (PAL) activity of reddish purple mangosteen fruit stored at 6°C for 9 days (◆) and transferred to room temperature for 3 days in normal air (●) and low O <sub>2</sub> (■) conditions. Data are means ± S.D. of three replications.	66
21	Cinnamyl alcohol dehydrogenase (CAD) activity of reddish purple mangosteen fruit stored at 6°C for 9 days (◆) and transferred to room temperature for 3 days in normal air (●) and low O <sub>2</sub> (■) conditions. Data are means ± S.D. of three replications.	67
22	Peroxidase (POD) activity of reddish purple mangosteen fruit stored at 6°C for 9 days (◆) and transferred to room temperature for 3 days in normal air (●) and low O <sub>2</sub> (■) conditions. Data are means ± S.D. of three replications.	68
23	Alignment of partial deduced amino acid sequences of mangosteen PAL cDNA fragment with other plants, American mangrove (AAW51923.2), cassava (AAK60275.1), populus (AAN52279.1), citrus (CAB42793.1), tobacco (ABG75911.1) and beetroot (CAH17686.1). The asterisk (*) means the amino acid residues conserved in that column are identical in all sequences in the alignment. The colon (:) means that conserved substitutions have been observed. The dot (.) means that semi-conserved substitutions are observed.	72
24	Alignment of partial deduced amino acid sequences of mangosteen LgPOD cDNA fragment with other plants, tomato (CAA50597.1), tobacco (P11965), populus (BAA06335.1), sweet potato (CAB494692.1), and gossypium (AAQ67366.1). The asterisk (*) means the amino acid residues conserved in that column are identical in all sequences in the alignment. The colon (:) means that conserved substitutions have been observed. The dot (.) means that semi-conserved substitutions are observed.	73

## LIST OF FIGURES (Continued)

<b>Figure</b>		<b>Page</b>
25	Northern hybridization analysis of (A) PAL and (B) LgPOD expression in reddish purple mangosteen fruit stored at 6 and 12°C for 12 days. Equal loading of RNA was checked by hybridization with a mangosteen 18S rRNA probe.	74
26	Northern hybridization analysis of (A) PAL and (B) LgPOD expression in reddish purple mangosteen fruit stored at 6°C in normal air and low O <sub>2</sub> conditions and measured immediately and after transfer to room temperature for 3 days. Equal loading of RNA was checked by hybridization with a mangosteen 18S rRNA probe.	75
27	Northern hybridization analysis of (A) PAL and (B) LgPOD expression in reddish purple mangosteen fruit stored at 6°C for 9 days and transferred to room temperature for 3 days in normal air and low O <sub>2</sub> conditions. Equal loading of RNA was checked by hybridization with a mangosteen 18S rRNA probe.	76
28	Semi-quantitative RT-PCR of PAL and LgPOD expression in reddish purple mangosteen fruit stored at 6 and 12°C for 12 days.	77
29	Semi-quantitative RT-PCR of PAL and LgPOD expression in reddish purple mangosteen fruit stored at 6°C in normal air and low O <sub>2</sub> conditions and measured immediately and after transferred to room temperature for 3 days.	78
30	Semi-quantitative RT-PCR of PAL and LgPOD expression in reddish purple mangosteen fruit stored at 6°C for 9 days and transferred to room temperature for 3 days in normal air and low O <sub>2</sub> conditions.	79
31	Relative expression of PAL gene in reddish purple mangosteen fruit stored at 6°C (□) and 12°C (■) for 12 days.	82
32	Relative expression of LgPOD gene in reddish purple mangosteen fruit stored at 6°C (□) and 12°C (■) for 12 days.	83

**LIST OF FIGURES (Continued)**

<b>Figure</b>		<b>Page</b>
33	Relative expression of PAL gene in reddish purple mangosteen fruit stored at 6°C in normal air (□,■) and low O <sub>2</sub> (▣,▤) conditions and measured immediately (□,▣) and after transferred to room temperature for 3 days (■,▤).	84
34	Relative expression of LgPOD gene in reddish purple mangosteen fruit stored at 6°C in normal air (□,■) and low O <sub>2</sub> (▣,▤) conditions and measured immediately (□,▣) and after transferred to room temperature for 3 days (■,▤).	85
35	Relative expression of PAL gene in reddish purple mangosteen fruit stored at 6°C for 9 days and transferred to room temperature for 3 days in normal air (□) and low O <sub>2</sub> (■) conditions.	86
36	Relative expression of LgPOD gene in reddish purple mangosteen fruit stored at 6°C for 9 days and transferred to room temperature for 3 days in normal air (□) and low O <sub>2</sub> (■) conditions.	87



## LIST OF ABBREVIATIONS

1-MCP	=	1-methylcyclopropene
4CL	=	4-coumaric acid CoA ligase
AIP	=	2-aminoindan-2-phosphonic acid
APEP	=	(1-amino-2-phenylethyl) phosphonic acid
BSA	=	bovine serum albumin
C3H	=	coumaric acid 3-hydroxylase
C4H	=	cinnamic acid 4-hydroxylase
CAD	=	cinnamyl alcohol dehydrogenase
CCoA3H	=	coumaroyl-CoA 3-hydroxylase
CCR	=	cinnamoyl-CoA reductase
CDP-Star	=	disodium 2-chloro-5-(4-methoxyspiro{1,2-dioxetane-3,2'-(5'chloro) tricycle[3.3.1.1 <sup>3,7</sup> ] decan}-4-yl)-1-phenyl phosphate
CI	=	chilling injury
DIG	=	digoxigenin-11-dUTP
F5H	=	ferulic acid 5-hydroxylase
FAA	=	formalin-acetic alcohol
GAPDH	=	glyceraldehydes-3-phosphate dehydrogenase
ILOS	=	initial low oxygen stress
IPTG	=	isopropyl- $\beta$ -D-thiogalactopyranoside
LB medium	=	Luria-Bertani medium
LgPOD	=	lignin peroxidase
OMT	=	<i>O</i> -methyltransferase
PAL	=	phenylalanine ammonia lyase
POD	=	peroxidase
PPO	=	polyphenol oxidase
PVPP	=	polyvinyl polypyrrolidone
RT-PCR	=	reverse transcriptase polymerase chain reaction
TBA	=	tertiary butyl alcohol
X-Gal	=	5-bromo-4-chloro-3-indolyl- $\beta$ -D-galactopyranoside