

LITERATURE CITED

- Adashi, A., K. Morita and T. Okano. 2006. Efficiency of rice bran for the removal of Fe^{3+} from solution. **J. of Health Science**. 52 (4): 455-458.
- Brinch-Pedersen, H., L.D. Sarensen and P.B. Holm. 2002. Engineering crop plants: getting a handle on phosphate. **Trends in Plant Science**. 7 (3): 118-125.
- Chen, P.S., J.R., T.Y. Toribara and H. Warner. 1956. Microdetermination of phosphorus. **Analytical Chemistry**. 28 (11): 1756-1758.
- Cheryan, M. 1980. Phytic acid interactions in food system . **Crit Rev Food Sci Nutr**. 13 (4): 297-335.
- Cochrane, M.P. 2000. Seed carbohydrates, pp. 85-120. In Black M. and Bewley J.D. (Eds). **Seed technology and its biological basis**. New York. Seeffield Academic Press.
- Copeland, L.O. and M.B. McDonald. 1985. The chemistry of seeds, pp. 46. Burgess Publishing Company. **Principles of seed science and technology**. U.S.A.
- Dellapenna, D. 1999. Nutritional genomics : Manipulating plant micronutrients to improve human health. **Science**. 285: 375-379.
- Eardman, J.W. 1981. Bioavailability of trace minerals from cereals and legumes. **American Association of Cereal Chemists, Inc.** 58 (1); 21-26.
- Gibson, D.M. and AHJ. Ullah. 1988. Purification and characterization of phytase from

cotyledons of germinating soybean seeds. **Arch Biochem Biophys.** 260 (2): 503-513.

Glahn, R.P. and G.M. Wortley. 2002. Inhibition of iron uptake by phytic acid, tannic acid and ZnCl₂: Studies using an in vitro digestion/caco-2 cell model. **J. Agric. Food. Chem.** 50: 390-395.

Glahn, R.P., Z. Cheng and R.M. Welch. 2002. Comparison of iron bioavailability from 15 rice genotypes: Studies using an in vitro digestion/caco-2 cell model. **J. Agric. Food. Chem.** 50: 3586-3591.

Greiner, R. and Others. 2001. Purification and characterization of a phytate-degrading enzyme from germinated Faba beans (*Vicia Faba* var. Alameda). **J. Agric. Food Chem.** 19: 2234-2240.

Greiner, R., M.L. Alminger, N.G. Carlsson, M. Muzquiz, C. Burbano, C. Cuadrado, M.M. Pedrosa and C. Goyoaga. 2002. Pathway of dephosphorylation of myo-inositol hexakisphosphate by phytases of legume seeds. **J. Agric. Food. Chem.** 50: 6865-6870.

Harland, B.F. and E.R. Morris. 1995. Phytate: A good or a bad food component. **Nutritional Research.** 15 (5): 733-754.

Harland, B.F. and G. Narula. 1999. Food phytate and its hydrolysis products. **Nutritional Research.** 19(6): 947-961.

Hatzack, F. and S.K. Rasmussen. 1999. High-performance thin-layer chromatography method for inositol phosphate analysis. **J. of Chromatography B.** 736: 221-229.

- Hayakava, T., Y. Toma and I. Igaue. 1989. Purification and characterization of acid phosphatases with or without phytase activity from rice bran. **Agric.Biol.Chem.** 53(6): 1475-1483.
- Hegeman, C.E. and E.A. Grabau. 2001. A novel phytase with sequence similarity to purple acid phosphatase is expressed in cotyledons of germinating soybean seedling. **Plant Physiology.** 126: 1598-1600 .
- Hubel, F. and B. Erwin. 1996. Maize Root Phytase : Purification, characterization and localization of enzyme activity and its putative substrate. **Plant Physiol.** 112: 1429-1436.
- Kennedy, B.M. and M. Schelstraete. 1975. Chemical, physical, and nutritional Properties of high-protein flours and residual kernel from the overmilling of uncoated milled rice.III. Iron, calcium, magnesium, phosphorus, sodium, potassium, and phytic acid. **American Association of Cereal Chemists.** 52: 173-182.
- Laboure, A.M., J. Gagnon and A.M. Lescure. 1993. Purification and characterization of a phytase (myo-inositol-hexakisphosphate phosphohydrolase) accumulated in maize (*Zea mays*) seedlings during germination. **Biochem J.** 295: 413-149 .
- Larson, S.R., J. N. Rutger, K.A. Young and V. Raboy. 2000. Isolation and genetic mapping of a non-lethal rice (*Oryza sativa* L.) low phytic acid1 mutation. **Crop Sci.** 40: 1397-1405.
- Larson, S.R., K.A. Young, A. Cook, T.K. Blake and V. Raboy. 1998. Linkage mapping


- of two mutations that reduce phytic acid content of barley grain. **Theor Appl Genet.** 97: 141-146.
- Latta, M. and M. Eskin. 1980. A Simple and Rapid Colorimetric Method for Phytate Determination. **American Chemical Society.** 28 (6): 1313-1315.
- Liu, J.C., I. Ockenden, M. Truax and J.N.A. Lott. 2004. Phytic acid-phosphorus and other nutritionally important mineral nutrient elements in grains of wild-type and low phytic acid (*lpa1-1*) rice. **Seed Science Research.** 14: 109-116.
- Loewus, F.A., P.P.N. Murthy. 2000. myo-inositol metabolism in plants. **Plant Science.** 150: 1-19 .
- Lott, J.N.A., I. Ockenden, V. Raboy and G.D. Batten. 2000. Phytic acid and phosphorus in crop seeds and fruit : a global estimate. **Seed Science Research.** 10: 1-33.
- Lott, J.N.A., J.S. Greenwood and G.D. Batten. 1995. Mechanisms and regulation of mineral nutrient storage during seed development, pp. 215-235. In Kigel, J.; Galili, G. (Eds). **Seed development and germination.** New York, Marcel Dekker, Inc.
- Lucca, P., R. Hurrell, I. Potrykus. 2001. Genetic engineering approaches to improve the bioavailability and the level of iron in rice grain. **Theor Appl Genet.** 102 : 392-397.
- Maga, J.A. 1982. Phytate : Its chemistry, occurrence, food interactions, nutritional significance and methods of analysis. **J.Agric.Food Chem.** 30 (1): 1-9.

- Maiti, I.B., A.L. Majumder and B.B. Biswas. 1974. Purification and mode of action of Phytase from *Phaseolus aureus*. **Phytochemistry**. 13: 1047-1051.
- Maiti, I.B. and B.B. Biswas. 1979. Further characterization of phytase from *Phaseolus aureus*. **Phytochemistry**. 18: 316-318 .
- Maugenest, S., I. Martinez and A.M. Lescure. 1997. Cloning and characterization of a cDNA encoding a maize seedling phytase. **Biochem J**. 322: 511-517.
- Maugenest, S. and Others. 1999. Structure of two maize phytase genes and their spatio-temporal expression during seedling development. **Plant Molecular Biology**. 39: 503-514.
- Mukherji, S. and Others. 1971. Changes in phosphorus fraction and phytase activity of rice seeds during germination. **Physiol Plant**. 25: 94-97 .
- O' Dell, B.L., A.R. de Boland and S.R. Koirtyohann. 1972. Distribution of phytate and nutritionally important elements among the morphological components of cereal grains. **J. AGR. FOOD CHEM**. 20 (3): 718-721.
- Otegui, M.S., R. Capp and L.A. Staehelin. 2002. Developing seeds of Arabidopsis store different minerals in two types of vacuoles and in the endoplasmic reticulum. **The Plant Cell**. 14: 1311-1327.
- Palmiano, E.P. and B.O. Juliano. 1973. Changes in the activity of some hydrolases, peroxidase and catalase in the rice seed during germination. **Plant Physiol**. 52: 247-277.

- Raboy, V. 2000. Low-phytic-acid grains. **Food and Nutrition Bulletin**. 21 (4): 423-427.
- Raboy, V. 2001. Seeds for a better future : low phytate grains help to overcome malnutrition and reduce pollution. **Trend in Plant Science**. 6(10): 458-462.
- Raboy, V., P.F. Gerbasi, K.A. Young., S.D. Stoneberg, S.G. Pickett, A.T. Bauman, P.P.N. Murthy, W.F. Sheridan and D.S. Ertl. 2000. Origin and sees phenotype of maize low phytic acid 1-1 and low phytic acid 2-1. **Plant Physiology**. 124: 355-368.
- Raboy, V., K.A. Young, J.A. Dorsch and A. Cook. 2001. Genetics and breeding of seed phosphorus and phytic acid. **J. of Plant Physiology**. 158: 489-497.
- Rasmussen, S.K. and F. Hatzack. 1998. Identification of two low-phytate barley (*Hordeum vulgare* L.) grain mutants by TLC and genetic analysis. *Hereditas*. 129: 107-112
- Rutger, J.N., V. Raboy, K.A.K. Moldenhauer, R.J. Bryant, F.N. Lee and J.W. Gibbons. 2004. Registration of KBNT lpa1-1 low phytic acid germplasm of rice. **Crop Sci**. 44: 362-363.
- Sanberg, A.S., N.G. Carlsson and U. Svanberg. 1989. Effects of inositol tri-, tetra-, penta- and hexaphosphates on in vitro estimation of iron availability. **Journal of Food Sciences**. 54 (1):159-161,186.
- Sanberg, A.S and Others. 1999. Inositol phosphate with different numbers of phosphate groups in fluence iron adsorption in humans. **Am.J.Clin.Nutr**. 70: 240-246.

- Shi J., H. Wang., Y. Wu., J. Hazebroek., R.B. Meeley and D.S. Ertl. 2003. The maize low-phytic acid mutant lpa2 is caused by mutation in ana inositol phosphate kinase gene. **Plant Physiology**. 131: 507-515.
- Verwoerd, T.C. and Others. 1995. Stable accumulation of *Aspergillus niger* phytase in transgenic tobacco leaves. **Plant Physiol**. 109: 1199-1205.
- Welch, R.M. and R.D. Graham. 2004. Breeding for micronutrients in staple food crops from a human nutrition perspective. **Journal of Experimental Botany**. 55 (396): 353-364.
- White, P.J. and M.R. Broadley. 2005. Biofortifying crops with essential mineral element. **Trends in Plant Science**. 10 (12): 586-593.
- Wilcox, J.R., G.S. Premachandra, K. A. Young and V. Raboy. 2000. Isolation of high seed inorganic P, low-phytate soybean mutants. **Crop Sci**. 40: 1601-1605.
- Williams, S.G. 1970. The role of phytic acid in the wheat grain. **Plant Physiol**. 45: 376-381.
- Yoshida, T., K. Tanaka and Z. Kasai. 1975. Phytase activity associated with isolated Aleurone particles of rice grains. **Agr.Biol.Chem**. 39 (1): 289-290.
- Yoshida, K.T., T. Wada, H. Koyama, R. Mizobuchi-Fukuoka and S. Naito. 1999. Temporal and spatial patterns of accumulation of the transcript of myo-inositol-1-phosphate synthase and phytin-containing particles during seed development in rice. **Plant Physiology**. 119: 65-72.

Appendix

Mutagenic breeding

CURRICULUM VITAE

NAME : Miss Tida Amon

BIRTH DATE : September 18, 1972

BIRTH PLACE : Bangkok, Thailand

EDUCATION : YEAR INSTITUTION DEGREE/DIPLOMA)

1994	Sri Nakharinwirot	B.Ed. (Chemistry)
1998	Sri Nakharinwirot	M.S. (Biological Chemistry)

POSITION/TITLE : lecturer

WORK PLACE : Faculty of Science, Chemistry program,
Bansomdejchaopraya Rajabhat University