

THE MAN WHO MADE AN IMPACT -THE SCIENTIFIC WORK OF DR SANGAD BUNOPAS

Clive Burrett^{1,*}

¹*Palaeontological Research and Education Centre, Mahasarakham University,
Kantarawichai, Maha Sarakham, 44150 Thailand*

**Corresponding author: cfburrett@gmail.com*

Received 29 December 2020 ; Accepted 15 January 2021.

Abstract

Sangad Bunopas published extensively on the geology of Thailand and left a legacy of maps, reports stratigraphic nomenclature and insightful papers covering most of the kingdom. His 1981 PhD thesis published in 1982 laid the foundation of modern Thai stratigraphy. His highly cited papers on the tectonics of South East Asia, often with Professor Paul Vella, were based on Sangad's extraordinary first-hand knowledge of Thai geology which allowed him to draw the first reliable tectonic subdivisions of the whole country. Although controversial at the time, these tectonic subdivisions and hypotheses have since been accepted as generally correct in that the Shan-Thai Terrane (as part of Sibumasu) rifted from Australian crust in the early Permian, drifted and warmed as it crossed the Palaeotethyan Ocean and collided with the Indochina Terrane in the late Triassic. His stimulating palaeomagnetically constrained, combined palinspastic-palaeogeographic maps for the Permian to Cenozoic were well in advance at the time and have not been attempted again in the subsequent 40 years.

His controversial later work, which attempted to explain the widespread, anomalous loess in northeast Thailand, the Australasian tektite strewn field, the fluvial deposits in Nakhon Ratchasima (Khorat) Province and their middle Pleistocene faunas and flora as the result of a cometary impact at 788 ka has received some support and some criticism but has nevertheless stimulated a considerable research and a lot of discussion.

Over more than 50 years, Sangad made many major contributions to Thai geology, South East Asian tectonics and meteoritic research. His wit, wisdom and iconoclastic approach to earth and planetary science will be sorely missed by his numerous friends and colleagues.

Keywords: *biography, cometary impact, loess, Quaternary stratigraphy, tectonics, terrane, Thailand*

1. INTRODUCTION

Dr Sangad Bunopas (1937-2020) was born and raised in Chaiyaphum near the edge of the Khorat Plateau in north eastern Thailand, surrounded by areas of scenic geology and geomorphology. To the east lay the Mesozoic sandstones of the Khorat Plateau and to the west the Central Plains with Cenozoic cover of marine Palaeozoic rocks. After graduating from Chulalongkorn University in 1960, Sangad joined the Department of Mineral Resources (DMR) in July 1961. Within a few years, he was sponsored to study abroad by the DMR and supported initially by a Colombo Plan scholarship. He was encouraged and helped in his overseas studies by his wife and by senior DMR staff Pisoot Sudasna Na Ayudhaya

(who became Deputy Permanent Secretary of the Minister of Industry), Saman Buravas, Manus Veraburras and significantly, in the light of his pioneering work on tektites, by Dr Kaset Pitakpaivan who later became Director-General of the Department of Industrial Promotion.

Sangad completed the rigours of a BSc (Honours) degree before progressing to an MSc programme (1965-6) in New Zealand. He returned to regional mapping in Thailand and then from 1977 to 1981 completed a PhD at the Victoria University of Wellington supervised by Professor Paul Vella. Sangad was lucky to study at Victoria University as the geology of New Zealand is diverse, rugged, spectacular and active (**Figs. 2-3**) and the local geologists are renowned for their excellence,

enthusiasm for fieldwork and all-round abilities. This was excellent training which he later put to good use working on Thai geology. He graduated with an MSc in 1966 at a time when the theory of plate tectonics was being developed and then applied to orogenic belts around the world. Sangad's regional mapping and his PhD work on palaeomagnetism would soon lay the foundations for his new ideas on the tectonics of South East Asia



Fig. 1 Sangad Bunopas. Photos kindly supplied by the Bunopas family

2. REGIONAL GEOLOGY OF THAILAND

His early work for the DMR, based in Bangkok, was in regional mapping and special projects. Unlike today, there was little infrastructure in Thailand in the late 1960's to 1970's. Roads were very bad, vehicles old and poorly maintained, road cuts were limited and thick jungle still covered large areas of the country. Communist insurgency made some areas dangerous for government personnel. "Traverses of the Western Mountains and some parts of the Sukhothai Foldbelt often require transportation by elephant..." (Raksakulwang and Bunopas, 1985, p.9). With some understatement they reported that progress on foot and by elephant made mapping very slow.

His early assignments were in several provinces on a wide variety of rocks and economic deposits. He investigated the geology and mineral resources of Amphoe Mae Thaeng in Chiang Mai Province in 1961, lignite and oil shale in Li, Lamphun Province (1962 and 1963), the geology of Song in Phrae Province in 1967, the geology of the Nam Phrom Dam in Chaiyaphum Province (Bunopas, 1971) and tin deposits in Rayong Bay in eastern Thailand in 1973. With DMR colleagues, Sangad produced 1: 250,000 maps and reports for geologically significant quadrangles in central Thailand such as Phitsanalok (Bunopas, 1974b and 1976a), Nakhon Sawan (Bunopas, 1976b, 1980) and Suphanburi (Bunopas, 1976c). Part of the Suphanburi mapping project led to an important paper in the first volume of the Journal of the

Geological Society of Thailand on the fascinating geology of Bo Phloi and Uthai Thani in western Thailand (Bunopas and Bunjitradyula, 1975). They mapped a variety of Palaeozoic rocks, metamorphic and igneous rocks with comments on the economic resources including the famous Bo Phloi sapphires. They discussed the confusing lithostratigraphic nomenclature at the time concerning the Kanchanaburi Series, Phuket Group and Tanaosri Group and most significantly recognised the several thrust faults such as the large, low-angle Chong Insi Thrust where the Siluro-Devonian rocks are overthrust on gneisses. The inclusion of structural data, and especially the recognition of thrust faults, was very unusual for Thai geology at that time and was a feature of many of Sangad's (with DMR colleagues) regional papers including the still very interesting structural geology chapter 7 of his 1981 thesis. An example of his emphasis on structural geology is shown in **Fig.4**.

An important study on the famous (and as a prison, infamous) island of Ko Tarutao in southern Thailand followed in 1978 with DMR colleagues Sahat Muenlek and Veerapong Tansuwan (Bunopas et al., 1978). Cambrian fossils had previously been found there by DMR director Saman Buravas in 1958 and described by Teichii Kobayashi but little work had been done on these important and fossiliferous Lower Palaeozoic sections in the intervening 20 years. They mapped and described the sandstones and limestones and recognised important faults thereby laying the foundations of scores of palaeontological papers, reports and theses in subsequent years. Work on Ko Tarutao and in mainland Satun was strongly encouraged and supported by Sangad. These foundational studies and discoveries eventually led to Satun being recognised as the first UNESCO Geopark in Thailand.

Another regional study of note was written-up by Lertsin Raksakulwang and Sangad in 1985 on the complex geology of 3000 square kilometres of Mae Hong Son Province in northwest Thailand. They included structural, lithostratigraphic and sedimentological details including many cross sections, stratigraphic columns and Landsat maps of lineaments with much of the work carried out on elephants and on foot across mountainous terrain (Raksakulwang and Bunopas, 1985). Within a few years this area became another well-studied and palaeontologically and tectonically significant area.

Sangad also produced several important and influential syntheses of stratigraphic correlations across Thailand (Bunopas 1974a,

1976, 1983, 1992b, 1994). Some of these papers were written prior to his massive encyclopedia of Thai geology (his PhD thesis) and others subsequent to Bunopas (1981, 1982). All were influential and there is hardly a chapter of recent books on the geology of Thailand that do not quote these studies extensively.

A paper with Sirot Salypongse, Father Henri Fontaine and Daniel Vachard (Bunopas et al., 1983) summarised the Permian stratigraphy of

Thailand and included important new palaeontological data. Sangad also worked with Father Fontaine on the fossiliferous Devonian limestone of Loei Province, in northeastern Thailand. This was followed by many significant papers and monographs by Fontaine, Vachard and DMR colleague Dr Voravudh Suteethorn, mainly on the Devonian to Permian palaeontology of Thailand.



Fig. 2 Gladstone East Coast Road. Cold and wet South Island, New Zealand – a long way from tropical Thailand. (From Bunopas, 1966, MSc thesis, Victoria University, New Zealand).



Fig. 3 Eringa Cyclothems, near Eringa Station, New Zealand. The area was studied in Wairarapa, a little north of Wellington. In the Wainuiru River Valley in Gladstone near Masterton. Miocene and Pliocene mudstones younging west (left hand), are overlain by

limestone cyclothem with coquina limestones and the terrestrial Hawera Series (From Bunopas, 1966, Victoria University, Wellington, MSc thesis).




Fig. 4 From Bunopas (1982, fig.42) showing an example of the structural cross-sections in his thesis. Section from north of Lampang to west of Phrae, northern Thailand.

3. TECTONICS OF SOUTH EAST ASIA

Armed with an enormous data-base of Thai geology acquired after detailed studies in every corner of the kingdom, Sangad developed a new tectonic model for the whole of South East Asia with his mentor and supervisor Professor Paul Vella. This model was first published in Bunopas and Vella (1978) and enlarged on in subsequent years (Bunopas and Vella 1983a, Bunopas et al., 1990, 2003, 2004). The model was developed fully in chapter 9 of his 801 page tour de force PhD from Wellington which was then published as special paper number 5 of the Geological Survey of the DMR (Bunopas, 1981, 1982). Unusually for a PhD thesis, the first 443 pages is a comprehensive, country-wide summary and it may well be considered the first book on the geology of Thailand. It has been cited over 400 times. Any geologist working in Thailand can still dip into these chapters and find something significant that they either did not know or had forgotten.

The first 443 pages of Sangad's thesis contains numerous maps and useful lithostratigraphic and correlation charts with the text full of new paleontological lists and sedimentological insights. Sangad included thin section descriptions and numerous chemical analyses of igneous rocks. He redefined and clarified old, loosely defined lithostratigraphic units and on almost every page named and


defined numerous new formations and groups, many of which are still in every-day use. Bunopas (1982) may be regarded as the foundation of modern Thai stratigraphy.

The 100 pages of Chapter 8 of his thesis concern Sangad's palaeomagnetic studies on both Palaeozoic and Mesozoic rocks. His preliminary palaeomagnetic results were published with Dr Kaset Pitakpaivan and J. Sukroo in 1978 (Bunopas et al., 1978). His conclusions on the rotation of Indochina and his Mesozoic results have been confirmed by later work. Unfortunately many of his Permian samples had high alpha 95's ('statistical uncertainty' or scatter) which were unacceptably large but even his low inclination results with moderate alpha 95's on Permian limestones appear to have been validated by recent studies. Sangad fully realised the limitations of his Palaeozoic palaeomagnetic studies and on returning from Wellington collected a large number of orientated cores on Cambrian and younger Palaeozoic rocks. Sadly, hardware and software problems meant that no further reliable results could be obtained.

Generally Sangad's basic tectonic concepts have stood the test of time. He drew his Shan-Thai Terrane to include western Thailand and Malaysia and eastern Myanmar and regarded what is now termed the Inthanon Zone or Terrane as a back-arc basin of his Sukhothai Arc. He identified the ophiolitic Nan Suture and for the first time suggested its extension to the ophiolite of

Sra Kaeo in eastern Thailand which he briefly described - also for the first time. He emphasised the contrast between the late Carboniferous and early Permian faunas of the Shan-Thai (cool Gondwana) and Indochina (tropical Cathaysian) terranes which were separated by the Nan-Sra Kaeo suture. He subdivided Thailand into distinctive tectonic belts, offset by major transcurrent faults and showed their palaeogeographic evolution through time (Fig.5).

Sangad suggested Shan-Thai rifted from the Australian sector of Gondwana, with the deposition of the glacial pebbly mudstones of the Kaeng Krachan Group (Bunopas and Vella, 1984, 1989) in the early Permian, drifted into lower and warmer palaeolatitudes during the middle Permian to Early Triassic and collided with Indochina in the Norian (Late Triassic).



Silurian-Devonian lithofacies (excepting peninsular Thailand). cs, carbonate shelf facies; cc, continent derived clastics, sandstone and shale; l, land; bab, back-arc basin (chert and shale); va, metavolcanics (metatuff and agglomerate); fab, fore-arc basin (tuff, greywacke, marble and chert); t, trench deposits, pelitic greenschist. PSA, Pha Som Ophiolite and SKO, Sra Kaeo Ophiolite thought to have been emplaced in the Triassic; cgs, shelf sandstone, shale and limestone. Note: No Silurian-Devonian thick marginal deposits are known east of the ophiolite belt. MP, Mae Ping Fault Zone; TP, Three Pagodas Fault Zone.

Fig. 5 From Bunopas (1982, fig 141b) showing one of several diagrams outlining Sangad's major tectonic subdivisions of Thailand (for the Siluro-Devonian) offset by major transcurrent faults. Note the recognition, mapping and offset of the Sra Kaeo ophiolite in eastern Thailand and the proposed back-arc basin in northern Thailand in what we would now term the Inthanon Zone (or terrane).

Sangad did not confine himself to the Palaeozoic-Triassic but wrote an interesting paper on Cenozoic rifting and the development of

the Gulf of Thailand (Bunopas and Vella, 1983b). Unusually, he attempted to remove Mesozoic deformation and Cenozoic faults and their movements and incorporated his palaeomagnetic results, showing block rotations, in order to draw combined palinspastic palaeogeographic restorations for South East Asia (Bunopas 1981, figs 151-155b).

These restorations (Fig.6) have not been attempted to the same level of detail since and contrast markedly with the inchoate 'blobs' drawn on many recent palaeogeographic maps. A feature here is that Sangad bravely extended his initial ideas where most conventional geologists would have finished. This pushing of hypotheses to their limits - and sometimes beyond was also characteristic of his later work on impacts.

Apart from the excision of the Upper Palaeozoic limestones and Devonian to Triassic cherts of the Inthanon Zone from the Shan-Thai Terrane, arguments on subduction directions and the inclusion of Shan-Thai in an enlarged Sibumasu Terrane, his initially controversial proposals, have been confirmed by most subsequent studies. Today few would disagree with his major conclusion that Shan-Thai (as a part of Sibumasu) was part of northwest Australian crust that rifted off in the early Permian, warmed-up in its drift across Palaeotethys and collided with Indochina in the Late Triassic.

4. IMPACTS, TEKTITES, LOESS AND COMETS

DMR Director General Dr Kaset Pitakpaivan was one of the first to study tektites in Thailand and it was not surprising that he influenced the young Sangad to eventually study them. Dr Kaset worked with Dr Virgil Barnes from Texas and made the first regional scientific study of these fascinating black glasses at about the same time that Sangad joined the DMR (Barnes and Pitakpaivan, 1962). At the time, many prominent and brilliant planetary scientists argued that tektites were extraterrestrial with the most prominent theory being that they were blasted from volcanoes on the moon (Chapman and Larsen, 1963; O'Keefe, 1976). Barnes and Pitakpaivan collected tektites throughout northeast Thailand and emphasised the importance of layered or Muong Nong tektites including the large 12.8 kg specimen donated by Dr Kaset to the DMR museum (Barnes, 1971).

Huge samples of Muong Nong tektites which were offered for sale by Mr Boomun Poonyathiro, owner of the House of Gems, Bangkok weigh up to 24 kg (Fig. 7). Detailed field

studies in Ubon Ratchathani Province by P. Fiske, Prinya Phutthapiban and John Wasson (1996) confirmed the abundance and stratigraphic position of Muong Nong tektites on a lateritic

surface overlain by loess. All subsequent workers showed conclusively that Australasian strewn field tektites are not extraterrestrial but the result




Fig. 6 A novel and iconoclastic palinspastic reconstruction of Thailand adjacent to northwest Australia in the early Carboniferous, figure 144 in Bunopas (1981, 1982). This is one of many stimulating and provocative palinspastic reconstructions in his thesis.

of bolide impact or impacts at 788 ka, centred in the Indochina region that blasted tektites all the way to Tasmania and as microtektites across the Indian Ocean, western Pacific Ocean and all the way to Antarctica. Tektites are so abundant in South East Asia that in some areas almost every villager has a collection and jewellery shops in Charoen Krung, Bangkok literally have buckets full of them.

In 1987 Dr Ramsay Ford (with the writer and Sangad) toured northeast Thailand (Isarn) and collected tektites from almost every province. We found a large disk tektite horizontally embedded in the middle of a 3- thick section of loess near Khon Kaen and found that the majority of tektites were restricted to the junction of a distinctive laterite and an overlying section of loess or rarely within the loess itself (Ford, 1988). The loess covers NE Thailand extends to the

Central Plain and varies between 4 m and 1 m in thickness and compares with the famous yellow earth of China but differs in detail (Sibava, 1993; Nichol and Nichol, 2015, see Bunopas, 2021 this volume). An interesting study by Paul Vella, Sangad and Vorakul Kaewyana that does not appear to have been followed-up, showed that loess extends to northern Thailand and covers river terrace deposits. This potentially provides constraints on dating river terraces and also provides clues to uplift rates (Vella et al., 1987). Loess in Europe and China is related to windy environments close to glaciated areas and its ubiquity in northeastern Thailand is anomalous and difficult to explain on current models of loess formation (Nichol and Nichol, 2015). Sangad was already interested in the tektite problem and he soon wrote papers on the catastrophic effects of a large impact in Indochina close to the area of the

large Muong Nong tektites eastern Isarn, Laos and Vietnam (Bunopas, 1989, 1990, 1992a). He specifically included the loess as a result of an impact and therefore suggested a c.800 ka age on the basis of finding pristine tektites within and just below this enigmatic deposit. Sirot Salyapongse and Kaset Pitakpaivan studied the Thai loess and compared it to Chinese loess. They found glass beads in the Thai loess and also, interestingly, in Pleistocene trees (Salyapongse and Pitakpaivan, 1999).



Fig. 7 Probably the largest Muong Nong tektite yet found and was on display at the House of Gems in Charoen Krung, Bangkok. This and related specimens may be from Ubon Ratchathani Province on the Thai-Lao border or possibly from Laos. This specimen was shown to Sangad Bunopas, Ramsay Ford, Clive Burrett and Kieren Howard at various times by Mr Boomun Poonyathiro. Weight is 24 kg. 1999 photograph kindly supplied by Prof. Kieren Howard.

Sangad coined the term *catastroloess* for the Thailand loess. He included fluvial deposits from along the palaeo-Mun River valley, which clearly are not loess and it is unlikely that this term will survive.

He soon extended the catastrophic impact model to include the fluvial sands along the Mun River Valley of Nakhon Ratchasima Province (Khorat) containing large charcoalified and partially burnt logs, small plant material and pollen (some burnt) and numerous vertebrate fossils (Bunopas and Khositantont, 1999c, Bunopas et al., 1997, 1998, 1999 a, b, 2001, 2002a, b, 2003, 2004, 2005 and 2007; Charusiri et al., 2002). When Sangad showed me the Khorat sandpits and their abundant apparently burnt logs, I was immediately struck by the resemblance to the multitude of trees felled by the huge air-blast of the Tunguska, Siberia explosion in 1908.

Sangad's correlation of the Khorat sands with the 788 ka impact was a step too far for many as it was already apparent that many but not all of the vertebrates in the sandpits are Miocene in age and not Pleistocene (Chaimanee, 2004, 2007; Duangkrayom et al., 2017). The situation is complicated by the sandpit operators who use high-powered hoses to sort and collect the poorly consolidated sands. As a result, vertebrate fossils are rarely found in situ but are found either much lower in the stratigraphy or at the bottom of the pits. The biostratigraphic situation was clarified by Sato (2002) and it appears that definite in situ though of course, fluviially transported, Miocene fossils are generally found in the basal parts of the pits and middle Pleistocene such as *Stegodon*, crocodiles and turtles in the upper parts (e.g. Lauprasert et al., 2019; Naksri et al., 2019). Of course, in a flood-prone, fluvial environment older vertebrates may be reworked into younger sands so the biostratigraphic and taphonomic context becomes complicated. The situation in northeast Thailand and beyond is also complicated by the difficulties dating Quaternary deposits with all available methods being subject to considerable problems, caveats and uncertainties.

Sangad encouraged detailed work on the Khorat sand-pits by Kieren Howard (1999) and then with Howard collaborated with Dr Peter Haines (Australia) and palaeomagnetist Professor Jason Ali (Hong Kong) (helped by the late Dr Somboon Khositantont and the writer) in a magnetostratigraphic study of the pits. The advantage of magnetostratigraphy is that if consecutive units are present then they have not been reworked or reset. The disadvantage is that short geomagnetic excursions, such as within the Brunhes Epoch, may be mistaken for epoch boundaries or vice versa. Haines et al., (2004) revealed a transition from present-day Normal magnetization to Reversed at 14 m depth at Ban Tha Chang and at 9 m depth 40 km away at Chum Phuang (**Fig.8**). This was interpreted as the transition from the Matuyama Epoch (R) to the Brunhes Chron (N) which polarity reversal is now dated at 773 +/- 5 ka (Haneda et al., 2020). The magnetism of the older samples from the pits below the Reversed level was equivocal and so the reversal could be the result of short-lived magnetic excursions during the late Pleistocene as interpreted by magnetostratigraphic studies at Khok Sung - also in Nakhon Ratchasima (Khorat) Province (Suraprasit et al., 2015). However, the discovery of an in situ, rare, un-abraded, delicate and hollow sphere, and definitely unworked

Australasian tektite, at the reversal level (**Fig.8**) strongly supports Sangad's and Peter Haines et al's (2004) interpretation. The stratigraphic situation in the Khorat sandpits is further complicated by the detailed palynological work of Yang and Grote (2018 a, b) who reported thermoluminescence dating on 7 samples and two C14 dates ranging from 172,739 +/- 22,400 a BP to 27,332 +/- 3000 a BP in a 3.25 m section from about 13.5 m to 10.0 m depths which covers the same depth interval as the study by Haines et al., (2004). Yang and Grote (2018 b) suggest major vegetational and palaeoclimatic changes from cool to warm to cool, and correlation to parts of two glacial periods (Riss and Wurm) during the deposition of this 3.25m of sediments.

Whether or not Sangad (and Howard, 1999, 2011a, b; Howard et al., 2003 and Haines et al., 2004) were correct in ascribing some parts of the upper sections of the sandpits and their contained biota to the 788 ka Australasian impact event, the importance of the studies in the Khorat sandpits to a wide variety of earth, planetary and biological sciences suggest that more efforts should be made on dating studies along with palaeontological, sedimentological and palynological studies in the sandpits. The pits are ephemeral and most, if not all, will end up filled with water. More careful interdisciplinary studies such as at Khok Sung by Prof. Jaroon Duangkrayom et al., (2014) are needed. If Sangad was correct, then northeastern Thailand contains globally unique evidence of the catastrophic

environmental effects of a major impact event. This is, of course, of global scientific interest but is also important as a possible tourist attraction that will bring employment to the local community.

With characteristic intellectual exuberance Sangad then extended his hypothesis to include events far from Khorat. He included the demise of Miocene vertebrates throughout Thailand (including Khorat) in his model which is highly unlikely. However, his speculation should prompt all who work in Khorat to carefully demarcate the Miocene biota from the Pleistocene and, if possible, the early from the later Pleistocene faunas and floras. He further suggested that the well-known, almost certainly, Miocene, fish beds in Phetchabun Province and shell beds near Bangkok, most likely to be Holocene, were also the result of the impact. These speculations are possible but unlikely on present evidence.

He further extended his hypothesis to the extinction of Miocene faunas in Australia. Although the widespread occurrence of unworked tektites is a very useful horizon in Australian Quaternary stratigraphy and the 788 ka event may well have had environmental and biotic consequences on that huge continent, any correlation with Quaternary megafaunal extinctions in Australia is very unlikely.






Fig. 8 Stratigraphic sections through the 22 m of the Ban Tha Chang and Chum Phuang sandpits in Nakhon Ratchasima (Khorat) Province showing Normal (N=Brunhes) and Reversed (R= Matuyama) magnetostratigraphy. Unabraded, hollow sphere, tektite in Chum Phuang section shown as T_n and reworked abraded tektites as T_a. From Haines et al., (2004). Colour slide kindly supplied by Dr Peter Haines (Australia).

However, his suggestion that our hominid ancestor *Homo erectus* witnessed and probably suffered from the tektite fall is well supported by studies in Java and in Yunnan (Yamei et al., 2002). The effects throughout SE Asia of the 788 ka 'cataclysm' on *Homo erectus* are recently well documented by Li et al., (2021) and provide considerable support for another of Sangad's 'wild' ideas.

Sangad suggested that the eroded pillars known as sao din (earth pillars) (**Fig.9**) found from northern Thailand to Sra Kaeo include deposits of loess caused by the 788 ka impact. These deposits remain undated directly but archaeological excavations in the Nanoi Basin, south of Nan in northern Thailand have found stone tools very similar to those in the Chinese Bose Basin closely associated with tektites as discussed by Li et al., (2021) and Yamei et al., (2002). Zeitoun et al., (2012) suggest an age of 1 to 0.5 Ma for the sao din stone tools so perhaps this is another of Sangad's 'wild ideas' that will soon be proven correct.




Fig. 9 'Sao Din' earth pillars at Nanoi, south of Nan city northern Thailand.

Sangad also claimed that the 788 ka impact in Indochina led to epeirogenesis in northern Thailand which he named the Inthanon Uplift. However, the uplifted deposits in northern Thailand (see Bunopas, 2021, this volume) are not well dated and this idea must, on present knowledge, be labelled as speculative. However, the huge Indochina impact or impacts must have caused severe ground shaking and the writer and others have observed faults in the Pleistocene

sections of northeastern Thailand that may be due to this event. The tectonic effects of this cataclysmic impact event should certainly be considered.

Almost a hundred years ago the famous Australian geologist Sir Edgeworth David and colleagues noted that several tektite-like glasses plotted on a great circle band around the earth and suggested an extraterrestrial origin (David et al., 1927). On the basis of extensive geochemical studies all meteoritic workers now accept that tektites are molten target rocks due to impacts by either asteroids or comets. Ramsay Ford (1988) suggested a similar path from SE Asia to Tasmania for the Australasian tektites. A date of 816 +/- 7 ka for the impact glass known as Darwin Glass scattered around Darwin Crater in western Tasmania (Howard, 2004) almost overlaps (to within 18 ka) with the best date for Australasian tektites (788 +/- 3 ka) and, with dates for the Zhamanshin crater in Kazakhstan (about 920-650 ka) led to the great circle idea being resurrected by several authors. Influenced by the work of Professor John Wasson (UCLA), Sangad postulated a cometary impact. In 1994, Sangad was strongly impressed by the well documented impact of huge fragments of the Shoemaker-Levy comet on Jupiter. Sangad envisaged a similar mechanism for the possibly simultaneous production of the Zhamanshin, Indochina and Darwin craters, their impact glasses and strewn fields. He envisaged an initial impact of a comet fragment in Kazakhstan, followed by an impact in Indochina and then at Mount Darwin, Tasmania (Bunopas, 2021, this volume). This scenario is possible and the Shoemaker-Levy comet break-up and massive impacts into Jupiter's atmosphere substantially increases the plausibility of Sangad's hypothesis. However, general acceptance of the idea is delayed because of the at least 18,000 year difference between the Darwin Crater and Australasian tektites and the as yet poorly dated glasses and crater at Zhamanshin.

The recent announcement of the discovery of a possible Australasian tektite source crater beneath Quaternary basalt of the Bolaven Plateau in Laos would, if correct, be the final crater to be identified and associated with a major tektite strewn field (Sieh et al., 2020). The crater is in the most likely position within the strewn field based on tektite morphology, microtektite and tektite geochemistry and sandstone target rock age and chemistry (Ford, 1988; Howard, 2011a, b). Our work on the same outcrops as Professor

Sieh and colleagues does not support their conclusions but if Sieh et al., (2019) are found to be correct then their small 15 km diameter crater would probably support the late Professor John Wasson's hypothesis of multiple small impact craters rather than one major Indochina crater (Wasson, 1991). If correct, then the Bolaven Crater impact was of sufficient size to account for the events summarised by Sangad in northeast Thailand but possibly not large enough to account for the whole Australasian strewn field.

5. MENTORING

One of Sangad's greatest contributions was to encourage and support both the study and research of Thai geologists at overseas universities and foreign geologists both senior and junior to carry on research in Thailand. He mentored, helped and encouraged DMR geologists Drs Thanis Wongwanich and Pol Chaodumrong to pursue PhD studies at the University of Tasmania in Australia and Somkiet Maranate and Worakul (Vorakul) Kaewayna to complete MSc degrees respectively on palaeomagnetism and Quaternary sedimentology at Victoria University in New Zealand.

He helped and encouraged Clive Burrett, Steve Carey, Ramsay Ford, Peter Haines, John Long and John Shergold (from Australia), John Wasson and Gudron Wasson (from USA) and Shigeki Hada (from Japan) and initially Father Henri Fontaine (from France) on a wide variety of fruitful projects all of which led to many publications in the international scientific literature.

Sangad helped students from Tasmania who studied the Permian glacial deposits in Phuket (John Hills) and the Lower Palaeozoic on Ko Tarutao (Tim Akerman and Roger Mason) and Kieren Howard who studied the fluvial deposits in Khorat.

6. CONCLUSIONS

Had Sangad retired without publishing his work on tectonics and impacts he would be remembered as a helpful, inspiring, pioneering, widely read, highly productive and influential geologist and as the father of Thai stratigraphy. By using his unrivalled and hard-won knowledge of Thai geology to construct, tectonic subdivisions of Thailand and an original and provocative tectonic model of South East Asia he placed Thai geology on a global stage, mentored generations of young Thai geologists and induced many

foreign scientists to carry-out collaborative research in the kingdom.

Like the pioneer of continental drift theory, Alfred Wegener, Sangad extended his impact theory too far, making outrageous claims that could be easily criticized often leading to the dismissal of all of his ideas. Although some of Sangad's extreme ideas on the effects of the 788 ka impact will be consigned to the waste basket of disproved hypotheses (joining some of Wegener's) many of his ideas on the Quaternary have gained support and are being currently cited and favourably discussed (e.g. Li et al., 2021; Zak et al., 2019).

Sangad had and continues to have a huge influence on Thai geoscience and with his seminal work on tectonics and cometary collisions continues to have a big impact on international earth and planetary science.

ACKNOWLEDGEMENTS

Dr Sangad welcomed me to Thailand and inducted me into Thai geology. He generously imparted his vast knowledge of Thai geology and later meteoritics and encouraged me with characteristic quiet humour and patience. His DMR teams particularly Dr Thanis Wongwanich, Dr Pol Chaodumrong and the late Dr Somboon Khositantont and Mr Charoen Butrnuang supported me in numerous ways. Professor Che Aziz Ali, Professor Kieren Howard, Dr Apsorn Sardud and Dr Pol Chaodumrong kindly read the manuscript and made many useful comments. Dr Peter Haines, Professor Kieren Howard, Dr Pol Chaodumrong and the Bunopas family helped with figures and photographs

REFERENCES

- Barnes, V. E. 1971. Description and origin of 12.8-kg layered tektite from Thailand (abstract). *Meteoritics*. 6 (4), 249.
- Barnes, V.E. & Pitakpaivan, K. 1962. Origin of Indochinite tektites. *Proc. Nat. Acad. Sci. USA*. 48, 947-955.
- Bunopas, S. 1961. Report on the geology and mineral deposits at Huai Tad-Bang Kud, Chiangmai. DMR Bangkok.
- Bunopas, S. 1962. Report on investigations of lignite and oil shale deposits, Li, Lamphun. Department of Mineral Resources, Bangkok.
- Bunopas, S. 1967. Report on the investigation of geology of Amphoe Song – Rong Khwang, Phrae Province, Department of Mineral Resources, Bangkok.
- Bunopas, S. 1969. Report on the geology of Amphoe Tha Pla-Nam Pat and Pha Som Dam. Department of Mineral Resources, Bangkok.
- Bunopas, S. 1966. Upper Miocene – Pleistocene Stratigraphy and the Paleontology of Gladstone, Wararapa New Zealand. Unpublished M.Sc.Thesis, Geology Department, Victoria University of Wellington, New Zealand.
- Bunopas, S. 1971a. On the geology and stratigraphy of the Nam Phrom Dam and its vicinities. *Geol. Soc. Thailand Newsletter*, 4, 17-40.
- Bunopas, S. 1971b. Jurassic beach sandstone. *Mineral Resources Gazette*. 16 (1), 40-48.
- Bunopas, S. 1972a. Preliminary report on the geological survey of Phitsanalok Quadrangle, 1:250,000 map. Department of Mineral Resources, Bangkok.
- Bunopas, S. 1972b. Brief note on the Phitsanalok Sheet. *Geological Society of Thailand Newsletter*. 4, (1-3), 17-40.
- Bunopas, S. 1973. On the tin deposit in Rayong Bay, eastern gulf of Thailand. *Geological Society of Thailand Newsletter*. 6 (5-6), 120-128.
- Bunopas, S. 1974 a. On the Lower Palaeozoic rocks in Thailand. *Geological Society of Thailand Newsletter*. 7 (3), 61-83.
- Bunopas, S. 1974 b. Geological map of Changwat Phitsanalok, 1:250,000. Department of Mineral Resources, Bangkok.
- Bunopas, S. 1975 a. On the Carboniferous volcanic sequence and red beds at Khao Luang, Sukhothai. Dept. Geol. Sci. University Chiangmai. Special Paper 1. Proceedings of the Conference on the Geology of Thailand, 1973. 2, 39-40.
- Bunopas, S. 1975 b. Permo-Triassic rocks in the Lansang Fault Zone. Dept. Geol. Sci. University Chiangmai. Special Paper 1. Proceedings of the Conference on the Geology of Thailand, 1973. 1, 39-40
- Bunopas, S. 1976 a. On the stratigraphic successions in Thailand- A preliminary summary. *Journal. Geol. Soc. Thailand*. 2, 31-58.
- Bunopas, S. 1976 b. Geology and Mineral Resources of Phitsanalok. Department of Mineral Resources, Bangkok. Report on Investigations. 16 Parts 1 and 2, 1-217, 30 plates.
- Bunopas, S. 1976 c Geological map of Nakhon Sawan, 1:250,000 series. Department of Mineral Resources, Bangkok.
- Bunopas, S. 1976 d Geological map of Suphanburi, 1:250,000 series. Department of Mineral Resources, Bangkok.
- Bunopas, S. 1980 Geology and mineral resources of Changwat Nakhon Sawan. Department of Mineral Resources, Geological Survey Report 1. Bangkok.
- Bunopas, S. 1985. Review on tectonic events of northeastern Thailand (abstract). *Conf. Geol. Min. Res. Dev. NE Thailand*, Khon Kaen University.
- Bunopas, S. 1980. Geology and Mineral Resources of Changwat Nakhon Sawan Quadrangle, Department of Mineral Resources, Geological Survey Report 1.
- Bunopas, S. 1981. Paleogeographic history of western Thailand and adjacent parts of Southeast Asia - a plate tectonics interpretation, Ph.D Thesis, 810 pp., University of Wellington, New Zealand, 1981. 810 pp., 164 figs., 21 tables.
- Bunopas, S. 1982. Palaeogeographic history of Western Thailand and adjacent parts of Southeast Asia. Geological Survey of Thailand Paper 5, Royal Thai Department of Mineral Resources, Thailand. 1-810 pp 164 figs., 21 tables.
- Bunopas, S. 1983. Paleozoic succession in Thailand. In: Nutalaya, P., ed., Proceedings of the Workshop on Stratigraphic Correlation of Thailand and Malaysia (Vol. 1). Haad Yai, Thailand. 39–76.
- Bunopas, S. 1989. Tektites-Their origin and the continental catastrophic destruction in NE Thailand and Indochina. Abstract. Pacific Science Congress, Seoul, Korea.
- Bunopas, S. 1990. Tektites-Their origin and the continental catastrophic destruction in NE Thailand and Indochina. Proc. of the 16th Conf. on Sciences and Technology of Thailand, 25-27 October, 1990, Bangkok. p. 512-513.
- Bunopas, S. 1992a. Tektites - Their origin and the Early Quaternary catastrophic destruction in Northeastern Thailand and south Indochina. Proc. 29th Intern. Geol.

- Congress. 24 Aug.- 3 Sept. 1992. Kyoto, Japan, Abstracts Vol. 3 (3), II-12-4 0-8 2114, p. 650.
- Bunopas, S. 1992b. Regional Stratigraphic Correlation in Thailand. In: Piancharoen, C., ed., Proceedings of the National Conference on Geological Resources of Thailand: Potential for Future Development. Department of Mineral Resources. Bangkok, Thailand: 189-208.
- Bunopas, S. 1994. The Regional Stratigraphy, Paleogeographic and tectonic events of Thailand and Continental Southeast Asia. In: Angsuwathana, P., Wongwanich, T., Tansathien, W., et al., eds., Proceeding of the International Symposium on Stratigraphic Correlation of Southeast Asia, 2-24
- Bunopas, S., & Bunjitradulya, S. 1975. Geology of Amphoe Bo Phloi, north Kanchanaburi, with special notes on the 'Kanchanaburi Series'. Journal of the Geological Society of Thailand 1, 51-67.
- Bunopas, S., Burrett, C. & Hada, S. 1990. Stratigraphy and paleogeographic evolution of the upper Paleozoic- early Mesozoic continental collision in Thailand (abstract). Proc. Dev. Geol. Thailand year 2000, Chulalongkorn University, Bangkok.
- Bunopas, S., Fontaine, H., Salyapongse, S. & Vachard, D. 1983. Permian paleogeography in southeast Thailand evidenced by new discoveries. Journal Geological Society of Thailand. 6, 17-21.
- Bunopas, S., & Khositantont, S. 1999. Geology and National Development in accordance with the King's new theory for self-sufficient agriculture as viewed from the Buntharik event (in Thai). In: The Special Memorial Book for the King's 72 Birthday in 2542 (1999), contribution made to the King, "The Geographer of The Nation" by The Rajabhat Institute Buriram: 3 pp.
- Bunopas, S., Khositantont, S., & Wasson, J. T. 1997. Evidence of the Early Global Disaster and Destruction from Extraterrestrial Impacts Comet in Northeast Thailand and South Indochina within the Australian Tektites Field: The last Mass Extinction, (extended abstract). Proc. of the national conf. on stratigraphy and tectonics evolution of Southeast Asia and the South Pacific. 19-24 August. 1997, Bangkok, Thailand, 434-435.
- Bunopas, S., Muenlek, S., & Tansuwan, V. 1978. Geology of Tarutao Island. Journal of the Geological Society of Thailand. 6, 121-138.
- Bunopas, S., Pitakpaivan, K., & Sukroo, J. 1978. Preliminary palaeomagnetic results from Thailand sedimentary rocks. In: Proceedings of the 3rd Regional Conference on Geology and Mineral Resources of Southeast Asia, Nutalaya, P. (ed.). Bangkok. 24-32.
- Bunopas, S., & Vella, P. 1978. Late Palaeozoic and Mesozoic structural evolution of northern Thailand: a plate tectonic model. In: Proceedings of the 3rd Regional Conference on Geology and Mineral Resources of Southeast Asia, Nutalaya, P. (ed.). Bangkok, 133-140.
- Bunopas, S., & Vella, P. 1983a. Tectonic and geologic evolution of Thailand. In: Nutalaya, P., (ed.) Proceedings Of the Workshop on stratigraphic correlation of Thailand and Malaysia, Haad Yai, Thailand, 8-10 September 1983, Tech. Paper 1, 307-323.
- Bunopas, S., & Vella, P. 1983b. Opening of the Gulf of Thailand - rifting of continental Southeast Asia, and late Cenozoic Tectonics, Journal of the Geological Society of Thailand, 6, 1-12.
- Bunopas, S., & Vella, P. 1984. Phuket-Kaeng Krachan Group, A rifted continental margin deposit with effect of ice-rafting from Gondwana. Abstr. paper presented at GEOSEA V., 9-13 April, Kuala Lumpur.
- Bunopas, S., & Vella, P. 1989. "Pebble mudstones", A composited assemblage of ice-rafted sediments on the western rifted marginal basin of Shan-Thai while drifting away from Gondwana during the Upper Paleozoic. Proceedings of the 4th Intern. Symp. On Pre-Jurassic East Asia, IGCP Project 224. Hat Yai and Langkawi Island, Thailand and Malaysia, 24 April-2 May 1989.
- Bunopas, S., Wasson, J. T., Vella, P., Fontaine, H., Hada, S., Burrett, C., Suphajanya, Th. & Khositantont, S. 1999a. Catastrophic loess, mass mortality and forest fires suggest that a Pleistocene cometary impact in Thailand caused the Australasian tektite field. Journ. Geol. Soc. of Thailand. 1999 (1), 142-151.
- Bunopas, S., Wasson, J. T., Vella, P., Fontaine, H., Hada, S., Burrett, C., Suphajanya, Th. & Khositantont, S. 1999c. Ancient analogs of burial alive extinction of the mastodons in catastroloess in Thailand, and of the last dinosaurs (in eggs) in Gobi Desert: further on tektites. Symposium on Mineral, Energy and Water Resources of Thailand: Towards the year 2000, October 28-29 1999, Bangkok, Thailand: 168-177.
- Bunopas, S., Wasson, J. T., Vella, P., Fontaine, H., Hada, S., Burrett, C. & Khositantont, S., 1998. The early extraterrestrial collision with a whole comet in the Australasian tektite field, apparent evidences and discoveries from Thailand and East Asia (abstract). Geosea '98. Ninth Regional Congress on Geology, Mineral and Energy Resources of Southeast Asia. 17-19 August 1998, Shangri-La Hotel, Kuala Lumpur, Malaysia. Programme and Abstracts. 191-192.
- Bunopas, S., Wasson, J. T., Vella, P., Fontaine, H.; Hada, S., Burrett, C., Suphajanya, Th. & Khositantont, S. 1999a. Catastrophic loess, mass mortality and forest fires suggest that a Pleistocene cometary impact in Thailand caused the Australasian tektite field. Journ. Geol. Soc. of Thailand 1999 (1): 1-17.
- Bunopas, S., Yaemniyom, Ng. & Khositantont, S. 1999b: Catastroloess and its derivatives, the life time sustainable construction sands was originally high atmospheric settling in the Quaternary cometary impact in Thailand and Asia. Symposium on Mineral, Energy and Water Resources of Thailand: Towards the year 2000.
- Chaimanee, Yaovaluk, 2004. Fossils from Khok Soong and Ban Tha Chang, Khorat. In: Numerous fossils in Paleontology of Thailand, in 114 years of the Department of Mineral Resources. (in Thai)
- Chaimanee, Yaovaluk, 2007. Diversity of Cenozoic Mammals in Thailand: Paleoenvironment and Age Updated. (GEOTHAI'07) International Conference on Geology of Thailand: Towards Sustainable Development and Sufficiency Economy. Geothai 2007, pp. 73-79. 21-22 November. Imeral Hotel, Bangkok, Thailand.
- Chapman, D. R., & Larson, H. K. 1963. On the Lunar origin of tektites. J. Geophys. Res. 68, 4305 - 4358.
- Charusiri P, Daorerk V, Krowchan V, Klongsara N, Kosuwan S, Sriattanachatchawan V, & Santatiwongchai U. 2002. Quaternary tektites and their sediment hosts at Ban Tachang Sand Pit, Chaloeam Prakiat, Nakhon Ratchasima, NE Thailand: Stratigraphy and TL Ages. In: Mantajit N, et al. editors. The Symposium on Geology of Thailand. 2002 August 26-31; Bangkok, Thailand: Department of Mineral Resources, Bangkok; p. 178-180.
- David T. W. E., Summers H. S. & Ampt G. A. 1927. The Tasmanian tektite - Darwin glass. Proceedings of the Royal Society of Victoria. 39, 167- 190.
- Duangkrayom J, Ratanasthien B, Jintasakul P, & Carling P.A. 2014. Sedimentary facies and paleoenvironment of a Pleistocene fossil site in Nakhon Ratchasima province, northeastern Thailand. Quaternary International. 325, 220-238. doi: 10.1016/j.quaint.2013.07.048
- Duangkrayom, J., Wang, S.-Q., Deng, T., & Jintasakul, P. 2017. The first Neogene record of Zygodolopodon (Mammalia, Proboscidea) in Thailand: implications for the mammutid evolution and dispersal in Southeast Asia. Journal of Paleontology. 91 (1), 179-193.

- Fiske P. S., Putthapiban P., & Wasson J. T. 1996. Excavation and analysis of layered tektites from northeast Thailand: Results of 1994 field expedition. *Meteoritics & Planetary Science*. 31, 36–41
- Ford, R., J. 1988: An empirical model for the Australasian tektite field. *Australian Journal of Earth Science*. 35, 483-490.
- Haines, P. W., Howard, K.T., Ali, J.R., Burrett, C., & Bunopas, S. 2004. Flood deposits penecontemporaneous with ~0.8 Ma tektite fall in NE Thailand: impact-induced environmental effects? *Earth and Planetary Science Letters*. 225, 19-28.
- Haneda, Y., Okada, M., & Kitamura, T. 2020. A full sequence of the Matuyama-Brunhes reversal in the Chiba composite section, Central Japan. *Progress in Earth and Planetary Sciences*. 7 no.44, doi.org/10.1186/s40645-020-00354-y
- Howard, K. T. 1999. Distal environmental effects of the 770ka impact event. Unpublished Hons. Thesis at the University of Tasmania, Australia. 175 pp.
- Howard, K. T. 2004. Origin of Darwin Glass. Unpublished Ph. D. Thesis at the University of Tasmania, Australia. 362 pp.
- Howard K. T. 2011a. Volatile enhanced dispersal of high velocity impact melts and the origin of tektites. *Proceedings of the Geologists Association*. 122, 363–382.
- Howard, K.T. 2011b. Tektites in: Ridd, M.F., Barber, A.J., Crow, M.J. (Eds.), *The Geology of Thailand*, The Geological Society of London, pp. 574-591. doi.org/10.1144/GOTH.21
- Howard, K. T., Haines, P.W., Burrett, C., Ali, J., & Bunopas, S. 2003. Sedimentology of 0.8 Ma log-bearing flood deposits in northeast Thailand and mechanisms for pre-flood deforestation, *Proceedings, 8th International Congress on Pacific Neogene Stratigraphy*, Chiang Mai, Thailand, 2003, pp. 49-67.
- Lauprasert, K., Watchajittaphan, P., Juangnam, S., & Bhuttarach, S. 2019. Freshwater crocodile, *Crocodylus siamensis* Schneider, 1801, from the Middle Pleistocene deposits in Chaloe Phrakiat District, Nakhon Ratchasima, Thailand. *Annales de Paléontologie*. 105 (4), 269-274. doi:10.1016/j.annpal.2019.05.002
- Li, H., Lotter, M., Kuman, K., Lei, L., & Wang, W. 2021. Population dynamics during the Acheulian at 0.8 Ma in East and SE Asia: Considering the influence of two geological cataclysms. *Palaeogeography, Palaeoclimatology, Palaeoecology* 562, https://doi.org/10.1016/j.palaeo.2020.109927. (online-in press)
- Naksri, W., Tong, H., Lauprasert, K., Jintasakul, P., Suteethorn, V., Vidthayanon, C., & Claude, J. 2019. Plio-Pleistocene giant tortoises from Tha Chang sandpits, Nakhon Ratchasima Province, Thailand. *Annales de Paléontologie*. 105 (4), 257-267. doi:10.1016/j.annpal.2019.06.001
- Nutalaya, P., Vella, P., Bunopas, S., & Kaewyana, V., 1986. Quaternary processes in Thailand. In: Thiramongkol (ed.), *Proceeding of the Workshop on Economic, Geology, Tectonics Sedimentary Processes and Environmental of the Quaternary of Southeast Asia*. p. 23-33, Bangkok.
- Nichol, J.E., & Nichol, D. W. 2015. Character and provenance of aeolian sediments in northeast Thailand. *Aeolian Research*. 19 (A), 5-14. <https://doi.org/10.1016/j.aeolia.2015.09.001>
- O'Keefe, J. A., 1976. *Tektites and Their Origin*. Elsevier. Amsterdam. 354 pp.
- Raksaskulwang, L., & Bunopas, S. 1985. Stratigraphy and tectonic evolution in the Mae Hong Son Province, north western Thailand. Unpublished report of the Royal Thai Department of Mineral Resources. 1-39.
- Sato, Y. 2002. Preliminary report on the occurrence of fossil mammals in Nakhon Ratchasima, northeast Thailand, in: N. Mantajit (Ed.). *Proceedings of the Symposium on Geology of Thailand*, Department of Mineral Resources. Bangkok, 2002, pp. 230-232.
- Sibava, V. 1993. Quaternary Sequences in Southeast Asia and their regional correlation. In: *ESCAP Atlas of Stratigraphy XII., Quaternary Stratigraphy of Asia and the Pacific*, IGCP 296. Mineral Resources Development Series, No 62. United Nations. pp. 1 -29.
- Sieh, S., Herrin, J., Jicha, B., Schonwalder, D., Moore, J., Banerjee, P., Wiwegwin, W., Sihavong, V., Singer, B., Chualaowanich T., & Charusiri, P. 2020. Australasian impact crater buried under the Bolaven volcanic field, Southern Laos. *Proceedings of the National Academy of Science*. 117 (3), 1346-1353. <https://doi.org/10.1073/pnas.1904368116>
- Vella, P., Bunopas, S., & Kaewyana, W. 1987. A loess covered alluvial gravel terrace near Chiangmai, Northern Thailand. *Geological Society of Thailand Newsletter*. 1987 (3), 29-31.
- Wasson J. T. 1991. Layered tektites: A multiple impact origin for the Australasian tektites. *Earth and Planetary Science Letters* 102, 95–109.
- Yamei, H., Potts, R., Baoyin, Y., Zhengtang, G., Deino, A., Wang, W., Clark, J., Guangmao, X., & Weiwen, H., 2000. Mid-Pleistocene Acheulean-Like Stone Technology of the Bose Basin, South China. *Science*, 287, 1622-1626.
- Yang, F-C., & Grote, P.J. 2018 a. Variation of fluorescence intensity in Pleistocene pollen and the correlation of fluvial deposits in the Khorat Plateau, Thailand, *Palynology*. 42 (1), 28-35, DOI: [10.1080/01916122.2017.1303796](https://doi.org/10.1080/01916122.2017.1303796)
- Yang, F-C., & Grote, P.J. 2018 b. Riverine vegetation and environments of a Late Pleistocene river terrace, Khorat Plateau, Southeast Asia. *Palynology*. 42 (2) 158-167. doi: 10.1080/01916122.2017.1296044
- Žák, K., Skála, R., Pack, A., Ackerman, L., & Křížová, Š. 2019. Triple oxygen isotope composition of Australasian tektites. *Meteoritics & Planetary Science*. 54 (5), 1167–1181. doi:10.1111/maps.13251
- Zeitoun, V., Forestier, H., Auetrakulvit, P., Khaokhiew, C., Rasse, M., Davtian, G., Winayalai, C., Tiamtinkrit, C. 2012 Discovery of a prehistoric site at Sao Din (Nanoi, Nan province, Northern Thailand): Stone tools and new geological insights. *Découverte d'un site préhistorique à Sao Din (Nanoi, province de Nan, Nord de la Thaïlande): études lithique et géologique préliminaires*. *Comptes Rendus Palevol*. 11, 575-580. doi:10.1016/j.crpv.2012.06.007