

TABLE OF CONTENTS

	Page
TABLE OF CONTENTS	i
LIST OF TABLES	iii
LIST OF FIGURES	iv
LIST OF ABBREVIATIONS	v
INTRODUCTION	1
LITERATURE REVIEWS	3
Chlorophyll mutation	3
Plant characterization from chlorophyll	4
Plant characterization from chlorophyll fluorescence	4
Seed growth and development	7
Seed desiccation	9
DNA markers	10
DNA markers based on hybridization	11
DNA markers based on PCR	11
Bulk segregant analysis for mapping or tagging gene	14
Linkage map	15
MATERIALS AND METHODS	16
Plant materials	16
Determination of chlorophyll content	16
Determination of chlorophyll fluorescence	17
Seed weight and seed moisture content	18
Data analysis	18
Seed cell morphology	19
DNA extraction	19
SSR analysis	20
ISSR analysis	20
AFLP analysis	21
Gel electrophoresis and detection	22
Analysis of genetic distance	23

TABLE OF CONTENTS (Continued)

	Page
RESULTS	24
Determination of chlorophyll content	24
Determination of chlorophyll fluorescence	26
Seed growth and development	28
Seed cell morphology	35
Inheritance of opaque leaf and linkage test	41
Molecular markers linked to opaque leaf	43
DISCUSSION	46
Determination of chlorophyll content	46
Determination of chlorophyll fluorescence	47
Seed growth and development	48
Seed cell morphology	49
Inheritance of opaque leaf and linkage test	50
Molecular markers linked to opaque leaf	51
CONCLUSION	52
LIERATURE CITED	54
APPENDIX	70

TABLE OF CONTENTS

Table	Page
1 Chlorophyll fluorescence in F_v/F_m and F'_v/F'_m values of normal leaf and mutant leaf genotypes.	27
2 Chlorophyll fluorescence in NPQ value of normal leaf and mutant leaf genotypes.	27
3 Number of F_2 plants segregating for opaque leaf, petiole color, and growth habit.	42
4 Linkage test between two traits in F_2 plants.	43
 Appendix Table	
1 Chlorophyll a, b, total (a+b) contents (g/m^2) and a/b ratio.	71
2 Pod fresh weight (g/pod) of normal and opaque mungbean.	72
3 Pod wall fresh weight (g/pod) of normal and opaque mungbean.	72
4 Seeds fresh weight (g/pod) of normal and opaque mungbean.	73
5 Seed coats fresh weight (g/pod) of normal and opaque mungbean.	73
6 Cotyledons fresh weight (g/pod) of normal and opaque mungbean.	74
7 Pod dry weight (g/pod) of normal and opaque mungbean.	74
8 Pod wall dry weight (g/pod) of normal and opaque mungbean.	75
9 Seed dry weight (g/pod) of normal and opaque mungbean.	75
10 Seed coat dry weight (g/pod) of normal and opaque mungbean.	76
11 Cotyledon dry weight (g/pod) of normal and opaque mungbean.	76
12 SSR primers used for parent screening.	77
13 ISSR primers used for parent screening.	80
14 Nucleotide sequences of adapter and primers used for AFLP analysis.	81
15 AFLP combination primers used for PCRI and PCRII reactions.	82
16 Ninety-six AFLP combination primers (<i>EcoRI</i> / <i>MseI</i>) obtained from parental screening.	83

FIGURE OF CONTENTS

Figure	Page
1 Chlorophyll a, b, total (a+b) contents (g/m ²), and a/b ratio in opaque leaf, Opaque, OP531, and OP541, and normal leaf plant genotypes, Berken, N531, and N541.	25
2 Accumulation of pod fresh weight (a) and dry weight (b) of opaque vs. normal leaf plants at 3 to 18 days after flowering.	30
3 Accumulation of pod wall fresh weight (a) and dry weight (b) of opaque vs. normal leaf plants at 3 to 18 days after flowering.	31
4 Accumulation of seed fresh weight per pod (a) and dry weight per pod (b) of opaque vs. normal leaf plants at 3 to 18 days after flowering.	32
5 Accumulation of seed coat fresh weight per pod (a) and dry weight per pod (b) of opaque vs. normal leaf plants at 3 to 18 days after flowering.	33
6 Accumulation of cotyledons fresh weight per pod (a) and dry weight per pod (b) of opaque vs. normal leaf plants at 3 to 18 days after flowering.	34
7 Seed water content per pod of opaque vs. normal leaf plants at 3 to 18 days after flowering.	35
8 Photographs from light microscope of pod and seed cross section from opaque leaf plant at 3, 6, 9, 12, 13, and 14 days after flowering.	37
9 Photographs from light microscopes of pod and seed cross sections from normal leaf plant at 3, 6, 9, 12, 15, and 17 days after flowering.	38
10 Photographs from light microscopes of cotyledonary transfer cells from opaque and normal leaf plant.	39
11 Photographs from light microscopes of cotyledon cross sections from opaque and normal leaf plant.	40
12 Photograph of plants with opaque leaf (O) vs. normal leaf (N) at seedling growth stage.	42
13 A partial linkage map between AFLP markers and <i>op</i> gene.	45

LIST OF ABBREVIATIONS

cM	=	centi-Morgan
NPQ	=	non-photochemical quenching
F_v/F_m	=	maximum quantum efficiency of PSII photochemistry
F'_v/F'_m	=	PSII maximum efficiency
RFLP	=	restriction fragment length polymorphism
PCR	=	polymerase chain reaction
RAPD	=	random amplified polymorphic DNA
ISSR	=	inter simple sequence repeat
SSRs	=	microsatellite or simple sequence repeats
AFLP	=	amplified fragment length polymorphism