

Evaluation of Atmospheric PM₁₀ in the Southwest Region of Nigeria

Mohammed Mohammed Ndamitso¹, Yahaya Ahmed Iyaka¹, Aishat Abdulkadir² and Francis Olawale Abulude^{1,3*}

¹Department of Chemistry, Federal University of Technology, Minna, Niger State, Nigeria

²Department of Geography, Federal University of Technology, Minna, Niger State, Nigeria

³Science and Education Development Institute, Akure, Ondo State, Nigeria

Received: 22 March 2020, Revised: 10 August 2020, Accepted: 8 September 2020

Abstract

Particulate matter (PM), according to World Health Organization (WHO), has caused several millions of deaths, and both the young and old have been affected. To mitigate this problem, the compositions of the elements that make up the PM and their sources must be determined. These can provide the necessary information needed by stakeholders to work on. To this end, the study determined the mass concentrations of the PM₁₀ obtained from the Federal University of Technology, Akure (FUTA), National Museum and Monuments and Oba-Ile; for a period of ten months (January-October, 2018). A total of thirty samples was collected. The PM concentration was calculated gravimetrically and the data were subjected to statistical analyses. The results for PM in $\mu\text{g}/\text{m}^3$ were: FUTA (39.10 ± 0.31 - 133.22 ± 0.21), Museum (49.71 ± 0.11 - 196.70 ± 0.01), and Oba-Ile (34.50 ± 0.31 - 161.30 ± 0.42). The results were well above the WHO standard limits. Air Quality Index (AQI) was calculated for each location, and the results for the air quality in Akure showed that the locations were unhealthy for sensitive groups.

Keywords: particulate matter; AQI; anthropogenic activities; WHO; Akure; Nigeria
DOI.....

1. Introduction

Air pollution is a major global issue due to its impact on human health and property [1-4]. Due to urbanization and industrial development, environmental quality has declined throughout the globe in recent times [5, 6]. It is thought that more than 2.4 million individuals die each year globally due to PM pollution issues [4, 7, 8]. According to Ilyas *et al.* [9] and Ndamitso *et al.* [4], PM moves into the lungs when inhaled, and thus has adverse effects that ultimately cause health-related problems such as cancer, asthma, and cardiovascular and respiratory diseases.

It has been confirmed that PM in Europe is a pollutant and has surpassed the recommended normal limits [10, 11]. Schweizer *et al.* [12] mention that PM is a complex mixture of chemical

*Corresponding author: Tel: +2348034458674
Email: walefut@gmail.com

species from anthropogenic and natural emission sources and from secondary pollutants which form in the atmosphere. In particular, in developing nations, rapid urbanization and population growth are part of the causes of the world-wide PM issue. This statement was verified by Arnfield [13], who indicated that by 2025, more than 45% of the population will live in metropolitan towns around the globe. Balogun *et al.* [14] and Owoeye and Ibitoye [15] revealed that city migration could have risen to over 30 billion over the next 30 years. A town established in 1976, Akure is no exception. Its population has been estimated at 476,159 by the National Population Commission [16] since its founding, the population, business, industrial, farming and petroleum products (bitumen) have increased which may have an impact on particulate matter and other anthropogenic pollutants.

In Akure, Nigeria, several road constructions, farming, industrial activities, and heavy traffic have been on the increase during the past years. In these areas, no doubt, metals, gas particles, and dust are transported by wind or runoff waters. Often these particles can cause negative effects on man and buildings. To this effect, it is essential to characterize the distributions and concentrations of these metals and atmospheric deposits in the environment especially air and aquatic. This research evaluates the concentration of particulate matter (PM10) obtained in Akure, Ondo State, Nigeria.

2. Materials and Methods

Three different locations [National Museum and Monuments (005 11 40.2 E, 07 15 11.6 N), Obale (005 14 29.1 E, 07 16 04.4 N), and Federal University of Technology-FUTA (005 08 06.5 E, 07 18 07.6 N)] were selected for the sampling within the town (Figure 1). These locations represent commercial, residential, and educational settings respectively. Samplings took place once every month from 8 am to 4 pm (8 h) for 10 months (January to October 2018) using a "Gent" stacked filter unit. The PM10 were filtered through quartz filters (47mm) and kept in a desiccator at room temperature. By the end of the sampling period, thirty samples (ten per sampling area) were collected monthly. The filters were put in a chamber kept at 20°C, and 35% relative humidity (RH) for 48 h to stabilize the weight. The particulate mass was determined gravimetrically (the difference of filter's weight before and after sampling) using an analytical balance (0.001mg) (Model Sartorius Microbalance ME 5 Balance).

The PM mass concentration was calculated based on these parameters:

The average flow rate = averaging flow rate at the beginning and end of 8 h sampling (1)

Average PM mass concentration ($\mu\text{g}/\text{m}^3$) = Mass/total volume (2)

The data obtained from the study data were extrapolated and scaled up to 24 h mean before computing the air quality index. The AQI (Air quality index) for the particulate matter was calculated by using the equation given by Osimobi *et al.* [17].

$$AQI = \left[\frac{AQI_{max} - AQI_{min}}{PM_{max} - PM_{min}} \times (PM_{meas} - PM_{min}) \right] + AQI_{min} \quad (3)$$

Where:

PM_{meas} is the 24 h measured average concentration of particulate matter ($\mu\text{g}/\text{m}^3$),

PM_{max} is the maximum concentration AQI color category containing PM_{meas}

PM_{min} is the minimum concentration of AQI color category containing PM_{meas}

AQI_{max} is the maximum AQI value for color category corresponding to PM_{meas}

AQI_{min} is the minimum AQI value for color category corresponding to PM_{meas}

The results were statistically analyzed using Minitab 16 Statistical Software.




Figure 1. Map of the sampling points

3. Results and Discussion

Table 1 depicts a summary of the statistical results of the PM collected and analyzed in this study. The PM ranged between 39.10 and 133.20 (FUTA), 49.70 and 196.70 (Museum), and 34.50 and 161.30 (Oba-Ile) with mean values of 100.50, 125.20, and 112.80 $\mu\text{g}/\text{m}^3$ respectively. The Standard Error (SE) and Standard Deviation (SD) range were 10.9, 13.1 and 10.4 and 34.3, 41.3 and 33.0, while the Coefficient of Variation in percent (CV%) range was 34.17, 33.02, and 29.25, for FUTA, Museum, and Oba-Ile, respectively. The Skewness and Kurtosis gave -0.98, -0.01, -1.29 and -0.65, 0.68, and 3.80 for FUTA, Museum, and Oba-Ile, respectively. From Table 1, it could be deduced that the low standard error showed that the variability in the PM values was low during the sampling periods, while the CV (%) showed moderate variability. The variations could be due to the variations from the different months of sampling. The coefficients of variation of the PMs show that there was not much variation with locations, but the variance depicts high values due to the differences (variations) in the values obtained in each location as shown in the minimum and maximum values from each location.

The highest PM concentration occurred during the summer months when there was high temperature, solar radiation, low humidity and low wind speed. The most polluted months were April and October, which could have been due to the harsh weather conditions and high secondary transformations, as noted by Wang *et al.* [18]. The former could have been due to emissions from traffic and also agricultural burning during the April land as part of the preparation towards planting season, while the latter could have been due to the October harvest season [1]. The high PM values

Table 1. Summary of the statistical results of PM (particulate matter) ($\mu\text{g}/\text{m}^3$) measured during January to October 2018

	FUTA	Museum	Oba Ile
January	68.39	174.69	144.11
February	39.06	85.42	108.80
March	119.47	49.69	34.48
April	52.08	196.67	161.30
May	115.94	114.22	108.49
June	106.88	124.27	115.73
July	131.35	133.86	127.55
August	112.40	118.07	112.69
September	133.23	115.73	104.43
October	126.20	139.48	110.83
Mean	100.50	125.20	112.80
SE	10.90	13.10	10.40
SD	34.30	41.30	33.00
CV (%)	34.17	33.02	29.25
Min	39.10	49.70	34.50
Max.	133.20	196.70	161.30
Q1	64.30	107.00	107.50
Q3	127.50	148.30	131.70
Skewness	-0.98	-0.01	-1.29
Kurtosis	-0.65	0.68	3.80

obtained in the Museum area could have been a result of the accumulation of re-entrained and re-suspended dust resulting from vehicular emissions (low, medium and high) and unpaved terrain activities. The variations in the emissions, meteorological conditions, type of long-distance transport and secondary production are associated with the pollution levels in different seasons [19-21].

According to US. EPA [22], the air quality index of PM is categorized into good, moderate, unhealthy for sensitive groups, unhealthy, very healthy, and hazardous categories. The results were compared with this index and the results whose values were above 50 AQI range were grouped under the 'Moderate' and 'Unhealthy' groups while those with values less than this were classified as "Good" (Table 2). The average AQI in January through October of the three locations in Akure were 126.17, 131.34, and 138.96 (museum, Oba-Ile, and FUTA, respectively). All the locations were categorized as 'unhealthy for sensitive groups'. The health-related problem of PM is linked to particle sizes and particularly PM_{10} [21]. The PM deposits considered in this study were PM_{10} , which was found in reasonable quantities in the sampling locations of this study and formed the basis of the classifications given in this study.

The values recorded in all the locations were more than $100 \mu\text{g}/\text{m}^3$ recommended by India's Central Pollution Control Board [23], and more than three times greater than the European limit [24]. Our results (minimum-maximum values) were compared with other authors - Greilinger *et al.* [25] reported between 10 and $45 \mu\text{g}/\text{m}^3$ for Masenberg and from 20 to above $60 \mu\text{g}/\text{m}^3$ for Bockberg, Austria; Mehta [26] gave 3.19 and $152.88 \mu\text{g}/\text{m}^3$, for Ahmedabad, India; Abe and Miraglia [27] put that of São Paulo, Brazil to be 8 - $131 \mu\text{g}/\text{m}^3$, and the Eastern part of Nigeria (14 - $504 \mu\text{g}/\text{m}^3$) by Osimobi *et al.* [17]; Contini *et al.* [28] reported of South-Eastern Italy as 5.6 - $127.8 \mu\text{g}/\text{m}^3$. It was observed that the minimum values from our study were higher than those reported by

Table 2. Air quality index of the selected sampling locations

Sampling Locations	Air Quality Index Categories					
	0-50 (Good)	51-100 (Moderate)	101-150 (Unhealthy for sensitive groups)	151-200 (Unhealthy)	201-300 (Very Unhealthy)	>301 (Hazardous)
FUTA	-	-	138.94	-	-	-
Museum	-	-	126.17	-	-	-
Oba-Ile	-	-	131.34	-	-	-

these authors. It is worthy of note that the maximum results obtained by Mehta [26] for Ahmedabad, India, and Osimobi *et al.* [17] were higher than ours. Also, the maximum results reported in this study, however, were lower than $109.60 \pm 10.83 - 453.45 \pm 62.92 \mu\text{g}/\text{m}^3$ reported by Mandal *et al.* [29] for Delhi's industrial region and $584.8 \mu\text{g}/\text{m}^3$ by Jain *et al.* [30] in Delhi, India. The variations between these results and those of others could be due to the climatic conditions of the areas, the methodology used and the natural and anthropogenic activities of the different locations. Also, the differences could be due to the differences in the weather conditions (temperature, humidity, and air exchange rates) [31].

The locational contributions of the PM to the total amount obtained in the study are shown in Figure 2. It could be noted that Oba-Ile, a residential area contributed 47% to the PM as against 36% (Museum) and 17% (FUTA) in February. The reason for the high contribution by Oba-Ile could have been the result of traffic diversion to this sampling location as a result of an accident that occurred on the Akure-Owo expressway during the sampling period. This increase in anthropogenic activities might have contributed immensely to the increase observed in the PM₁₀ deposit of the said area during the said period (February). Likewise, the elevations in FUTA PM₁₀ results could have been due to the waste dumps around the vicinity and also the on-going renovation activities of a lecture theater besides the sampling area [31].

4. Conclusions

To conclude, the evaluation of PM in this study shows that the concentrations were high. The results were well above the WHO standard limits. The highest value was obtained within the Museum vicinity, which could have been due to anthropogenic (dust, soil, traffic, and commercial) activities. The results, when compared with International Air Quality Index, indicate that the PM quality in Akure is unhealthy for sensitive groups. Constant monitoring of the vicinities is recommended.




Figure 2. Distribution of PM₁₀ in the locations (January-October, 2018)

5. Acknowledgements

The authors are grateful to the curator of the National Museum Commission, Akure, the Dean of Faculty of Science, Federal University of Technology, Akure (FUTA) for the permission granted to use their premises for sample collections and Mr. Edah Sunday, Department of Geography, Federal University of Technology, Minna, Niger State, Nigeria for drawing the maps.

References

- [1] Zhang, J., Cheng, M., Ji, D., Liu, Z., Hu, B., Sun, Y. and Wang, Y.S., 2016. Characterization of submicron particles during biomass burning and coal combustion periods in Beijing, China. *Science and Total Environment*, 562, 812-821.
- [2] Tuankruea, V., Tongdeenog, P., Tangtham, N., Aungsuratana, P. and Witthawatchuetikul, P., 2014. Assessment of aerosol-cloud-rainfall interactions in Northern Thailand. *Proceedings of the International Academy of Ecology and Environmental Sciences*, 4(4), 134-147.
- [3] Zhao, L., Hu, G., Yan, Y., Yu, J., Wang, X. and Yan, Y., 2019. Source apportionment of heavy metals in urban road dust in a continental city of eastern China: Using Pb and Sr isotopes combined with multivariate statistical analysis. *Atmospheric Environment*, 201, 201-211.
- [4] Ndamitso, M.M., Abdulkadir, A. and Abulude, F.O. 2016. Source apportionment: Case studies of selected African countries. *International Journal of Advancements in Research and Technology*, 5(3), 1-25.
- [5] Wang, Q., Kwan, M., Zhou, K., Fan, J., Wang, Y. and Zhang, D., 2019. The impacts of urbanization on fine particulate matter (PM_{2.5}) concentrations: Empirical evidence from 135 countries worldwide. *Environmental Pollution*, 247, 989-998.
- [6] Latif, M.T., Azmi, S.Z., Noor, A.D.M., Ismail, A.S., Johny, Z., Idris, S., Mohamed, A.F. and Mokhtar, M., 2011. The impact of urban growth on regional air quality surrounding the Langat river basin, Malaysia. *Environmentalist*, 31, 315-324.
- [7] WHO, 2012. *World Health Report-Reducing Risk, Promoting a Healthy Life*. Geneva: World Health Organization.
- [8] Yun, G., He, Y., Jiang, Y., Dou, P. and Dai, S., 2019. PM_{2.5} Spatiotemporal Evolution and Drivers in the Yangtze River Delta between 2005 and 2015. *Atmosphere*, 10, 55, <https://doi.org/10.3390/atmos10020055>.
- [9] Ilyas, S.Z., Khattak, A.I., Nasir S.M., Qurashi T. and Durrani R., 2009. Air pollution assessment in urban areas and its impact on human health in the city of Quetta, Pakistan. *Clean Technologies and Environmental Policy*, 12(3), 291-299.
- [10] Harrison, R.M. and Yin, J., 2008. Sources and processes affecting carbonaceous aerosol in central England. *Atmospheric Environment*, 42, 1413-1423.
- [11] Farao, C., Caneparis, S., Perrino, C. and Harrison, R.M., 2014. Sources of PM in an industrial area: Comparison between receptor model results and semi empirical calculations of source contributions. *Aerosol and Air Quality Research*, 14, 1558-1572.
- [12] Schweizer, D., Cisneros, R. and Buhler, M., 2019. Coarse and fine particulate matter components of wildland fire smoke at Devils Postpile National Monument, California, USA. *Aerosol and Air Quality Research*, 19, 1463-1470.
- [13] Arnfield, A.J., 2003. Two decades of urban climate research: a review of turbulence, exchanges of energy and water, and the urban heat island. *International Journal of Climatology*, 23(1), 1-26.

- [14] Balogun, I.A., Adeyewa, D.Z., Balogun, A.A. and Morakinyo, T.E., 2011. Analysis of urban expansion and land use changes in Akure, Nigeria using RS and GIS techniques. *Journal of Geography and Regional Planning*, 4(9), 533-541.
- [15] Owoeye, J.O. and Ibitoye, O. A., 2016. Analysis of Akure Urban land use change detection from remote imagery perspective. *Urban Studies Research*, 2016, <http://dx.doi.org/10.1155/2016/4673019>
- [16] National Population Commission (NPC). 2006. *The 1963, 1991 and 2006 Population Census Reports*. Abuja: National Population Commission.
- [17] Osimobi, O.J., Yorkor, B. and Nwankwo, C.A., 2019. Evaluation of daily pollutant standard index and air quality index in a university campus in Nigeria using PM10 and PM2.5 particulate matter. *Journal of Science, Technology and Environment Informatics*, 7(2), 517-532.
- [18] Wang, L., Li, W., Sun, Y., Tao, M., Xin, J., Song, T., Li, X., Zhang, N., Ying, K. and Wang, Y., 2018. PM_{2.5} characteristics and regional transport contribution in five cities in Southern North China Plain, during 2013-2015. *Atmosphere*, 9(157), <https://doi.org/10.3390/atmos9040157>
- [19] Jia, M., Zhao, T., Cheng, X., Gong, S., Zhang, X., Tang, L., Liu, D., Wu, X., Wang, L. and Chen, Y., 2017. Inverse relations of PM_{2.5} and O₃ in air compound pollution between cold and hot seasons over an urban area of east China. *Atmosphere*, 8(3) , 59, <https://doi.org/10.3390/atmos8030059>
- [20] Abulude, F.O., Ndamitso, M.M., Iyaka, Y.A. and Abdulkadir, A., 2018. Study of particulate matter of Akure, Nigeria using a sharp-cut inertial filter combined with an impactor-a preliminary study. *Journal of Atmospheric and Solar-Terrestrial Physics*, 179, 396-404.
- [21] Liu, H., Schneider, P., Haugen, R. and Vogt, M., 2019. Performance assessment of a low-cost PM_{2.5} sensor for a near four-month period in Oslo, Norway. *Atmosphere*, 10(41), <https://doi.org/10.3390/atmos10020041>
- [22] US. EPA, 2009. *Patient Exposure and the Air Quality Index. Particle Pollution and Your Patients' Health*. [online] Available at: <https://www.epa.gov/pmcourse/patient-exposure-and-air-quality-index#what>
- [23] CPCB, 2008. *Stationary Sources Emission Profiles. Central Pollution Control Board, New Delhi, India*. [online] Available at: http://cpcb.nic.in/Stationary_Sources_Emission_Profiles.xls.
- [24] EEA (European Environment Agency), 2014. *Air Quality in Europe-2014 Report*. Copenhagen: European Environment Agency.
- [25] Greilinger, M., Zbira, J. and Kasper-Giebl, A., 2019. Desert dust contribution to PM10 loads in Styria (Southern Austria) and impact on exceedance of limit values from 2013-2018. *Applied Sciences*, 9(11), 2265, <https://doi.org/10.3390/app9112265>
- [26] Mehta, U., 2020. Statistical analysis of suspended and respirable suspended particulate matter (PM₁₀ and PM_{2.5}) concentrations in urban region of Ahmedabad, India. *International Journal of Science and Research*, 9(2), 605-609.
- [27] Abe, K.C. and Miraglia, S.G.E.K. 2016. Health impact assessment of air pollution in São Paulo, Brazil. *International Journal of Environmental Research and Public Health*, 13(7), 694, <https://doi.org/10.3390/ijerph13070694>
- [28] Contini, D., Cesari, D., Donato, A., Chirizzi, D. and Belosi, F., 2014. Characterization of PM10 and PM2.5 and their metals content in different typologies of sites in South-Eastern Italy. *Atmosphere*, 5(2), 435-453.
- [29] Mandal, P., Saud, T., Sarkar, R., Mandal, A., Sharma, S.K., Mandal, T.K. and Bassin, J.K., 2014. High seasonal variation of atmospheric C and particle concentrations in Delhi, India. *Environmental Chemistry Letters*, 12(1), 225-230.

- [30] Jain, S., Sharma, S.K., Mandal, T.K. and Saxena. M., 2018. Source apportionment of PM₁₀ in Delhi, India using PCA/APCS, UNMIX, and PMF. *Particuology*, 37, 107-118.
- [31] Kulashrestha, A., Massey, D.D., Masih, J. and Taneja, A., 2014. Source characterization of trace elements in an indoor environment at urban, rural and wayside sites in a semi-arid region of India. *Aerosol and Air Quality Research*, 14, 1738-1751.