

LIST OF TABLES

Table		Page
1	Secondary metabolites from coprophilous fungi.....	21
2	Frequency of various dung samples collected from different provinces	28
3	Dung samples from thirteen wildlife and domestic animals collected from various locations.....	29
4	Species of plant pathogenic fungi from various diseased fruits and vegetables used for antagonistic activity test.....	39
5	Fractionation of the crude extract of <i>Ascodesmis macrospora</i> and <i>Sordaria fimicola</i>	43
6	Frequency of coprophilous fungi and other microfungi found on 60 dung samples.....	46
7	Occurrence of coprophilous and other microfungi on 60 dung samples of 13 wildlife and domestic animals.....	47
8	Number of isolates of coprophilous Zygomycetes and other microfungi obtained from different dung types.....	48
9	Number of isolates of coprophilous Ascomycetes and other microfungi obtained from different dung types.....	50
10	Number of isolates of coprophilous Hyphomycetes and other microfungi obtained from different dung types.....	52
11	Number of isolates of Coelomycetes obtained from different dung types.....	57

LIST OF TABLES (Cont'd)

Table		Page
12	Number of isolates of coprophilous Agonomycetes and other microfungi obtained from different dung types.....	58
13	Percent inhibition on mycelial growth of nine plant pathogenic fungi by <i>Ascodesmis macrospora</i> cultivated on PDA in dual culture test at 28 °C for 14 days.....	133
14	Percent inhibition on mycelial growth of nine plant pathogenic fungi by <i>Sordaria fimicola</i> cultivated on PDA in dual culture test at 28 °C for 14 days.....	135
15	¹³ NMR data (CDCl ₃ , 75.47 MHz) of 3-keto derivative of ergosterol...	141
16	¹³ NMR data (CDCl ₃ , 75.47 MHz) of ergosterol.....	142
17	¹³ NMR data (CDCl ₃ , 75.47 MHz) of 7,8-dehydroergosterol.....	143