

TABLE OF CONTENTS

	Page
TABLE OF CONTENTS	i
LIST OF TABLES	ii
LIST OF FIGURES	v
INTRODUCTION	1
OBJECTIVES	4
LITERATURE REVIEW	5
MATERIALS AND METHODS	14
Materials	14
Methods	14
RESULTS AND DISCUSSION	20
CONCLUSION	43
LITERATURE CITED	45
APPENDIX	49

LIST OF TABLES

Table		Page
1	Vegetative and reproductive stages of a maize plant	7
2	The general analysis of variance combine in split-plot design	19
3	Yield and traits average of eight hybrids in three water treatments: full irrigation (control), mild stress and severe stress.	21
4	Grain yields at 15% moisture content (Mg ha^{-1}) of eight hybrids in three water treatments: full irrigation (control), mild stress and severe stress. The hybrids are ranked according to the yield of the control; the averages of two seasons are presented.	24
5	Grain dry weight per plant (g) of eight hybrids in three water treatments: full irrigation (control), mild stress and severe stress. The hybrids are ranked according to the yield of the control at an area basis, see Table 3; the averages of two seasons are presented.	26
6	Kernel number per hectare (10^4) of eight hybrids in three water treatments: full irrigation (control), mild stress and severe stress. The hybrids are ranked according to the yield of the control at an area basis, see Table 4; the averages of two seasons are presented.	28
7	Kernel number per plant of eight hybrids in three water treatments: full irrigation (control), mild stress and severe stress. The hybrids are ranked according to the yield of the control at an area basis, see Table 4; the averages of two seasons are presented.	29
8	1000 kernels dry weight (g) of eight hybrids in three water treatments: full irrigation (control), mild stress and severe stress in 2003 and 2004. The hybrids are ranked according to the yield of the control at an area basis, see Table 4.	31

LIST OF TABLES (continued)

Table		Page
9	Harvest index of eight hybrids in three water treatments: full irrigation (control), mild stress and severe stress in 2003 and 2004. The hybrids are ranked according to the yield of the control at an area basis, see Table 4.	33
10	Days to 50 percent silking of eight hybrids in three water treatments: fullirrigation(control), mild stress and severe stress.	35
11	Tasseling-silking interval of eight hybrids in three water treatments: full irrigation (control), mild stress and severe stress in 2003 and 2004. The hybrids are ranked according to the yield of the control at an area basis, see Table 4.	37
12	Ear height (cm) of eight hybrids in three water treatments: full irrigation (control), mild stress and severe stress.	39
13	Correlation coefficients calculated among traits measured on 8 tropical maize hybrids grown under water stress.	42
Appendix Table		
1	Analysis of variance for grain yield at 15 % moisture content of 8 hybrids maize	50
2	Analysis of variance for grain dry weigh per plant of 8 hybrids maize	51
3	Analysis of variance for kernel number per hectare of 8 hybrids maize	52
4	Analysis of variance for kernel number per plant of 8 hybrids maize	53

LIST OF TABLES (continued)

Appendix Table		Page
5	Analysis of variance for 1000 kernels dry weight of 8 hybrids maize	54
6	Analysis of variance for harvest index of 8 hybrids maize	55
7	Analysis of variance for days to 50 percent tasseling of 8 hybrids maize	56
8	Analysis of variance for days to 50 percent silking of 8 hybrids maize	57
9	Analysis of variance for tasseling-silking interval (ASI) of 8 hybrids maize	58
10	Analysis of variance for plant height of 8 hybrids maize	59
11	Analysis of variance for ear height of 8 hybrids maize	60

LIST OF FIGURES

Figure		Page
1	Plant responses to environmental stress in correspondence with stress and plant characteristics	11
2	Field layout irrigation management of pre-anthesis drought experiment with split plot design in the dry season (November-February) of 2003/2004 and 2004/2005	16
Appendix Figure		
1	Growth of corn plants which gave irrigation (water) every week interval (W1,Control)	61
2	Growth of corn plants which reduced water 50 % of the control (W2,mild stress)	62
3	Growth of corn plants which did not gave water from the 6 - 7 leaf stage until one week before anthesis thereafter the weekly full furrow irrigation was resumed. (W3, severe stress)	63