

Original Article

Phenolic contents and antioxidant activities of leaf extracts from
Elaeocarpus submonoceras Miq.

Franxisca Mariani^{1,2,3}, Rachaneekorn Tammachote^{1,3}, Irawan Wijaya Kusuma⁴,
Warinthorn Chavasiri⁵, Hunsa Punnapayak^{2,6}, and Sehanat Prasongsuk^{2,6*}

¹ Biological Sciences Program, Faculty of Science,
Chulalongkorn University, Pathum Wan, Bangkok, 10330 Thailand

² Plant Biomass Utilization Research Unit, Department of Botany, Faculty of Science,
Chulalongkorn University, Pathum Wan, Bangkok, 10330 Thailand

³ Division of Genetics, Department of Botany, Faculty of Science,
Chulalongkorn University, Pathum Wan, Bangkok, 10330 Thailand

⁴ Forest Products Chemistry Laboratory, Faculty of Forestry,
Mulawarman University, Samarinda, 75119 Indonesia

⁵ Center of Excellence in Natural Product Chemistry, Department of Chemistry, Faculty of Science,
Chulalongkorn University, Pathum Wan, Bangkok, 10330 Thailand

⁶ Department of Biology, Faculty of Science and Technology,
Universitas Airlangga, Surabaya, 60115 Indonesia

Received: 10 June 2019; Revised: 26 March 2020; Accepted: 6 April 2020

Abstract

The phenolic contents and antioxidant activities of leaf extracts from *Elaeocarpus submonoceras* Miq. were determined for two alternative extractions, namely maceration with ethanol and successive extraction by Soxhlet, with various solvents. Ethanol extract showed the highest phenolic content and strong antioxidant activities against DPPH and ABTS, followed in rank order by methanol and ethyl acetate extracts. The ethyl acetate extract showed no cytotoxic effect towards HaCaT, a human skin non-carcinoma cell. Based on thin layer chromatography, cytotoxic and antioxidant activities, ethyl acetate extract was chosen for isolation. The structure of an isolated compound was identified as gallic acid from ¹H-NMR. This study provides the first scientific information regarding the phenolic content and antioxidant activities of *E. submonoceras* Miq.

Keywords: *Elaeocarpus submonoceras*, phenolic content; antioxidant, cytotoxicity, gallic acid

*Corresponding author
Email address: sehanat.p@chula.ac.th

1. Introduction

Elaeocarpus submonoceras Miq., a member of the Elaeocarpaceae family, is a native tree from Kalimantan and West Java, Indonesia, with edible fruits, and is found in primary or secondary forests, swamps, or dipterocarp forests (Coode, 2001; Uji, 2004). This plant is a sub-canopy tree with alternate, simple, and penni-veined leaves, and white flowers (Figure 1). The seed is oval with a cone shape at the end and has a dark brown color (Lailati & Ekasari, 2015). *E. submonoceras* is also called “Katulampa” by local people in West Java and “Nkodoi” by the Dayak tribe, a local people in Temula Village, West Kutai, East Kalimantan, Indonesia. The leaf has long been used by the Dayak tribe for the treatment of skin diseases, wound healing, and face washing for cleansing and skin brightening purposes. Genus *Elaeocarpus* has approximately 200 species, some of which have been investigated for their chemical constituents (Chand, Dasgupta, Chattopadhyay, & Ray, 1977). Phenolic compounds, such as myricetin and gallic acid, have also been found in this genus (Chand *et al.*, 1977). Phenolic compounds are well known to be related to many biological activities, including antioxidant activities (Fawole, Makunga, & Opara, 2012).

The demand for cosmetic and skin care products has grown significantly in recent years due to the increasing interests in health and beauty, including for maintaining healthy skin. People are also demanding the use of natural ingredients (i.e., organic matter or derivatives, mainly plants) in cosmetic products (Fonseca-Santos, Corrêa, & Chorilli, 2015). Natural products displayed significant scavenging activities against reactive oxygen species, which can cause many skin care problems, such as photoaging, dyschromia, rhytides, and actinic damage (Bowe & Pugliese, 2014). In addition, synthetic ingredients, for example the whitening agent named hydroquinone, can cause ochronosis in long-term use and this is also a carcinogen (Sarkar, Arora, & Garg, 2013).

Consequently, this research aimed to investigate the phenolic content, antioxidant activities and cytotoxic effects of *E. submonoceras* Miq. leaf extracts. Moreover, isolation of the key compound and its structural identification were carried out.



Figure 1. Leaves and flowers of *Elaeocarpus submonoceras* Miq.

2. Materials and Methods

2.1 Plant material

Fresh leaves of *E. submonoceras* Miq. were collected in the rainy season (January to May) from Temula village, West Kutai, East Kalimantan, Indonesia. Mature leaves were collected from 3 to 6 years old trees. A reference specimen was deposited in Wood Chemistry Laboratory, Faculty of Forestry, Mulawarman University, East Kalimantan, Indonesia.

2.2 Leaf extraction

The leaves were air-dried under the shade and then ground. Ten grams of leaf powder were extracted with ethanol (95% v/v, 3x100 mL) for 48 h with constant shaking. The solution was then filtered through Whatman No. 1 filter paper (Sigma-Aldrich, USA) and evaporated at 40°C in a rotary evaporator to obtain ethanol crude extract.

Five-hundred grams of dried ground leaves were extracted successively using Soxhlet (4 L, 12 h) by hexane, dichloromethane, ethyl acetate and methanol, respectively. The solution was evaporated at 40°C in a rotary evaporator to obtain four different extracts.

2.3 Total phenolic content (TPC)

TPC was determined using a method reported by Pientaweeratch, Panapisal, and Tansirikongkol (2016) with modification. The extracts were diluted in 1% dimethyl sulfoxide (DMSO) in distilled water. The final concentration of samples was 50 µg/mL, while gallic acid as a reference was used in the range of 0.195-50 µg/mL. Twenty microliters of extracts and 100 µL of 0.2 N Folin-Ciocalteu reagent (Loba Chemie Pvt. Ltd., India) were added into 96-well plate (#3599, Corning®, NY, USA), then incubated for 5 min. After incubation, 80 µL of sodium carbonate (BDH Chemicals, Toronto, Canada) (75 g/L) was added, slightly shaken and reincubated for 120 min in dark, at room temperature. The absorbance of the mixture reaction was measured at 760 nm by SpectraMax M3 multi-mode microplate reader (Molecular Devices, Sunnyvale, CA, USA) with Softmax software (SoftMax, San Diego, CA, USA). Blank solution contained 1% (v/v) DMSO in distilled water. A standard curve to calculate phenolic contents was made by using linear regression. The results are reported as mg gallic acid equivalent (GAE)/g of dried weight (DW) of extract.

2.4 Antioxidant assays

2.4.1 Total antioxidant capacity (TAC)

The method for determining TAC was modified from Prieto, Pineda, and Aguilar (1999). Final concentrations of ascorbic acid were varied from 5 to 100 µg/mL, while the extract concentration was 50 µg/mL. Forty microliters of each extract were pipetted into 1.5 mL tube, and 400 µL of TAC reagent solution (0.6 M H₂SO₄, 28 mM sodium phosphate and 4 mM ammonium molybdate) were added then incubated for 90 min at 95°C. Two-hundred microliters from each reaction tube were transferred into a 96-well plate, and allowed to cool

to room temperature. The absorbance of reaction mixture was measured at 695 nm against a blank (contained 1% DMSO in distilled water only). The ascorbic acid reference standard results were plotted to make a standard curve by linear regression. TAC values are expressed as mg ascorbic acid equivalent (AAE)/g of dried weight (DW) of extract.

2.4.2 DPPH (2,2-diphenyl-1-picrylhydrazyl) inhibition

DPPH inhibition assay was modified from Lin *et al.* (2014). Reaction was made by mixing fifty microliters of sample (various concentrations of extracts or the positive control, ascorbic acid (from zero to 1000 µg/mL)) dissolved in 1% DMSO-methanol with 150 µL of 100 µM DPPH reagent in methanol, then incubated for 30 min in the dark. The inhibition of DPPH in the samples caused discoloration from purple to pale yellow and was measured at 512 nm using SpectraMax M3 reader.

An equation to estimate the percentage of DPPH free radical inhibition was adopted from Benmehdi, Behlil, Memmou, and Amrouche (2017):

$$\frac{A_{\text{control}} - A_{\text{sample}}}{A_{\text{control}}} \times 100\% \quad (1)$$

where A_{sample} is the absorbance from the reaction of DPPH with the sample and A_{control} is the absorbance of DPPH and solution only (without sample). Results from this equation were plotted and normalized using linear regression to get IC_{50} (concentration of the sample when inhibiting 50% of DPPH). Results of percent inhibition were calculated for 100 µg/mL sample concentration.

2.4.3 ABTS (2,2'-azino-bis(3-ethylbenzothiazoline-6-sulfonic acid) diammonium salt) inhibition

The ability of the sample to scavenge ABTS radical was assayed based on Fu, Zhang, Guo, and Chen (2014) with slight modification in a solution used to dilute ABTS^{•+} and potassium persulfate reagents, by change to be distilled water. The absorbance was measured within three minutes. Equation from linear regression of percent inhibition of sample was used to calculate IC_{50} . Concentrations of sample and ascorbic acid varied from 0.78 to 1000 µg/mL. Result of percent inhibition was obtained for 100 µg/mL sample concentration.

2.5 Cytotoxicity effect

Cytotoxicity of the plant extracts was determined by the method modified from Ritto *et al.* (2017). HaCaT human keratinocyte cell was maintained in DMEM (Gibco, NY, USA) and supplemented with 1% antibiotic-antimycotic 100x (Gibco, NY, USA), 10% heat-inactivated fetal bovine serum (HIMEDIA, Mumbai, India), and 1% 200 mM glutamine (PAA, Pasching, Austria). The cells were grown at 37°C and 5% CO₂ in a cell culture incubator. In 96-well plates, 10,000 cells/well in 100 µL of media were seeded and incubated for 48 h. Samples were prepared in 1% DMSO-complete media. After adding the samples, the cells were reincubated for another 48 h and then changed to 10% PrestoBlue (Invitrogen, CA, USA)-complete media. After incubating for 2 h,

fluorescence units were measured at 570 (excitation) and 600 nm (emission) based on the manufacturer's reference. The cytotoxicity effect was determined from percentage of viable cells when treated with various final concentrations of extracts (10 to 1000 µg/mL). The IC_{50} value was obtained for 50% of cell viability by plotting the percentages from logarithmic regression in Excel.

2.6 Isolation of compound and structural identification

Ethyl acetate extract was isolated based on its thin layer chromatography, antioxidant activities and cytotoxic effect. The ethyl acetate extract (5.28 g) was eluted using silica gel 60 column chromatography with solvent system *n*-hexane-ethyl acetate (3:1, 2:1, 1:1, 1:3 v/v), ethyl acetate 100% followed by ethyl acetate-methanol (98:2, 95:5, 9:1, 7:3, 6:4, 5:5, 3:7, 2:8, 1:9 v/v) and methanol 100% to yield 13 fractions (E.1 – E.13). Fraction E.10 (324.9 mg) was then washed with dichloromethane and the precipitant (217.6 mg) was purified by crystallization with dichloromethane: methanol: water: acetic acid (90:10:1:0.1 v/v/v/v). The isolate compound was found in E10.11 (12.2 mg). The purified compound (E10.11) was analyzed using BRUKER proton nuclear magnetic resonance spectroscopy (¹H NMR) 400 MHz in acetone ((CD₃)₂CO).

2.7 Statistical analysis

One-way analysis of variance (ANOVA) with significant differences ($P < 0.05$) was used for statistical analysis in Microsoft Excel Office 365 and IBM SPSS statistics 22 with Duncan's post hoc test. Results are displayed as mean values with standard deviations (mean±SD) from three replications of each sample.

3. Results and Discussion

3.1 TPC, TAC, DPPH, and ABTS of extracts

The leaf ethanol extract had a high phenolic content and high antioxidant activities as listed in Table 1. The ethanol extract and ascorbic acid had similar inhibitions of DPPH and ABTS at the same concentration (100 µg/mL) in the range from 94.98% to 100%. The ethanol extract of this plant had an excellent ability to scavenge free radicals, with IC_{50} values of 3.81 µg/mL DPPH and 13.51 µg/mL ABTS, compared with 12.72 µg/mL and 3.07 µg/mL for ascorbic acid (a well-known antioxidant compound), respectively.

Ethanol is a solvent that is commonly used to extract phenolic compounds. Phenolics have several distinctive characteristics, including their color, flavor, odor, bitterness, and biological activities, which have led to their use in many food and medicinal applications (Naczka & Shahidi, 2006).

The present study determined that ethyl acetate extract was not significantly different from the methanol extract. Both extracts had the highest phenolic content with no significant difference (504.92 and 542.81 mg GAE/g DW). They also had excellent activity to scavenge DPPH and ABTS free radicals with an IC_{50} range from 7.67 to 14.00 µg/mL, while the least activity was for the hexane extract. Based on

Table 1. TPC, TAC, DPPH and ABTS inhibitions of *Elaeocarpus submonoceras* Miq. extracts. The percentage inhibition of DPPH and ABTS is given for a concentration of 100 µg/mL. All results in this table are shown as mean ± SD (n = 3). Significant differences ($P < 0.05$) are shown by different superscript letters.

Extraction process/sample	TPC (mg GAE/g DW)	TAC (mg AAE/g DW)	DPPH		ABTS	
			Inhibiton (%)	IC ₅₀ (µg/mL)	Inhibiton (%)	IC ₅₀ (µg/mL)
Positive control Ascorbic acid	-	-	96.60 ± 1.05 ^a	12.72 ± 0.32 ^c	100.00 ± 0.10 ^a	3.07 ± 0.04 ^a
Maceration Ethanol	578.45 ± 31.44 ^a	419.73 ± 3.95 ^a	94.98 ± 0.29 ^a	3.81 ± 0.06 ^a	96.85 ± 0.11 ^b	13.51 ± 0.67 ^b
Successive Hexane	7.65 ± 0.55 ^c	53.22 ± 3.08 ^e	35.37 ± 2.31 ^c	92.10 ± 4.69 ^d	11.04 ± 0.34 ^d	>1000
Soxhlet Dichloromethane	57.23 ± 4.08 ^c	90.11 ± 4.59 ^d	64.99 ± 2.69 ^b	307.33 ± 0.06 ^e	13.98 ± 0.76 ^e	827.82 ± 5.20 ^e
extraction Ethyl acetate	504.92 ± 43.23 ^b	274.21 ± 6.43 ^c	94.71 ± 0.66 ^a	7.81 ± 0.19 ^b	95.94 ± 1.10 ^b	14.00 ± 0.57 ^b
	Methanol	542.81 ± 50.11 ^{ab}	343.45 ± 1.51 ^b	94.07 ± 0.14 ^a	7.67 ± 0.09 ^b	96.75 ± 0.18 ^b

these results, it was determined that the antioxidant activities originated from polar compounds were related to the abundance of phenolics in the extract. Similar to another report, the successive extraction of three species of *Mentha* (*M. spicata*, *M. pulegium*, and *M. piperita*) also had DPPH inhibition only from polar extracts (methanol and water) rather than from non-polar ones (hexane and dichloromethane) (Barchan, Bakkali, Arakrak, Pagán, & Laglaoui, 2014). Polar extracts were related to the phenolic content. Li *et al.* (2017) identified 15 phenolic compounds and concluded that DPPH and ABTS scavenging activities had a positive correlation with the phenolic constituents.

Antioxidant activity is a biological activity highly related to phenolics due to the donation of a hydrogen cation from the phenolic hydroxyl group to free radicals, generating relatively stable free radicals (Prihantini, Tachibana, & Itoh, 2015). Studies of the antioxidant activity are often performed by phosphomolybdenum, DPPH, and ABTS scavenging methods, which are widely used in plant and food research (Basumatary, Das, Nanjian, & Sharma, 2015; Pientaweeratch *et al.*, 2016).

Phenolic compounds that are also active as antioxidants can be found in various *Elaeocarpus* species. For example, an ethanol extract of leaves of *E. lanceofolius* contained 4'-methylmyricetin and myricetin-3-O-rhamnoside (Ray, Dutta, & Dasgupta, 1976). Mearnsetin was found in *E. serratus*, *E. oblongus*, and *E. floribundus*. Quercetin, kaempferol, and ethyl gallate were found in *E. tuberculatus*. Ellagic acid was found in *E. serratus* and *E. ganitrus* Roxb. (Chand *et al.*, 1977). Ethanol extract from leaves of *E. sphaericus* also showed the presence of gallic acid (Garg, Goswami, & Khurana, 2012). Gallic acid, a simple phenolic compound, possesses strong antioxidant activity and is easily absorbed by the human metabolic system (Badhani, Sharma, & Kakkar, 2015). Its mechanism in the human body involves counteracting with reactive oxygen species to produce antioxidants. However, an imbalance in the amount of antioxidants in the body can cause many diseases (Kurutas, 2016). Therefore, natural antioxidant agents that may be in leaf extracts could have an important role to play in safeguarding human health. Ethanol extract from *E. submonoceras* Miq. is promising as an antioxidant agent and could inhibit ABTS better than *E. ganitrus* leaf ethanol extract. *E. ganitrus* showed 55.77% ABTS inhibition at a concentration of 500 µg/mL and had a higher IC₅₀ value (297.12 µg/mL). This species also had a lower TPC at 56.79 mg GAE/g DW (Kumar, Shanmugam, Palvannan, & Kumar,

2007). *E. ganitrus* bark ethanol extract was also found to have a higher IC₅₀ value (81.85 µg/mL) against DPPH free radicals and phenol compounds were detected in a phytochemicals test (Talukdar, Dutta, Chakraborty, & Das, 2017).

3.2 Cytotoxic effects of *E. submonoceras* Miq. extracts

For commercial applications, comprehensive information about the content, including cytotoxicity against normal cells, is needed (Fonseca-Santos *et al.*, 2015). The extracts of *E. submonoceras* Miq. in concentrations ranging from 10–1000 µg/mL reduced viability of HaCaT, human skin non-carcinoma cells, in a dose-dependent manner (data not shown). Interestingly, the calculated IC₅₀ values of ethanol, ethyl acetate, and methanol were 258.98, 446.25, and 230.27 µg/mL, respectively (Figure 2) and did not indicate a toxic effect compared to the IC₅₀ values of antioxidant assays (Table 1). These results were similar to those from a previous study, which involved treating HaCaT and fibroblast cells with a leaf ethanol extract of *Helianthus tuberosus* L. The proliferation of normal cells after extract treatment may be related to the high concentration of phenolic compounds (Nizioł-Lukaszewska, Furman-Toczek, & Zagórska-Dziok, 2018). Methanol extracts of *E. tuberculatus* and *E. serratus* also showed low cytotoxicity against Vero cells (normal African green monkey kidney epithelial cells), in a concentration dependent manner (Shailasree, Sampathkumara, Niranjana, & Prakash, 2014).




Figure 2. IC₅₀ of *Elaeocarpus submonoceras* Miq. leaf extracts when treating HACAT cells for 48 h. All results are represented as mean + SD (n = 3).

3.2 Isolation of the active compound

The isolation process started from the separation of the compounds based on their polarity, followed by conducting TPC and antioxidant assays. DPPH and ABTS inhibition assays were used to evaluate all the extracts and

fractions until the pure compound was obtained. To date, no information on the phenolic content and antioxidant activities related to successive extracts of *Elaeocarpus* genus have been reported. The ethyl acetate extract was selected for further fractionation, purification, and isolation of the pure compound because its phenolic content, antioxidant activities, TLC; and this extract was the most non-toxic when tested against HaCaT cells (section 3.2).

This compound displayed as blue color when reacted with Folin–Ciocalteu’s reagent (data not shown) and only one spot when confirmed by a TLC plate with variation of the solvent system (data not shown). The compound (E10.11) was obtained at a weight of about 12.2 mg as a white powder. ^1H NMR was used to identify the compound ($(\text{CD}_3)_2\text{CO}$, 400 MHz), δ (ppm): 7.15 (d, $J=1.2$ Hz, 2 H (H-2 and H-6)) (Figure 3), and the results revealed that it was gallic acid. The ^1H NMR result is similar to previous reports related to the *Elaeocarpus* genus. Leaf extracts of *E. sylvestris* and *E. tonkinensis* also contained gallic acid (Dao *et al.*, 2018; Prihantini *et al.*, 2015). Gallic acid is also one of the major compounds in *E. tuberculatus* and *E. ganitrus* (Chand *et al.*, 1977). Gallic acid or 3,4,5-trihydroxybenzoic acid is a member of the phenolic group, with one aromatic ring, one carboxylic acid, and three hydroxyl substituents. It has a low molecular weight with a simple structure, but possesses high antioxidant activities (Badhani *et al.*, 2015).

4. Conclusions

Information from the Dayak tribe regarding *E. submonoceras* Miq. used for skin and wound treatment was validated in this study. Leaf extracts from *E. submonoceras* Miq. showed abundant phenolic content, indicating it may be used as a safe antioxidant agent. Gallic acid was identified as one of the active compounds. Information of antioxidant activities from this study could help expand the utilization of this plant as a supplement for skin care or in cosmetics production, and could lead to a further study of the antioxidant activities of this plant extract *in vivo*.

Acknowledgements

Financial support from the East Kalimantan and ASEAN Scholarship, Chulalongkorn University to Franxisca Mariani are gratefully acknowledged. The authors thank Hastaniah, M.Agr., Laboratory of Dendrology and Forest Ecology, Faculty of Forestry, Mulawarman University for identifying the plant species.

References

- Badhani, B., Sharma, N., & Kakkar, R. (2015). Gallic acid: a versatile antioxidant with promising therapeutic and industrial applications. *RSC Advances*, 5(35), 27540-27557. doi:10.1039/C5RA01911G.
- Barchan, A., Bakkali, M., Arakrak, A., Pagán, R., & Laglaoui, A. (2014). The effects of solvents polarity on the phenolic contents and antioxidant activity of three *Mentha* species extracts. *International Journal of Current Microbiology and Applied Sciences*, 3(11), 399-412. Retrieved from <https://www.ijcmas.com/vol-3-11/A.Barchan,%20et%20al.pdf>
- Basumatary, S., Das, A. K., Nanjian, R., & Sharma, G. D. (2015). *In vitro* evaluation of antioxidant properties of *Hodgsonia heteroclita* (Cucurbitaceae) fruit. *Journal of Applied Pharmaceutical Science*, 5(3), 80-83. doi:10.7324/JAPS.2015.510.S14.
- Benmehdi, H., Behilil, A., Memmou, F., & Amrouche, A. (2017). Free radical scavenging activity, kinetic behavior and phytochemical constituents of *Aristolochia clematitis* L. roots. *Arabian Journal of Chemistry*, 10(1), S1402-S1408. doi: 10.1016/j.arabjc.2013.04.015.
- Bowe, W. P., & Pugliese, S. (2014). Cosmetic benefits of natural ingredients. *Journal of Drugs in Dermatology*, 13(9), 1019-1027. Retrieved from <https://jddonline.com/articles/dermatology/S1545961614P1021X>
- Chand, L., Dasgupta, S., Chattopadhyay, S. K., & Ray, A. B. (1977). Chemical investigation of some *Elaeocarpus* species. *Planta Medica*, 32(2), 197-199. doi:10.1055/s-0028-1097584.




Figure 3. ^1H NMR of the purified compound (E10.11)

- Coode, M. J. E. (2001). *Elaeocarpus* for flora Malesiana – the *E. stipulom* complex, *E. nitidus* group & *E. barbulatus*. *Kew Bulletin*, 56(3), 513-565. doi:10.2307/4117684.
- Dao, N. T., Jang, Y., Kim, M., Nguyen, H. H., Pham, D. Q., Dang, Q. L., . . . Hoang, V. D. (2019). Chemical constituents and anti-influenza viral activity of the leaves of Vietnamese plant *Elaeocarpus tonkinensis*. *Records of Natural Products*, 13(1), 71-80. doi:10.25135/rnp.76.18.04.266.
- Fawole, O. A., Makunga, N. P., & Opara, U. L. (2012). Antibacterial, antioxidant and tyrosinase-inhibition activities of pomegranate fruit peel methanolic extract. *BMC Complementary and Alternative Medicine*, 12(200), 1-11. doi:10.1186/1472-6882-12-200.
- Fonseca-Santos, B., Corrêa, M. A., & Chorilli, M. (2015). Sustainability, natural and organic cosmetics: consumer, products, efficacy, toxicological and regulatory considerations. *Brazilian Journal of Pharmaceutical Sciences*, 51(1), 17-26. doi:10.1590/S1984-82502015000100002.
- Fu, R., Zhang, Y., Guo, Y., & Chen, F. (2014). Antioxidant and tyrosinase inhibition activities of the ethanol-insoluble fraction of water extract of *Sapium sebiferum* (L.) Roxb. leaves. *South African Journal of Botany*, 93, 98-104. doi:10.1016/j.sajb.2014.04.003.
- Garg, K., Goswami, K., & Khurana, G. (2012). A pharmacognostical review on *Elaeocarpus sphaericus*. *International Journal of Pharmacy and Pharmaceutical Sciences*, 5(1), 3-8. Retrieved from <https://www.semanticscholar.org/paper/A-PHARM-ACOGNOSTICAL-REVIEW-ON-ELAEOCARPU-S-Garg-Goswami/2a7cc2d378eb83aefac941b9ac6feb6a80208682>
- Kumar, T. S., Shanmugam, S., Palvannan, T., & Kumar, M. B. (2007). Evaluation of antioxidant properties of *Elaeocarpus ganitrus* Roxb. leaves. *Iranian Journal of Pharmaceutical Research*, 7(3), 211-215. Retrieved from http://ijpr.sbmu.ac.ir/article_767_1b1ff80928d873f3d2e6f7ec25f6268f.pdf
- Kurutas, E. B. (2016). The importance of antioxidants which play the role in cellular response against oxidative/nitrosative stress: Current state. *Nutrition Journal*, 15(1), 71. doi:10.1186/s12937-016-0186-5.
- Lailati, M., & Ekasari, I. (2015). Unique and interesting seeds of Cibodas Botanic Garden Collections. Translate to English, add term *Proceedings of National Seminar on Indonesian Biodiversity Society I* (In Indonesian), 1328-1333. doi:10.13057/psnmbi/m01 0612
- Li, H., Zhang, D., Tan, L.-H., Yu, B., Zhao, S.-P., & Cao, W.-G. (2017). Comparison of the antioxidant properties of various solvent extracts from *Dipsacus asperoides* and identification of phenolic compounds by LC-ESI-QTOF-MS-MS. *South African Journal of Botany*, 109, 1-8. doi:10.1016/j.sajb.2016.12.018.
- Lin, F.-J., Yen, F.-L., Chen, P.-C., Wang, M.-C., Lin, C.-N., Lee, C.-W., & Ko, H.-H. (2014). HPLC-fingerprints and antioxidant constituents of *Phyllanthus nodiflora*. *The Scientific World Journal*, 2014(528653), 1-8. doi:10.1155/2014/528653.
- Naczki, M., & Shahidi, F. (2006). Phenolics in cereals, fruits and vegetables: occurrence, extraction and analysis. *Journal of Pharmaceutical and Biomedical Analysis*, 41(5), 1523-1542. doi:10.1016/j.jpba.2006.04.002.
- Nizioł-Lukaszewska, Z., Furman-Toczek, D., & Zagórska-Dziok, M. (2018). Antioxidant activity and cytotoxicity of Jerusalem artichoke tubers and leaves extract on HaCaT and BJ fibroblast cells. *Lipids in Health and Disease*, 17(1), 280. doi:10.1186/s12944-018-0929-8.
- Pientaweeratch, S., Panapisal, V., & Tansirikongkol, A. (2016). Antioxidant, anti-collagenase and anti-elastase activities of *Phyllanthus emblica*, *Manilkara zapota* and silymarin: an in vitro comparative study for anti-aging applications. *Pharmaceutical Biology*, 54(9), 1865-1872. doi:10.3109/13880209.2015.1133658.
- Prieto, P., Pineda, M., & Aguilar, M. (1999). Spectrophotometric quantitation of antioxidant capacity through the formation of a phosphor molybdenum complex: specific application to the determination of vitamin E. *Analytical Biochemistry*, 269(2), 337-341. doi:10.1006/abio.1999.4019.
- Prihantini, A. I., Tachibana, S., & Itoh, K. (2015). Antioxidant active compounds from *Elaeocarpus sylvestris* and their relationship between structure and activity. *Procedia Environmental Sciences*, 28, 758-768. doi:10.1016/j.proenv.2015.07.089.
- Ray, A. B., Dutta, S. C., & Dasgupta, S. (1976). Flavonoids of *Elaeocarpus lanceofolius*. *Phytochemistry*, 15(11), 1797-1798. doi:10.1016/S0031-9422(00)97498-3.
- Ritto, D., Tanasawet, S., Singkhorn, S., Klaypradit, W., Hutamekalin, P., Tipmanee, V., & Sukketsiri, W. (2017). Astaxanthin induces migration in human skin keratinocytes via Racl activation and RhoA inhibition. *Nutrition Research and Practice*, 11(4), 275-280. doi:10.4162/nrp.2017.11.4.275.
- Sarkar, R., Arora, P., & Garg, K. V. (2013). Cosmeceuticals for hyperpigmentation: what is available?. *Journal of Cutaneous and Aesthetic Surgery*, 6(1), 4-11. doi:10.4103/0974-2077.110089.
- Shailasree, S., Sampathkumara, K. K., Niranjana, S. R. & Prakash, H. S. (2014). Bioactive potential of medicinal plants from Western Ghats Region, India. *Journal of Herbs, Spices and Medicinal Plants*, 20(3), 221-234. doi:10.1080/10496475.2013.860070.
- Talukdar, N., Dutta, A. M., Chakraborty, R., & Das, K. (2017). Screening of phytochemicals, antioxidant and inhibitory effect on alpha-amylase by ethanolic extract of *Elaeocarpus ganitrus* (bark). *International Journal of Pharmaceutical Sciences and Research*, 8(12): 5270-75. doi:10.13040/IJPSR.0975-8232.8(12).5270-75.
- Uji, T. (2004). Species diversity, genetic resources, and potential of the indigenous fruits in Kalimantan. *BioSmart*, 6(2), 117-125. Retrieved from <https://www.scribd.com/document/349085794/B060210-pdf>