

ACKNOWLEDGEMENTS

This work was partially supported by Thesis and Dissertation support fund from the Graduate School, Kasetsart University, Commission on Higher Education, Chachoengsao Rubber Research Center, CIRAD (Centre de Coopération Internationale en Recherche Agronomique pour le Développement – French Agricultural Research Centre for International Development in Cooperation, France and Thailand) and French Embassy to Thailand.

I wish to express my profound gratitude to my thesis advisors Assistant Professor Dr. Sornparch Thanisawanyangkura, Associate Professor Kamolpun Namwongprom, Assistant Professor Dr. Poonpipope Kasemsap, Dr. Eric Gohet and Dr. Philippe Thaler for their supports, comments, suggestion, guidance and encouragement.

I would like to express my gratitude and appreciation to Mrs. Anne Clement-vidal, Dr. Agnes Guilliot, Dr. Andre Lacointe and Dr. Andre Clement-demange who help me a lot for their fruitful comments, suggestion and friendliness. And also thanks all staffs in INRA-UMR, PIAF and CIRAD for their helps and friendship during my work. Thanks you go to all of staffs in Plant Physiology of Horticulture, Plant Architecture Research Lab of Botany, Kasetsart University and CIRAD in Thailand for many helps.

I would also thank Directors of Chachoengsao Rubber Research Center : Mr. Suchin Maemuen and Mr. Theerachart Vichitchochai for their facilities support. Special thanks to my staffs who work both laboratory and field.

Finally, I would like to thank my warm and cheerful family: my parents, husband and son, relatives and friends who always stand by me.

Pisamai Chantuma

May, 2007