

เอกสารอ้างอิง

- เกศิณี ตระกูลทิวกร และ จันทร์เพ็ญ ศักดิ์สิทธิพิทักษ์, 2539, ศักยภาพในการต้านสารอนุมูลอิสระของสารสกัดจากผักพื้นบ้านไทย, รายงานการวิจัยประจำปี 2539-2542 สถาบันค้นคว้าและวิจัยพัฒนาผลิตภัณฑ์อาหาร มหาวิทยาลัยเกษตรศาสตร์, หน้า 13-23.
- Aharonson, Z., Shani, J., and Sulman, F.G., 1969, "Hypoglycemic effect of the salt bush (*Atriplex halimus*): a feeding source of the sand rat (*Psammomys obesus*)", **Diabetologia**, Vol. 5, pp. 379-383
- Alzoreky, N.S. and Nakahara, K., 2003, "Antibacterial activity of extracts from some edible plants commonly consumed in Asia", **International Journal of Food Microbiology**, Vol. 80, pp. 223– 230.
- An, R .B., Sohn, D.H., Jeong, G, J., and Kim, Y, C., 2008, "In vitro hepatoprotective compounds from *Sueada glauca*", **Archives of Pharmacol Research**, Vol. 31, No. 5, pp. 594-597.
- Anonymous, 1999, **Physiology & Biochemistry of Microorganisms**, SIUC/College of Science/Microbiology, November 2008,
<http://www.micro.siu.edu.micr425/425Notes/ 01-Introduction.html>.
- Asano, N.K., Yokoyama, M., Sakurai, K., Kizu, A., Kato, M., Arisawa, D., Hoke, B., Drager, A., and Watson, R.N., 2001, "Dihydroxynortropane alkaloids from calystegine-producingplants", **Phytochemistry**, Vol. 57, pp. 721-726.
- Atalay M, Gordillo G, Roy S, Rovin B, Bagchi D, Bagchi M, Sen CK. Anti-angiogenic property of edible berry in a model of hemangioma. *FEBS Lett.* 2003 Jun 5;544(1-3):252-7.
- Basilico, M.Z., and basilico, J.C., 1999, "Inhibitory effects of some spice essential oils on *Aspergillus ochraceus* NRRL 3174 growth and ochratoxin A production", **Letters in Applied Microbiology**, Vol. 29, pp. 238-241.
- Benedict, A.C., 2009, **Extraction of The Essential Oil of Aquilaria Malaccensis (Gaharu) Using Hydro-Distillation and Solvent Extraction Methods**, Faculty of Chemical & Natural Resources Engineering University Malaysia Pahang, Thesis.
- Benkeblia, N., 2004, "Antimicrobial activity of essential oil extracts of various onions (*Allium cepa*) and garlic (*Allium sativum*)", **Lebensmittel Wissenschaft und Technologie**, Vol. 37, pp. 263-268.

- Benwahhoud, M., Jouad, H., Eddouks, M., and Lyoussi, B., 2001, "Hypoglycemic effect of *Suaeda fruticosa* in streptozotocin-induced diabetic rats", **Journal of Ethnopharmacology**, Vol. 76, pp. 35-38.
- Ben-Yehosha, S., Rodov, V., and Perez, J., 1998, **Constitutive and Induced Resistance of Citrus Fruits Against Pathogens**, In Johnson, G.I., Highly, E., and Joyce, D.C. (Eds.), Disease resistance in fruits, ACIAR proceeding no. 80, Canberra, Australia: Australian Center for International Agricultural Research, pp. 78-89.
- Block, G., Patterson, B., and Subar, A., 1992, "Fruits vegetables and cancer prevention: A review of the epidemiological evidence", **Nutrition and Cancer**, Vol. 18, pp. 1-29.
- Braga, M.E.M., Leal, P.F., Carvalho, J.E., and Meireles, M.A., 2003, "Comparison of yield, composition, and antioxidant activity of turmeric (*Curcuma longa* L.) extracts obtained using various techniques", **Journal of Agricultural and Food Chemistry**, Vol. 51, pp. 6604-6611.
- Buxiang, S. and Fukuraha, M., 1997, "Effects of co-administration of butylated hydroxytoluene, butylated hydroxyanisole and flavonoids on the activation of mutagens and drug-metabolizing enzymes in mice", **Toxicology**, Vol. 122, pp. 61-72.
- Caccioni, D.R.L., Guizzardi, M., Biondi, D.M., Renda, A., and Ruberto, G., 1998, "Relationships between volatile components of citrus fruit essential oils and antimicrobial action on *Penicillium digitatum* and *Penicillium italicum*", **International Journal of Food Microbiology**, Vol. 43, pp. 73-79.
- Casey, R., West, S.I., Hardy, D., Robinson, D.S., Wu, Z., and Hughes, R.K., 1999, "New frontiers in food enzymology; recombinant lipoxygenases", **Trends in Food Science and Technology**, Vol. 10, pp. 297-302.
- Cateni, F., Falsone, G., Vitrotti, E., Zillic, J., Birkofer, L., Della Loggia, R., and Soa, S., 2000, "Constituents of Euphorbiaceae: An anti-inflammatory digalactosyl diacylglycerol from *Euphorbia peplis* L.", **Pharmaceutical and Pharmacological Letters**, Vol. 2, pp. 65-67.
- Cerruti, P. and Alzamora, S.M., 1996, "Inhibitory effects of vanillin on some food spoilage yeasts in laboratory media and fruit purees", **International Journal of Food Microbiology**, Vol. 29, pp. 379-386.

- Chandrasekaran, M., Kannathasan, K., and Venkatesalu, V., 2008, "Antimicrobial activity of fatty acid methyl esters of some members of Chenopodiaceae, **Zeitschrift für Naturforschung**, Vol. 63c, pp. 331-336.
- Cho, S.H., Lee, S.Y., Kim, J.W., Ko, G.H., and Seo, I.W., 1995, "Development and application of natural antimicrobial agent isolated from grapefruit seed extract-antimicrobial activity of grapefruit seed extract", **Journal Food Hygiene and Safety**, Vol. 10, pp. 33-39.
- Conolly, J. and Hill, R., 1996, "Triterpenoids", **Natural Products Report**, pp. 151-169.
- Corbo, M.R., Lanciotti, R., Gardini, F., Sinigaglia, M., and Guerzoni, M.E., 2000, "Effects of hexanal, E(2)-hexenal, and storage temperature on shelf-life of fresh sliced apples", **Journal of Agricultural and Food Chemistry**, Vol. 48, pp. 2401-2408.
- Delaquis, P.J., Stanich,K., Girard, B., and Mazza, G., 2002, "Antimicrobial activity of individual and mixed fractions of dill, citantro, coriander and eucalyptus essential oils", **International Journal of Food Microbiology**, Vol. 74, pp. 101-109.
- Devi, K.P., Suganthy, N., Kesika, P., and Pandian, S.K., 2008, "Bioprotective properties of seaweeds", **Malaysian Journal of Nutrition**, Vol. 8, No.2, pp. 167-177.
- Dorman, H.J.D. and Deans, S.G., 2000, "Antimicrobial against from plants; antimicrobial activity of plant volatile oils", **Journal of Applied Microbiology**, Vol. 88, pp. 308-316.
- Dorman, H.J.D., Kosar, M., Kahlos, K., Holm, Y., and Hiltunen, R., 2003, "Antioxidant properties and composition of aqueous extracts from *Mentha* species, hybrids, varieties, and cultivars", **Journal of Agricultural and Food Chemistry**, Vol. 51, pp. 4563-4569.
- Fahay, J.W. and Stephenson, K.K., 2002, "Pinostrobin from honey and Thai ginger (*Boesenbergia pandurata*): a potent flavonoid inducer of mammalian phase 2 chemoprotective and antioxidant enzymes", **Journal of Agricultural and Food Chemistry**, Vol. 50, pp. 7472-7476.
- Farag, R.S., Daw, Z.Y., Hewedi, F.M., and El-Baroty, G.S.A., 1989, "Antimicrobial activity of some Egyptian spice essential oils", **Journal of Food Protection**, Vol. 85, pp. 289-293.

- Gardini, F., Lenciotti, R., and Guerzoni, M.E., 2001, "Effect of (*E*)-2-hexenal on the growth of *Aspergillus flavus* in relation to its concentration, temperature and water activity", **Letters in Applied Microbiology**, Vol. 33, pp. 50-55.
- Gardini, F., Lenciotti, R., Belletti, N., and Guerzoni, M.E., 2002, "Use of natural aroma compounds to control microbial growth in foods", **Kerala: Global Research Network**, pp. 63-67.
- Gaspar, E., 2001, "SoldanelineB: The first acylated non-linear tetrasaccharide macrolactone from the European Convolvaceae plant *Calystegia soldanella*", **European, Journal of Organic Chemistry**, Vol. 2, pp. 369-694.
- Gill, A.O., Delaquis, P.J., Russo, P., and Holley, R.A., 2002, "Evaluation of antilisterial action of cilantro oils on vacuum packed ham", **International Journal of Food Microbiology**, Vol. 73, pp. 83-92.
- Gorham, J. and Wyn Jones, R.G., 1983, "Solute distribution in *Suaeda maritima*", **Planta**, Vol. 157, pp. 344-349.
- Gotta, H., Adolf, W., Opferkuch, H., and Hecker, E., 1984, "On the active principles on the Euphorbiaceae, IX. Ingenane type diterpene esters from five Euphorbia species", **Zeitschrift für Naturforschung**, Vol. 39b, pp. 683-694.
- Gurdip, S., Sumitra, C., Catalan, C., and Lampasona, M.P., 2004, "Chemical constituents antifungal and antioxidative effects of Ajwain essential oil and its acetone extract", **Journal of Agricultural and Food Chemistry**, Vol. 52, pp. 3292-3296.
- Hammer, K.A., Carson, C.F. and Riley T.V., 1999, "Antimicrobial activity of essential oils and other plant extracts", **Jounal of Applied Microbiology**, Vol. 86, pp. 985-990.
- Hemmers, H. and Gblz, P., 1986, "Epicuticular waxes from leaves of five Euphorbia species", **Phytochemistry**, Vol. 25, pp. 2103-2107.
- Hirose, M., Takesada, Y., Tanaka, H., Tamano, S., Kato, T., and Shirai, T., 1998, "Carcinogenicity of antioxidants BHA, caffeic acid, sesamol, 4-methoxyphenol and catechol at low doses, either alone or in combination and modulation of their effects in a rat medium-term multi-organ carcinogenesis model", **Carcinogenesis**, Vol. 19, pp. 207-212.
- Hsieh, P.C., Mau, J.L., and Huang, S.H., 2001, "Antimicrobial effect of various combinations of plants extracts", **Food Microbiology**, Vol. 18, pp. 35-43.

- Hyun-Jin, K., Feng, C., Changqing, W., Hau-Yin, C. and Zhenaju, J., 2004, "Evolution of antioxidant activity of Australian tea tree (*Melaleuca alternifolia*) oil and its components", **Journal of Agricultural and Food Chemistry**, Vol. 52, pp. 2849-2854.
- Ivanova, A., Nechev, J., and Stefanov, K., 2006, "Effect of soil salinity on the lipid composition of halophyte plants from the sand bar of Pomorie", **Plant Physiology** (Special Issue), pp. 125-130.
- Jitoe, A, Masuda, T, Tengah, I. G. P., Suprapta, Dewa N., Gara I. W., Nakatan, N., 1992 Antioxidant activity of tropical ginger extracts and analysis of the contained curcuminoids, **Journal of Agricultural and Food Chemistry**, Vol 40 (8), pp 1337-1340.
- Kikuzari, H. and Nikatani, N., 1993, "Antioxidants effects of some ginger constituents", **Journal of Food Science**, Vol. 58, pp. 1407-1410.
- Kim, J.M., Lee, Y.J., Hong, K.H., Kwon, Y.K., Sim, K.C., Lee, J.Y., Cho, H.Y., Kim, I.S., Han, S.B., Lee, C.W., Shin, I.S., and Cho, J.S., 2001, "Isolation of antimicrobial substances from natural products and their preservative effects", **Food Science Biotechnology**, Vol. 10, No. 1, pp. 59-71.
- Kubo, I. and Fujita, K., 2001, "Naturally occurring anti-Salmonella agents", **Journal of Agricultural and Food Chemistry**, Vol. 49, pp. 5750-5754.
- Kwang-Geun, L. and Takayuki, S., 2002, "Determination of antioxidant potential of volatile extracts isolate from various herbs and spices", **Journal of Agricultural and Food Chemistry**, Vol. 50, pp. 4947-4952.
- Lachowicz, K.J., Jones, G.P., Briggs, D.R., Bienvenu, F.E., Wan, J., Wilcock, A., and Conventry, M.J., 1998, "The synergistic preservative effects of essential oils of sweet basil (*Ocimum basilicum* L) against acid-tolerant food microflora", **Letters in Applied Microbiology**, Vol. 26, pp. 209-214.
- Lambert, R.J.W., Skandamis, P.N., Coote, P.J., and Nychas, G.J.E., 2001, "A study of the minimum inhibitory concentration and mode of action of oregano essential oils, thymol and carvacrol", **Journal of Applied Microbiology**, Vol. 91, pp. 453-462.

- Lanciotti, R., Corbo, M., Gardini, F., Sinigaglia, M., and Guerzoni, M.E., 1999, "Effect of hexanal on shelf-life of fresh apple slices", **Journal of Agricultural and Food Chemistry**, Vol. 47, pp. 4769-4776.
- Lanciotti, R., Gianotti, A., Patrignani, F., Belletti, N., Guerzoni, M.E., and Gardini, F., 2004, "Use of natural aroma compounds to improve shelf-life and safety of minimally processed fruits", **Trends in Food Science and Technology**, Vol. 15, pp. 201-208.
- Laohakunjit, N., Kerdchoechuen, O., Norajit, K., and Boonsong, P., 2008, "Essential oil from five Zingiberaceae for anti Food-borne Bacteria", **International Food Research Journal**, Vol. 15, No. 3, pp. 337-346.
- Lee, J.C., Kim, H.R., Kim, J., and Jang, Y.S., 2002, "Antioxidant property of an ethanolic extract of the stem of *Opuntia ficus-indica* var. *saboten*", **Journal of Agricultural and Food Chemistry**, Vol. 50, pp. 6490-6496.
- Ly, T.N., Shimoyamada, M., Kato, K., and Yamauchi, R., 2003, "Isolation and characterization of some antioxidative compounds from the rhizomes of smaller galanga (*Alpinia officinarum* Hance)". **Journal of Agricultural and Food Chemistry**, Vol. 51, pp. 4924-4929.
- Mansour, E.H. and Khalil, A.H., 2000, "Evaluation of antioxidant activity of some plant extracts and their application to ground beef patties", **Food Chemistry**, Vol. 69, pp. 135-141.
- Nanasombat, S. and Teckchuen, N., 2009, "Antimicrobial, antioxidant and anticancer activities of Thai local vegetables", **Journal of Medicinal Plant Research**, Vol. 3, No. 5, pp. 443-449.
- Paris, R. and Nothis, A., 1969, "Sur quelques plants de nouvelle Caledonie", **Planttes Medicinales et Phytotherapie**, Vol. 4, pp. 274-287.
- Quattara, B., Simard, R.E., Piette, G., Begin, A., and Holley, R.A., 2000, "Inhibition of surface spoilage bacteria in processed meats by application of antimicrobial films prepared with chitosan", **International Journal of Food Microbiology**, Vol. 62, pp. 139-148.
- Rafi, M.M. and Vastano, B.C., 2007, "Identification of a structure specific Bcl-2-phosphorylating homoisoflavone molecule from Vietnamese coriander (*Polygonum odoratum*) that induced apoptosis and G2/M cell cycle arrest in breast cancer cell lines", **Food Chemistry**, Vol. 104, pp. 332-340.

- Richheimer SL, Bernart MW, King GA, Kent MC, Bailey DT (1996) Anti-oxidant activity of lipid-soluble phenolic diterpenes from rosemary. *J Am Oil Chem Soc* 73:507-514.
- Rizk, A., Hammouda, F., El-Nasr, S., and Abou-Youssef, A., 1980, "Constituents of Egyptian Euphorbiaceae. Part. VIII. Phytochemical investigation of *Euphorbia peplis*", *Filoterapia*, Vol. 51, pp. 223-227.
- Sara, B., 2004, "Essential oils: Their antibacterial properties and potential applications in foods: A review", **International Journal of Food Microbiology**, Vol. 94, pp. 223-253.
- Sekar, T. and Francis, K., 1998, "Some plant species screened for energy, hydrocarbons and phytochemicals", **Bioresouce Technology**, Vol. 65, pp. 257-259.
- Sekiwa, Y., Kubota, K., and Kobayashi, A., 2000, "Isolation of novel glucosides related to gingerdiol from ginger and their antioxidative activities", **Journal of Agricultural and Food Chemistry**, Vol. 48, pp. 373-377.
- Shahidi and Naczk, 1995. Food phenolics: Sources, Chemistry, Effects, Applications, Technomic Publishing Company Inc., Lancaster PA., pp: 231-245.
- Singhal, R.S., Kulkarni, P.R., and Rege, D.V., 2001, **Quality Indices for Spice Essential Oils**, Handbook of Herbs and Spices, CRC Press, USA, pp. 30.
- Singleton, V.L. and Rossi, J.A., 1965, "Colorimeter of total phenolics with phosphomolybdic-phosphotungstic acid reagent", **American Journal of Enology and Viticulture**, Vol. 16, pp. 144-158.
- Sivropoulou, A., Papanikolaou, E., Nikolaos, C., Kokkini, S., Lanaras, T., and Arsenakis, M., 1996, "Antimicrobial and cytotoxic activities of *Origanum* essential oils", **Journal of Agricultural and Food Chemistry**, Vol. 44, pp. 1202-1205.
- Skandamis, P., Tsigarida, E., and Nychas, G.J.E., 2002, "The effcet of oregano essential oil on survival/death of *Salmonella typhimurium* in meat stored at 5°C under aerobic, vp/map conditions", **Food Microbiology**, Vol. 19, pp. 97-103.
- Smid, E.J., Gorris, L.G.M., 1999. Natural antimicrobials for food preservation. In: Rahman, M.S. (Ed.), **Handbook of Food Preservation**. Marcel Dekker, New York, pp. 285–308.

- Smith-Palmer, A., Stewart, J., and Fyfe, L., 2001, "The potential application of plant essential oils as natural food preservatives in soft cheese", **Food Microbiology**, Vol. 18, pp. 463-470.
- Sokmen, A., Gulluce, M., Askin Akpulat, H., Daferera, D., Tepe, B., Polissiou, M., Sokmen, M. and Sahin, F., 2004, "The in vitro antimicrobial and antioxidant activities of the essential oils and methanol extracts of endemic *Thymus spathulifolius*", **Food Control**, vol. 15, pp. 627-634.
- Song, J., Deng, W., Fan, L., Verschoor, J., Beaudry, R., and Gonry, J.R., 1997, **Aroma volatiles and quality changes in modified atmosphere packaging**, In: Proceeding of "Fresh-cut fruits and vegetables and MAP symposium, Davis, USA, Vol. 5, pp. 19-25.
- Song, J., Leepipattanawit, R., and Beaudry, R., 1996, "Hexanal vapor is a natural, metabolizable fungicide: inhibition of fungal activity and enhancement of aroma biosynthesis in apple slice", **Journal of American Society of Horticultural Science**, Vol. 121, pp. 937-942.
- Starkenmann, C., Luca, L., Niclass,Y., Praz,E. and Roguet, D., 2006, "Comparison of Volatile Constituents of *Persicaria odorata* (Lour.) Soja'k (*Polygonum odoratum* Lour.) and *Persicaria hydropiper* L. Spach (*Polygonum hydropiper* L.)", **Journal of Agricultural and Food Chemistry**, Vol. 54, No. 8, pp. 3067-3071.
- Tepe, B., Sokmen, M., Askin Akpulat, H. and Sokmen, M., 2003, "In vitro antioxidant activities of the methanol extracts of five *Allium* species from Turkey", **Food Chemistry**, Vol. 192, pp. 89-92.
- Tepe, B., Daferera, D.; Sokmen, M., Polissiou, M., and Sokmen, A., 2004, "In Vitro Antimicrobial and antioxidant activities of the essential oils and various extracts of *thymus eigii*", **Journal of Agricultural and Food Chemistry**, Vol. 52, pp. 1132-1137.
- Tori, M., Ohara, Y., nakashima, K., Sono, M., 2000, "Caffeic and coumaric acid esters from *Calystegia soldanella*", **Filoterapia**, Vol. 71, pp. 39-44.
- Villalobos, M. and Correal, C., 1992, "The Euphorbiaceae as a source of vegetable oils for the chemical industry", **Grassy Aceites**, Vol. 43, pp. 39-44.

- Wettasinghe, M. and Shahidi, F., 1999, "Evening primrose meal: a source of natural antioxidants and scavenger of hydrogen peroxide and oxygen-derived free radicals", **Journal of Agricultural and Food Chemistry**, Vol. 47, pp. 1801-1812.
- Yuste, J., and Fung, D.Y.C., 2002, "Inactivation of *Listeria monocytogenes* Scott A 49594 in apple juice supplemented with cinnamon", **Journal of Food Protection**, Vol. 65, pp. 1663-1666.
- Zheng, W. and Wang,S.Y., 2001, "Antioxidant Activity and Phenolic Compounds in Selected Herbs", **Journal of Agricultural and Food Chemistry**, Vol. 49, pp. 5165-5170.