

Chromosome Study of Slender Walking Catfish (*Clarias nieuhofii*) and African Catfish (*C. gariepinus*) F₁ Hybrid: Revealed by Conventional Staining and Ag-NOR Banding Techniques

Samnao Saowakoon^{1*} and Nisa Machoo²

¹ Faculty of Agriculture and Technology, Rajamangala University of Technology Isan, Surin Campus, Muang, Surin 32000, Thailand

² Faculty of Agricultural Technology, Songkhla Rajabhat University, Tambon Khoa-Roob-Chang, Muang Songkhla 90000, Thailand

* Corresponding author: Saowakoon130713@gmail.com Tel: 0898467202

(Received: 3rd January 2021, Revised: 17th May 2021, Accepted: 19th May 2021)

Abstract - The hybrid catfishes *Clarias nieuhofii* (♀) × *C. gariepinus* (♂) were cytogenetically studied using conventional staining and Ag-NOR banding techniques, with the aim to reveal the chromosome complement and chromosome marker of the hybrid. Specimens were kindly provided by faculty of Agricultural Technology, Songkhla Rajabhat University, Songkhla, Thailand. Mitotic chromosomes were prepared from kidney follow that of the standard protocol. The result showed that hybrid catfishes had chromosome diploid of $2n=61$. An intermediate chromosomal pattern incorporating those of the parental species was found in the probable hybrid, confirming its interspecific origin. NORs were observed in one large submetacentric pair and one small subtelocentric pair. The number of diploid chromosome, chromosomal morphology and NOR locations are being cytogenetic evidence to prove that a hybrid occurred from *C. nieuhofii* × *C. gariepinus*.

Keywords: *C. nieuhofii*, *C. gariepinus*, hybrid catfish, karyotype

1. Introduction

Catfishes of the genus *Clarias* Scopoli, 1777 are characterized by their anguilliform body, long dorsal and anal fins, flattened bony head and a broad mouth with 4 pairs of barbels (Hee, 1990). The genus *Clarias* is the most diverse group 56 species within the moderately diverse Old World catfish family Clariidae (113 species in 16 genera) (Ferraris, 2007). Members of the genus are naturally distributed in inland water bodies in both Africa and Asia, with the bulk of the species being found in the former. Nineteen species of Southeast Asian *Clarias* are currently recognized (Ng, 2004). The African catfish (*C. gariepinus*) is considered to have a rapid growth rate (in length and weight), the rate of which strongly depends on ambient conditions and habitat (Britz & Pienaar, 1992; Bruton & Allanson, 1980; Hecht & Appelbaum, 1987). Growth has been found to be positively density dependent (Hecht and Appelbaum, 1987). Individuals have been recorded to reach 200 mm SL within a year (Bruton and Allanson, 1980; Skelton, 2001). In females, the growth rate decreases after 3 years resulting in the males reaching larger sizes (Skelton, 2001). Individuals of this species are known to live for eight or more years (Bruton & Allanson, 1980).

The slender walking catfish (*C. nieuhofii*) command a higher market price compared to other *Clarias*, because of its good taste and stunningly attractive appearance. It is beautifully patterned and eel-like in motion. It is reddish brown in color, with 13 or 14 vertical rows of white spots, which make it a valuable species for aquariums in Thailand. However, their populations and density have decreased significantly during the last 2 decades,

mainly due to overexploitation, reduction in habitat area as a result of the reclamation of peat swamp forests, and the injudicious application of insecticides in paddy fields. They have been observed to be rare, and were identified as a vulnerable species by IUCN red list categories and criteria (Pechsiri & Vanichanon, 2015). There have been more than 20 types of catfish hybrids attempted, hybrid catfish has been one of the most popularly culture freshwater fish species in Thailand (Srithong *et al.*, 2015).

In *Clarias* species examined to date, diploid chromosome numbers range between $2n=48$ and $2n=56$, with the exception of *C. pachynema*, *C. nieuhofii* ($2n=66$) and one population of *C. batrachus* ($2n=100$) (Table 1). In this study, we observed remarkable chromosomal dynamism and chromosomal characteristics by conventional staining and Ag-NOR banding techniques. The result obtained not only confirmed the possibility hybridization of *C. nieuhofii* × *C. gariepinus*, but the basic genetic information obtained is also be used for developing of other hybrid catfish in the future.

2. Materials and methods

2.1 Sample collection

The *C. nieuhofii* is a native species which inhabit in swamp forest, crossbreeding between *C. nieuhofii* female and *C. gariepinus* male. Ten F_1 hybrid catfishes (five males and five females) were collected from Aquaculture program, Faculty of Agricultural Technology, Songkhla Rajabhat University, Thailand. All specimens were maintained in aerated, flowing freshwater aquaria until analysis.

2.2 Chromosome preparation

Chromosomes were prepared *in vivo* (Phimphan *et al.*, 2013) as follows. Colchicine (0.1%) solution was injected into the fish's intramuscular and/or its abdominal cavity and then left for one hour. The kidneys were gently minced into small pieces in 0.075 M KCl and were then incubated for 30 min. After centrifugation at 1,200 rpm for 8 min, the cells suspension was fixed in freshly prepared fixative (3 methanol: 1 glacial acetic acid). The fixation step was repeated usually three times. After that the pellet was re-suspended with 1 ml of fixative. The cells were then dropped onto a clean slide and air-dried.

2.3 Chromosome staining

Conventional staining was done using 20% Giemsa's solution for 30 min. Ag-NOR banding (Howell & Black, 1980) was performed by adding four drops of 50% silver nitrate followed by two drops of 2% gelatin on slides. The slides were then covered with a cover slip and incubated at 60 °C for 5 min. The slides were then washed in distilled water. After air-dried, the slides were stained with 20% Giemsa's solution for 1 min. Mitotic metaphases were observed under a light microscope and were photographed.

2.4 Chromosome checks

The length of short arms (Ls) and long arms (Ll) of chromosomes were measured and calculated for the length of total arm chromosomes (LT, $LT = Ls + Ll$). Relative length (RL) and centromeric index (CI) were estimated. CI was also computed to classify the types of chromosomes according

to Turpin and Lejeune (1965). Classification of microchromosomes is based on the presence of a size between groups of large chromosomes and smaller chromosomes, all the microchromosomes are very small are often less than 1.5 micrometers (Ezaz & Young, 2013). All parameters were used in karyotyping and idiogramming.

This study was carried out in accordance with Ethical Principles and Guidelines for the Use of Animals, National Research Council of Thailand.

3. Results and Discussion

No differences of chromosome complement between male and female were observed in the hybrid catfishes (*C. nieuhofii* × *C. gariepinus*) (Figure 1C). Interestingly, from the metaphase chromosomes plates and karyotype (Figure 2 and 3), they showed a chromosome number of $2n=61$ and a fundamental number (NF; number of chromosome arm) of 112 which comprised 19 metacentric, 17 submetacentric, 15 subtelocentric and 10 acrocentric chromosomes. Although chromosomal morphology of hybrid is similar to that of parental species, the diploid number is obviously different i.e., *C. nieuhofii* (Figure 1A) and *C. gariepinus* (Figure 1B) had the diploid of $2n=66$ and 56, respectively (Table 1). Here, we proposed that the diploid chromosome number of the hybrid catfishes are basically assembled from the haploid set of gametes of *C. nieuhofii* ($n=33$) and *C. gariepinus* ($n=28$) resulting in the diploid chromosome number of $2n=61$ in the hybrid. A half of genome hybridization was also found in other catfish, the natural hybrid from *C. gariepinus* and *C. macrocephalus* ($2n=54$) which produced a new hybrid of $2n=55$ (Maneechot *et al.*, 2016).

Table 1 Review of available data on $2n$, karyotypes and sex systems in the genus *Clarias*.

Species	$2n$	NF	Karyotype	Sex system	Locality	Reference
<i>C. albopunctatus</i>	48	75	4m+23sm+21a	ZW	Nigeria	(Eyo, 2005)
	48	74	4m+22sm+22a	ZZ	Nigeria	(Eyo, 2005)
<i>C. anguillaris</i>	56	91	8m+27sm+21a	ZW	Nigeria	(Eyo, 2005)
	56	90	8m+26sm+22a	ZZ	Nigeria	(Eyo, 2005)
	48	-	27m+10sm+3st+8t	-	Nigeria	(Wu <i>et al.</i> , 1986)
	56	-	33m+12sm+2st+9t	-	Nigeria	(Wu <i>et al.</i> , 1986)
	100	111	4m+7sm+77a+12m	XY	Thailand	(Cui <i>et al.</i> , 1991)
<i>C. batrachus</i>	56	-	-	-	China	(Rishi, 1978)
	100	110	4m+6sm+78a+12mc	XY	Thailand	(Cui <i>et al.</i> , 1991)
	50	88	16m+8sm+14st+12a	-	India	(Prasad, 1998)
	50	96	18m+20sm+8st+4a	-	India	(Hinegarder & Rosen, 1972)
	54	-	-	-	-	(Pandey & Lakra, 1997)
	50	89	16m+11sm+5st+1at	ZW	India	(Nagpure <i>et al.</i> , 2000)
	50	88	16m+10sm+6st+18a	ZZ	India	(Nagpure <i>et al.</i> , 2000)
	51	89	16m+11sm+5st+18a+1B-chromosome	ZW	India	(Nagpure <i>et al.</i> , 2000)
	51	88	16m+10sm+6st+18a+1B-chromosome	ZZ	India	(Nagpure <i>et al.</i> , 2000)
	50	96	12m+18sm+10st+10t	-	India	(Siraj <i>et al.</i> , 2009)
	50	90	11m+7sm+2st+34a	-	Malaysia	(Yaqoob <i>et al.</i> , 2012)
	54	74	12m+18sm+10st+14t	-	India	(Ozouf-Costaz <i>et al.</i> , 1994)
	104	-	2m+4sm+98st/a	-	Thailand	(Maneechot <i>et al.</i> , 2016)
<i>C. camerunensis</i>	54	-	-	-	Africa	(Luo <i>et al.</i> , 1986)
	56	-	22m+20sm+9st+5t	-	Nigeria	(Wu <i>et al.</i> , 1986)
<i>C. ebriensis</i>	50	-	-	-	Africa	(Luo <i>et al.</i> , 1986)
	48	77	6m+23sm+19a	ZW	Nigeria	(Eyo, 2005)
	48	76	6m+22sm+20a	ZZ	Nigeria	(Eyo, 2005)
<i>C. fuscus</i>	56	106	18m+24sm+8st+6a	XX	China	(Cui <i>et al.</i> , 1991)
	56	106	19m+23sm+8st+6a	XY	China	(Cui <i>et al.</i> , 1991)
	56	106	20m+22sm+8st+6a	XX	China	(Yu <i>et al.</i> , 1989)
	56	106	20m+22sm+8st+6a	XY	China	(Yu <i>et al.</i> , 1989)
	56	102	18m+14sm+14st+10a	XX,XY	China	(Arai & Hirano, 1974)
	56	88	32m/sm+24st/a	-	Japan	(Luo <i>et al.</i> , 1986)
<i>C. gariepinus</i>	56	89	8m+25sm+23a	ZW	Africa, Israel	(Teugels <i>et al.</i> , 1992)

Table 1 Review of available data on $2n$, karyotypes and sex systems in the genus *Clarias*. (cont.)

Species	$2n$	NF	Karyotype	Sex system	Locality	Reference
	56	88	8m+24sm+24a	ZZ	Africa, Israel	(Teugels <i>et al.</i> , 1992)
	56	87	14m+17sm+25a	ZW	Egypt	(Siraj <i>et al.</i> , 2009)
	56	88	14m+18sm+24a	ZZ	Egypt	(Siraj <i>et al.</i> , 2009)
	56	102	20m+16sm+10st+10a	-	India	(Siraj <i>et al.</i> , 2009)
	56	89	8m+25sm+23a	ZW	Nigeria	(Eyo, 2005)
	56	88	8m+24sm+24a	ZZ	Nigeria	(Eyo, 2005)
	56	96	21m+14sm+5st+16a	-	Malaysia	(Yaqoob <i>et al.</i> , 2012)
	56	-	25m+14sm+14st+3t	-	Nigeria	(Wu <i>et al.</i> , 1986)
	56	100	28m+6sm+10a+12t	-	Turkey	(Omotayo, 2012)
	56	100	24m+10sm+10a+12t	-	Turkey	(Omotayo, 2012)
	54	98	34m+10sm+10t	-	Nigeria	(Awodiran <i>et al.</i> , 2014)
	56	102	6m+12sm+28st+10a	-	Nigeria	(Awodiran <i>et al.</i> , 2014)
	56	98	30m+6sm+6st+14t	-	Thailand	(Donsakul & Magtoon, 1990)
	56	110	18m+20sm+16st+2a	-	Thailand	(Maneechot <i>et al.</i> , 2016)
<i>C. jaensis</i>	54	-	22m+12sm+5st+15t	-	Nigeria	(Wu <i>et al.</i> , 1986)
<i>C. macrocephalus</i>	54	104	24m+20sm+6st+4a	-	Thailand	(Cui <i>et al.</i> , 1991)
	54	98	22m+18sm+4st+10a	-	Malaysia	(Yaqoob <i>et al.</i> , 2012)
	54	104	22m+16sm+12st+4a	-	Thailand	(Maneechot <i>et al.</i> , 2016)
<i>C. macromystax</i>	49	-	27 m+10sm+11st+1t	-	Nigeria	(Wu <i>et al.</i> , 1986)
<i>C. platycephalus</i>	54	-	-	-	Africa	(Luo <i>et al.</i> , 1986)
<i>C. pachynema</i>	66	-	30m+10sm+16st+10 t	-	Nigeria	(Wu <i>et al.</i> , 1986)
<i>C. nieuhoftii</i>	66	94	19m +10sm+8st+30a	-	Thailand	(Donsakul & Magtoon, 1992)

Notes: $2n$ = diploid chromosome, NF = fundamental number and - not available.




Figure 1. General characteristics of the *Clarias nieuhoftii* (♀), $2n=66$ (A), *C. gariepinus* (♂), $2n=56$ (B), hybrid catfish *C. nieuhoftii* × *C. gariepinus*, $2n=61$ (C). Scale bars indicate 5 cm.




Figure 2. Metaphase chromosome plate of hybrid fish [*Clarias nieuhofii* (♀) × *C. gariepinus* (♂)] $2n=61$ (A) by conventional staining and Ag-NORs banding (B). Arrows indicate nucleolar organizer regions (NORs). Bar = 5 µm.




Figure 3. Karyotype of hybrid fish [*Clarias nieuhofii* (♀) × *C. gariepinus* (♂)] $2n=61$ by conventional staining. Bar = 5 µm.
Note: un = unpaired chromosome.

In the present study, NOR bearing chromosomes were observed in two pairs of the hybrid, one is located in the short arm of large submetacentric pair and the other in the short arm of small subtelocentric pair (Figure 2B). For *C. nieuhofii*, the information of NOR has not yet been reported. However, one reported in *C. gariepinus* in Turkey has been revealed that NOR were observed on 3 or 4 chromosome pairs (Karahana &

Ergene, 2011). Moreover, the chromosomes of *C. gariepinus* have been mapped with 5S and 18S rDNA sequences compared to *C. macrocephalus* and their hybrid. Accordingly, the supposed hybrid of *C. gariepinus* × *C. microcephalus* had the intermediate number for both rDNA probes (Maneechot *et al.*, 2016). In the same side, to understand more about genomic hybridization relating to NOR of the hybrid

(*C. nieuhoftii* × *C. gariepinus*) and the individual parental species, the molecular cytogenetic approach such as rDNA sequences need to be performed in the future study.

4. Acknowledgments

This work was supported by Faculty of Agriculture and Technology, Rajamangala University of Technology, Isan Surin campus, Thailand.

5. References

- Arai, R. & Hirano, H. (1974). First record of the clariid catfish, *Clarias fuscus*, from Japan. *The Japanese Journal of Ichthyology*, 21(2), pp. 53–60. https://www.jstage.jst.go.jp/article/jji1950/21/2/21_2_53/_pdf.
- Awodiran, M.O., Majolagbe, F.A., Komolafe, O.O., Adewumi, A.A., Oyebola, O.O. & Oyewole, R.O. (2014). Cytogenetic study and serum protein characterization of *Clarias gariepinus* (Burchell, 1822) and *Heterobranchus bidorsalis* (Geoffroy Saint-Hillaire, 1809) in South western Nigeria. *International Journal of Biological Chemistry*, 8(6), 77–86. <https://www.ajol.info/index.php/ijbcs/article/view/116631>.
- Britz, P. J. & Pienaar, A. G. (1992). Laboratory experiments on the effect of light and cover on the behaviour and growth of African catfish, *Clarias gariepinus* (Pisces: Clariidae). *Journal of Zoology*, 227(1), 43–62. <http://doi/abs/10.1111/j.1469-7998.1992.tb04343.x>.
- Bruton, M. N. & Allanson, B. R. (1980). Growth of *Clarias gariepinus* in Lake Sibaya, South Africa. *African Zoology*, 15(1), 7–15. <https://doi.org/10.1080/02541858.1980.11447678>.
- Cui, J. X., Ren, X. H. & Yu, Q. X. (1991). Nuclear DNA content variation in fishes. *Cytologia*, 56(3), 425–429. <https://doi.org/10.1508/cytologia.56.425>.
- Donsakul, T. & Magtoon, W. (1990). Karyotypes of *Prophagorus nieuhoftii* (family clariidae), *Tachysurus caelatus* and *hemipimelodus borneensis* (family tachysuridae) from Thailand. *Proceedings of 41st Kasetsart University Annual Conference* (pp. 289–297).
- Donsakul, T. & Magtoon, W. (1992). A karyotype of two clariid catfishes, *Clarias gariepinus* and interspecific hybridization between *C. gariepinus* and *C. macrocephalus*. *Proceedings of the 17th Congress on Science and Technology of Thailand* (pp. 406–407).
- Eyo, J.E. (2005). Cytogenetic variations in *Clarias* species (Clariidae: Siluriformes) of the anambra river using leucocytes culture techniques. *Animal Research International*, 2(1), 275–86. <https://doi.10.4314/ari.v2i1.40852>.
- Ezaz, T. & Young, M. (2013). *Brenner's encyclopedia of genetics (2nd eds)*. Elsevier.

- Ferraris, C.J. (2007). Checklist of catfishes, recent and fossil (Osteichthyes: Siluriformes), and catalogue of siluriform primary types. *Zootaxa*, 1418(1), 1–628. <https://www.biotaxa.org/Zootaxa/article/view/zootaxa.1418.1.1/0>.
- Hecht, T. & Appelbaum, S. (1987). Notes on the growth of Israeli sharp-tooth catfish (*Clarias gariepinus*) during the primary nursing phase. *Aquaculture*, 63(1-4), 195–204. [https://doi.org/10.1016/0044-8486\(87\)90071-8](https://doi.org/10.1016/0044-8486(87)90071-8).
- Hee, H.N. (1999). Two new species of catfishes of the genus *Clarias* from Borneo (Teleostei: Clariidae). *Raffles Bulletin of Zoology*, 47(1), 17–32.
- Hinegarde, R. & Rosen, D.E. (1972). Cellular DNA content and the evolution of teleostean fishes. *The American Naturalist*, 106(951), 621–644.
- Howell, W.M. & Black, D.A. (1980). Controlled silver-staining of nucleolus organizer regions with a protective colloidal developer: A 1-step method. *Experientia*, 36, 1014–1015. <https://doi.10.1007/BF01953855>.
- Karahan, A. & Ergene, S. (2011). Chromosomal differentiation between populations of *clarias gariepinus* (Burchell, 1822) from the Goksu Delta and Orontes river (Turkey). *Turkish Journal of Biology*, 35(1), 79–87. <https://doi.10.3906/biy-0903-4>.
- Luo, J.L., Wang, Z.X., Lin, S.P. & Yang, J. H. (1986). Studies on the karyotype of *Clarias fuscus*. *Journal of Fisheries of China*, 10, 441–446.
- Maneechot, N., Yano, F.C., Bertollo, C.A. L., Getlekh, N., Molina, F.W., Ditcharoen, S., Tengjaroenkul, B., Supiwong, W., Tanomtong, A. & Cioffi, B.M. (2016). Genomic organization of repetitive DNAs highlights chromosomal evolution in the genus *Clarias* (Clariidae, Siluriformes). *Molecular Cytogenetics*, 9(4), 1-10. <https://doi.10.1186/s13039-016-0215-2>.
- Nagpure, N.S., Kushwah, B., Srivastava, S.K. & Ponniah, A.G. (2000). Comparative karyomorphology of African catfish *clarias gariepinus* (Burchell) and Asian catfish *clarias batrachus* (Linn.). *Chromosome Science*, 4(2), 57–59. <http://krishi.icar.gov.in/jspui/handle/123456789/5189>.
- Ng, H.H. (2004). *Clarias sulcatus*, a new walking catfish (Teleostei: Clariidae) from Pulau Redang. *Ichthyological Exploration of Freshwaters*, 15(4), 289–294.
- Omotayo, F. (2012). Karyological analysis of the African catfish, *Clarias gariepinus* (Burchell, 1822). Wudpecker. *Journal of Agricultural Research*, 1, 674–677.
- Ozouf-Costaz, C., Parrent, M., Teugels, G. G. and Van der Berg, E. (1994). Further karyological data on African clariid catfishes (Ostariophysi, Siluroidei). *Abstr. VIII Congr. Soc Europ Ichthyol* (pp. 56).

- Pandey, N. & Lakra, W.S. (1997). Evidence of female heterogamety, B-chromosome and natural tetraploidy in the Asian catfish, *Clarias batrachus*, used in aquaculture. *Aquaculture*, 149 (1-2), 31–37. [https://doi.org/10.1016/S0044-8486\(96\)01427-5](https://doi.org/10.1016/S0044-8486(96)01427-5).
- Pechsiri, J. & Vanichanon, A. (2015). Genetic diversity in slender walking catfish (*Clarias nieuhofii*) populations: implications for population management. *Walailak Journal Science & Technology*, 13(7), 511–519. <https://wjst.wu.ac.th/index.php/wjst/article/view/1690>.
- Phimphan, S., Tanomtong, A., Jumrusthanasan, S., Supiwong, W., Siripiyasing, P. & Sanoamuang, L. (2013). First report of NOR polymorphism and chromosome analysis of John's snapper, *Lutjanus johnii* (Perciformes, Lutjanidae) in Thailand. *Cytologia*, 78(4), 335–344. <https://doi.10.1508/cytologia.78.335>.
- Prasad, R. (1998). *Clarias batrachus*. In: Ponniah A.G & John, G. (eds). *Fish chromosome atlas* (pp. 214–215). National Bureau of Fish Genetic Resources.
- Rishi, K.K. (1978). *Chromosomal homogeneity in two Indian catfishes (5th edition)*. Geobios.
- Siraj, S.S., Sukardi, R., Phaik Imm, C.C., Mei, K.H., Vellasamy, S. & Panandam, J.M. (2009). Chromosome analysis of walking catfish, *Clarias* spp. *Malaysian Applied Biology Journal*, 38(1), 61–64. <http://psasir.upm.edu.my/id/eprint/13593>.
- Skelton, P. (2001). *A complete guide to the freshwater fishes of Southern Africa*. Struik Publishers.
- Srithong, C., Musig, Y., Areechon, N. & Taparhudee, W. (2015). Water quality and growth performance of hybrid catfish (*Clarias macrocephalus* x *C. gariepinus*) comparison in two type of water recirculating system and awater exchange system. *Journal of Fisheries and Environment*, 39(3), 57–68.
- Teugels, G.G., Guyomard, R. & Legendre, M. (1992). Enzymatic variation in African clariid catfishes. *Journal of Fish Biology*, 40, 87–96. <http://doi.10.1111/j.1095-8649.1992.tb02556.x>.
- Turpin, R. & Lejeune. (1965). *Les chromosomes humains (caryotype normal et variations pathologiques)*. Gauthier-Villars.
- Wu, G.M., Xin, C.B. & Chen, Y.L. (1986). A comparative study on the karyotypes of four species of catfish. *Acta Genetica Sinica*, 3, 213–20. http://en.cnki.com.cn/Article_en/CJFDTOTAL-YCXB198603008.htm.
- Yaqoob, S., Mohi-uddin, T. & Swaleha, T. (2012). Karyotypic analysis of walking catfish *Clarias batrachus* (Linnaeus, 1758). *Researcher*, 4(5), 46–48. http://www.sciencepub.net/researcher/research0405/009_9317research0405_46_48.pdf.
- Yu, X.J., Zhou, T., Li, Y.C., Li, K. & Zhou, M. (1989). *Chromosomes of Chinese freshwater fishes*. Science Press.