

## LIST OF FIGURES

Figure		Page
1	Schematic representation of a well known catalytic reaction, the synthesis of ammonia over iron catalysts.....	4
2	Crystal structure of $\text{LaCoO}_3$ showing (a) atomic positions of the unit cell and (b) structure depicting $\text{CoO}_6$ corner-sharing octahedra, with the A cations occupying the central interstice, and the B cations residing in the centre of the octahedra.....	5
3	Two types of oxygen: $\bullet$ adsorbed oxygen ( $\alpha$ -oxygen), $\circ$ lattice oxygen ( $\beta$ -oxygen).....	6
4	The various steps of a sol – gel process.....	9
5	Synthesis of basic salen structure.....	16
6	Simplified scheme for the formation of a solid product from solution.....	18
7	The precipitation conditions for a given compound.....	19
8	Qualitative comparison of the dependency of the rates of nucleation and growth of precipitate particle as a function of the solubility parameter (increasing from left to right) or T (opposite direction).....	20
9	Fundamental vibrations of several molecules.....	25
10	Schematic of an FTIR system.....	26
11	Schematic of a Michelson interferometer.....	27
12	Schematic representation of waves and their phases, input, output, and the two arms of the interferometer as the scan goes from zero path difference condition to $\text{OPD}=\lambda$ . (a) $\text{OPD}=0$ case. (b) $\lambda/4$ OPD case. (c) $\lambda/2$ OPD case. (d) $3\lambda/4$ OPD case. (e) $1\lambda$ OPD case.....	29
13	Two wavelength source case.....	30
14	Interferogram.....	30

## LIST OF FIGURES (continued)

Figure		Page
15	(a) an electron is ejected from ground state, (b) other electron from one of the upper levels falls to ground state. ....	35
16	Bragg diffraction.....	36
17	Diagram of a powder x-ray diffractometer.....	37
18	Powder diffraction file (JCPDF) No. 25-1060 of LaCoO <sub>3</sub> .....	38
19	Photoionization process occurs in x-ray photoelectron spectroscopy.....	41
20	Photoionization process and Auger process.....	42
21	Basic components of a monochromatic XPS system.....	43
22	X-ray fluorescence spectrometer.....	47
23	Different forms of performing vibrational spectroscopy: a) transmission infrared on a wafer pressed from catalyst powder and b) diffuse reflectance infrared spectroscopy on catalyst powder.....	49
24	(a) Mechanisms generating the infrared spectrum of a powder (b) Ellipsoidal mirror used in the collection of diffuse radiation from the sample and focus them on the detector.....	50
25	pH effect on precipitation of La(OH) <sub>3</sub> and Co(OH) <sub>2</sub> (● = [La <sup>3+</sup> ] and ○ = [Co <sup>2+</sup> ]).....	62
26	X-ray diffraction patterns of LaCoO <sub>3</sub> at different pH (* = LaCoO <sub>3</sub> , + = La(OH) <sub>3</sub> and # = Co <sub>3</sub> O <sub>4</sub> ) .....	63
27	X-ray diffraction patterns of α-Al <sub>2</sub> O <sub>3</sub> calcined at 1573 K.....	68
28	Calibration curve for determination of instrument broadening (β <sub>reference</sub> ).....	69

## LIST OF FIGURES (continued)

Figure		Page
29	Schematic of the system set up for catalytic activity testing: (A) mass flow controller, (B) vaporizer, (C) mixing coil and (D) DRIFT spectrometer.....	72
30	Cross-section of the vaporizer (B) in the Schematic of the system set up for catalytic activity testing.....	73
31	An environmental chamber.....	74
32	TG curve of the sample prepared by the Pechini method.....	77
33	FTIR spectra of the samples prepared by the Pechini method at 383 K (a) PN383-12, (b) La-PN383-12, (c) Co-PN383-12 and (d) LC-PN383-12.....	79
34	The proposed mechanism of the Pechini method.....	81
35	The proposed structure of the ester resulted from the reaction of citric acid and ethylene glycol.....	82
36	The complexation of the metal ions with the carboxylate in precursor prepared by the Pechini method.....	82
37	FTIR spectra of the samples prepared by the Pechini method at various temperatures.....	84
38	X-ray diffraction patterns of the samples prepared by the Pechini method and calcined at various temperatures (a) LC-PN573-03, (b) LC-PN773-08, (c) LC-PN823-08 and (d) LC-PN973-08.....	86
39	TG curve of the sample prepared by Schiff base complex method.....	88
40	FTIR spectra of the samples prepared by Schiff base complex method at 383 K (a) SB383-12, (b) La-SB383-12, (c) Co-SB383-12 and (d) LC-SB383-12.....	90

## LIST OF FIGURES (continued)

Figure		Page
41	The proposed structure of the resulted Schiff base.....	92
42	The complexation of the metal ions with the azomethine in precursor prepared by Schiff base complex.....	93
43	FTIR spectra of the samples prepared by Schiff base complex method at various temperatures.....	94
44	X-ray diffraction patterns of the samples prepared by Schiff base complex method and calcined at various temperatures (a) LC-SB573-03, (b) LC-SB773-08, (c) LC-SB823-08 and (d) LC-SB973-08.....	97
45	TG curve of the sample prepared by coprecipitation.....	99
46	FTIR spectra of the samples prepared by coprecipitation at various temperatures.....	101
47	X-ray diffraction patterns of the samples prepared by the coprecipitation and calcined at various temperatures (a) LC-CP773-03, (b) LC-CP823-03, (c) LC-CP873-03, (d) LC-CP973-03 and (e) LC-PN973-08 [ $\blacklozenge$ = La(OH) <sub>3</sub> ].....	102
48	La 3d photoelectron spectra of the samples prepared by the Pechini method.....	107
49	La 3d photoelectron spectra of the samples prepared by Schiff base complex method.....	108
50	O 1s photoelectron spectra of the samples prepared by the Pechini method.....	110
51	O 1s photoelectron spectra of the samples prepared by Schiff base complex method.....	111
52	Co 2p photoelectron spectra of the samples prepared by the Pechini method.....	114

## LIST OF FIGURES (continued)

Figure		Page
53	Co 2p photoelectron spectra of the samples prepared by Schiff base complex method.....	115
54	<i>In situ</i> infrared spectra of LC-PN773-08 at 423 K: (a) after pretreated 1 hour, (b) before start oxidation, (c) during oxidation reaction at 20 minutes and (d) after evacuation for 10 minutes.....	119
55	Absorbance intensity at wavenumber 2920 cm <sup>-1</sup> of the unconverted toluene relative to the starting toluene from oxidation at 423 K on ■ LC-PN573-03, ● LC-PN773-08, ▲ LC-PN823-08 as function of time.....	122
56	Absorbance intensity at wavenumber 2920 cm <sup>-1</sup> of the unconverted toluene relative to the starting toluene from oxidation at 423 K on □ LC-SB573-03, ○ LC-SB773-08, △ LC-SB823-08 as function of time.....	123
57	Absorbance intensity at wavenumber 2920 cm <sup>-1</sup> of the unconverted toluene relative to the starting toluene from oxidation at 393 K on ● LC-PN773-08 and ○ LC-SB773-08 as function of time.....	125
58	The proposed mechanism of toluene oxidation.....	128
 <b>Appendix Figure</b>		
A1	FTIR spectrum of Co(NO) <sub>3</sub> · 6H <sub>2</sub> O calcined at 773 K.....	140
A2	X-ray diffraction pattern of expansion of peak at d <sub>110</sub> .....	142
A3	X-ray diffraction pattern of expansion of peak at d <sub>024</sub> .....	142
B1	C 1s photoelectron spectra of the samples prepared by the Pechini method.....	143

## LIST OF FIGURES (continued)

Appendix Figure		Page
B2	C 1s photoelectron spectra of the samples prepared by Schiff base complex method.....	145
C1	DRIFTS spectra of flow rate comparison of toluene at (a) 50, (b) 70, (c) 80, (d) 90 and (d) 100 rpm adsorbed on LC-SB-773.....	147
D1	FTIR spectra of sample prepared by (I)Pechini and (II) Schiff base complex at various temperatures (a) ligand, (b) 383, (c) 423, (d) 473, (e) 523, (f) 573, (g) 623, (h) 773 for 3 h, (i) 773 for 8 h, (j) 823, (k) 873 and (l) 973 K.....	153
D2	XRD pattern of (I) LCO-PC and (II) LCO-SB calcined at 573 K for 3 h, 773 K, 823 K and 973 K for 8 h.....	154
D3	Absorbance intensity at wavenumber 2920 $\text{cm}^{-1}$ of the unconverted toluene relative to the starting toluene from oxidation at 423 K on (I) ■PC573, ●PC773 , ▲PC823 and (II) □SB573, ○SB773, △SB823 as a function of time.....	154
D4	Absorbance intensity at wavenumber 2920 $\text{cm}^{-1}$ of the unconverted toluene relative to the starting toluene from oxidation at 393 K on ● PC773 and ○ SB773 as function of time.....	155