

## LIST OF TABLES

Table		Page
1	Developmental period of <i>Platydema waterhousei</i> Gelbien reared on dried Ling-Zhi mushroom at $25.7\pm 4.9^{\circ}\text{C}$ , $56.5\pm 11.6\%$ rh and under a 12:12 (L:D) natural photoperiod.....	22
2	Regression equation of probit mortality (y) on logarithm of exposure time ( $\log_{10}$ ) for eggs, larvae, pupae, and adults of <i>Platydema waterhousei</i> Gelbien exposed to each type of modified atmospheres, at $25.7\pm 4.9^{\circ}\text{C}$ , $56.5\pm 11.6\%$ rh and under a 12:12 (L:D) natural photoperiod.....	30
3	The $\text{LT}_{50}$ and $\text{LT}_{99}$ values for eggs, larvae, pupae, and adults of <i>Platydema waterhousei</i> Gelbien when exposed to modified atmospheres, at $25.7\pm 4.9^{\circ}\text{C}$ , $56.5\pm 11.6\%$ rh and under a 12:12 (L:D) natural photoperiod.....	31
<b>Appendix Table</b>		
1	Comparison of developmental period (day) between male and female of <i>Platydema waterhousei</i> Gelbien at $25.7\pm 4.9^{\circ}\text{C}$ and $56.5\pm 11.6\%$ rh and under a 12:12 (L:D) natural photoperiod.....	50
2	First preliminary test to determine the optimal range of exposure period when eggs of <i>Platydema waterhousei</i> Gelbien exposed to four modified atmospheres, at $25.7\pm 4.9^{\circ}\text{C}$ , $56.5\pm 11.6\%$ rh and under a 12:12 (L:D) natural photoperiod.....	51
3	Second preliminary test to determine the optimal range of exposure period when eggs of <i>Platydema waterhousei</i> Gelbien exposed to four modified atmospheres, at $25.7\pm 4.9^{\circ}\text{C}$ , $56.5\pm 11.6\%$ rh and under a 12:12 (L:D) natural photoperiod.....	51

## LIST OF TABLES (Continued)

Appendix Table	Page	
4	First preliminary test to determine the optimal range of exposure period when larvae of <i>Platydema waterhousei</i> Gelbien exposed to four modified atmospheres, at $25.7 \pm 4.9^\circ\text{C}$ , $56.5 \pm 11.6\%$ rh and under a 12:12 (L:D) natural photoperiod.....	52
5	Second preliminary test to determine the optimal range of exposure period when larvae of <i>Platydema waterhousei</i> Gelbien exposed to four modified atmospheres, at $25.7 \pm 4.9^\circ\text{C}$ , $56.5 \pm 11.6\%$ rh and under a 12:12 (L:D) natural photoperiod.....	52
6	First preliminary test to determine the optimal range of exposure period when pupae of <i>Platydema waterhousei</i> Gelbien exposed to four modified atmospheres, at $25.7 \pm 4.9^\circ\text{C}$ , $56.5 \pm 11.6\%$ rh and under a 12:12 (L:D) natural photoperiod.....	53
7	Second preliminary test to determine the optimal range of exposure period when pupae of <i>Platydema waterhousei</i> Gelbien exposed to four modified atmospheres, at $25.7 \pm 4.9^\circ\text{C}$ , $56.5 \pm 11.6\%$ rh and under a 12:12 (L:D) natural photoperiod.....	53
8	First preliminary test to determine the optimal range of exposure period when adults of <i>Platydema waterhousei</i> Gelbien exposed to four modified atmospheres, at $25.7 \pm 4.9^\circ\text{C}$ , $56.5 \pm 11.6\%$ rh and under a 12:12 (L:D) natural photoperiod.....	54
9	Second preliminary test to determine the optimal range of exposure period when adults of <i>Platydema waterhousei</i> Gelbien exposed to four modified atmospheres, at $25.7 \pm 4.9^\circ\text{C}$ , $56.5 \pm 11.6\%$ rh and under a 12:12 (L:D) natural photoperiod.....	54

## LIST OF TABLES (Continued)

<b>Appendix Table</b>	<b>Page</b>
10	The corrected percentage mortality of <i>Platydema waterhousei</i> Gelbien eggs when exposed to four modified atmospheres, at 25.7±4.9°C, 56.5±11.6% rh and under a 12:12 (L:D) natural photoperiod.....
	55
11	The corrected percentage mortality of <i>Platydema waterhousei</i> Gelbien larvae when exposed to four modified atmospheres, at 25.7±4.9°C, 56.5±11.6% rh and under a 12:12 (L:D) natural photoperiod .....
	55
12	The corrected percentage mortality of <i>Platydema waterhousei</i> Gelbien pupae when exposed to four modified atmospheres, at 25.7±4.9°C, 56.5±11.6% rh and under a 12:12 (L:D) natural photoperiod.....
	56
13	The corrected percentage mortality of <i>Platydema waterhousei</i> Gelbien adults when exposed to four modified atmospheres, at 25.7±4.9°C, 56.5±11.6% rh and under a 12:12 (L:D) natural photoperiod.....
	56
14	The lethal time values for eggs of <i>Platydema waterhousei</i> Gelbien when exposed to G1 (100%CO <sub>2</sub> ) atmospheres, at 25.7±4.9°C, 56.5±11.6% rh and under a 12:12 (L:D) natural photoperiod.....
	57
15	The lethal time values for eggs of <i>Platydema waterhousei</i> Gelbien when exposed to G2 (80%CO <sub>2</sub> : 20%N <sub>2</sub> ) atmosphere, at 25.7±4.9°C, 56.5±11.6% rh and under a 12:12 (L:D) natural photoperiod.....
	58
16	The lethal time values for eggs of <i>Platydema waterhousei</i> Gelbien when exposed to G3 (60%CO <sub>2</sub> : 40%N <sub>2</sub> ) atmospheres, at 25.7±4.9°C, 56.5±11.6% rh and under a 12:12 (L:D) natural photoperiod.....
	59
17	The lethal time values for eggs of <i>Platydema waterhousei</i> Gelbien when exposed to G4 (100%N <sub>2</sub> ) atmosphere, at 25.7±4.9°C, 56.5±11.6% rh and under a 12:12 (L:D) natural photoperiod.....
	60

## LIST OF TABLES (Continued)

Appendix Table	Page	
18	The lethal time values for larvae of <i>Platydema waterhousei</i> Gelbien when exposed to G1 (100%CO <sub>2</sub> ) atmospheres, at 25.7±4.9°C, 56.5±11.6% rh and under a 12:12 (L:D) natural photoperiod.....	61
19	The lethal time values for larvae of <i>Platydema waterhousei</i> Gelbien when exposed to G2 (80%CO <sub>2</sub> : 20%N <sub>2</sub> ) atmospheres, at 25.7±4.9°C, 56.5±11.6% rh and under a 12:12 (L:D) natural photoperiod.....	62
20	The lethal time values for larvae of <i>Platydema waterhousei</i> Gelbien when exposed to G3 (60%CO <sub>2</sub> : 40%N <sub>2</sub> ) atmospheres, at 25.7±4.9°C, 56.5±11.6% rh and under a 12:12 (L:D) natural photoperiod.....	63
21	The lethal time values for larvae of <i>Platydema waterhousei</i> Gelbien when exposed to G4 (100%N <sub>2</sub> ) atmospheres, at 25.7±4.9°C, 56.5±11.6% rh and under a 12:12 (L:D) natural photoperiod.....	64
22	The lethal time values for pupae of <i>Platydema waterhousei</i> Gelbien when exposed to G1 (100%CO <sub>2</sub> ) atmospheres, at 25.7±4.9°C, 56.5±11.6% rh and under a 12:12 (L:D) natural photoperiod.....	65
23	The lethal time values for pupae of <i>Platydema waterhousei</i> Gelbien when exposed to G2 (80%CO <sub>2</sub> : 20%N <sub>2</sub> ) atmospheres, at 25.7±4.9°C, 56.5±11.6% rh and under a 12:12 (L:D) natural photoperiod.....	66
24	The lethal time values for pupae of <i>Platydema waterhousei</i> Gelbien when exposed to G3 (60%CO <sub>2</sub> : 40%N <sub>2</sub> ) atmospheres, at 25.7±4.9°C, 56.5±11.6% rh and under a 12:12 (L:D) natural photoperiod.....	67
25	The lethal time values for pupae of <i>Platydema waterhousei</i> Gelbien when exposed to G4 (100%N <sub>2</sub> ) atmospheres , at 25.7±4.9°C, 56.5±11.6% rh and under a 12:12 (L:D) natural photoperiod.....	68

## LIST OF TABLES (Continued)

<b>Appendix Table</b>	<b>Page</b>	
26	The lethal time values for adults of <i>Platydema waterhousei</i> Gelbien when exposed to G1 (100%CO <sub>2</sub> ) atmospheres, at 25.7±4.9°C, 56.5±11.6% rh and under a 12:12 (L:D) natural photoperiod.....	69
27	The lethal time values for adults of <i>Platydema waterhousei</i> Gelbien when exposed to G2 (80%CO <sub>2</sub> : 20%N <sub>2</sub> ) atmospheres, at 25.7±4.9°C, 56.5±11.6% rh and under a 12:12 (L:D) natural photoperiod.....	70
28	The lethal time values for adults of <i>Platydema waterhousei</i> Gelbien when exposed to G3 (60%CO <sub>2</sub> : 40%N <sub>2</sub> ) atmospheres, at 25.7±4.9°C, 56.5±11.6% rh and under a 12:12 (L:D) natural photoperiod.....	71
29	The lethal time values for adults of <i>Platydema waterhousei</i> Gelbien when exposed to G4 (100%N <sub>2</sub> ) atmospheres, at 25.7±4.9°C, 56.5±11.6% rh and under a 12:12 (L:D) natural photoperiod.....	72