

LIST OF FIGURES

Figure	Page
1 Structure of silica (a) A unit structure of silica containing SiO_4 unit in which one silicon is surrounded by four oxygen atoms in a tetrahedral geometry. (b) The expanded structure showing the coordination of oxygen atom between two silicon atoms.	3
2 Structure of amorphous silica (SiO_2).	4
3 Schematic diagram showing the various steps of a sol-gel process.	5
4 Sol-gel technologies and their products.	6
5 Electron microscopic view of silica nanotubes. (a) SEM image of silica nanotubes (1000 X). (b) TEM image of the cross section of the tubes.	7
6 TEM images of single silica nanotubes. (a) Low-magnification and (b) high-magnification	8
7 Electron microscopic view of silica products (a) TEM image of spherical silica particles obtained at $75\text{ }^\circ\text{C}$ (b) SEM image of silica tube formed at $0\text{ }^\circ\text{C}$ (c) TEM image of hollow silica tube obtained at $0\text{ }^\circ\text{C}$	9
8 TEM image of the final silica nanotubes synthesized from kaolin clay	10
9 TEM images of hollow silica particles: (a) spherical particles by cubic CaCO_3 template; (b) tubular particles by needle-like CaCO_3 template	11
10 FE-SEM images of the silica nanotubes having been annealed at $100\text{ }^\circ\text{C}$ for (a) 2h, and (b) 5 h after calcination and complete removal of the alumina template	12

LIST OF FIGURES (cont'd)

Figure	Page
11 Structure of molecule for self-assembly process (a) structure of <i>p</i> -dodecanoyl aminophenyl- β -D-glucopyranoside (b) structure of <i>p</i> -aminophenyl glucopyranoside and (c) a possible self-assembling model in the bilayered chiral fiber from the mixed gel of a and b	14
12 TEM images of the double-helical silica nanotube after calcinations (a and b), and (c) schematic representation of the double-helical structure of the silica nanotubes	14
13 The structure of product appended cholesterol (a) The structure of dimeric azobenzene and (b) SEM picture of the silica nanotube obtained from 1-octanol gel 1 after calcinations.	15
14 Schematic representation for the creation of the helical structure of the silica from the organogel state 1 .	15
15 Image of silica tubes from cocondensation method. (a) FE-SEM image. (b)FE-TEM image and (c) EDX spectrum of dye 1 prepared in 1-butanol	16
16 FE-SEM images illustrating the whisker-like morphology of the synthesized product. And small picture is a high-magnification image of the ends of two single hollow whiskers with a scale bar of 100 nm.	17
17 TEM images of nasnosilica particles ; 50,000X	18
18 Comparison the FTIR spectra of silica using tartaric acid template under shaking condition (SiTa_B)	25

LIST OF FIGURES (cont'd)

Figure	Page
19 FTIR spectra of calcined silica agitated by shaking condition. (a) without template and (b)-(h) with template	27
20 FT-IR spectra of calcined silica agitated by stirring condition. (a) without template and (b)-(h) with template	28
21 The powder XRD pattern of calcined silica using (a) without acid, (b) tartaric acid and (c) mucic acid as a template.	29
22 TG-DTA thermogram of uncalcined silica materials without template (SiW_B).	30
23 TG-DTA thermogram of uncalcined silica materials with mucic acid template (SiMu_B).	31
24 TG-DTA thermogram of uncalcined silica materials with succinic acid template (SiTa_B).	31
25 TGA thermogram of calcined silica materials (SiTa_B).	32
26 SEM images of silica materials from stirring method using organic acid templates	34
27 SEM images of silica materials from shaking method using organic acid templates.	35
28 TEM images of silica nanotubes that using tartaric acid as a template under stirring method	39
29 TEM images of silica nanotubes using mucic acid as a template under stirring condition (a) 4,000X and (b) 5,000X	40
30 TEM images of silica nanotubes using tartaric acid as a template under shaking condition	41

LIST OF FIGURES (cont'd)

Figure	Page
31 TEM images of silica nanotubes using mucic acid as a template under shaking condition (a) 4,000X, (b) 5,000X and (c and d) show EDS spectra of sphere and wall of silica products.	42
32 TEM images of silica materials using acid template under shaking condition	43
33 Propose mechanism for silica nanotubes formation using tartaric acid as template	45