

TABLE OF CONTENTS

	Page
TABLE OF CONTENTS	i
LIST OF TABLES	ii
LIST OF FIGURES	iv
INTRODUCTION	1
OBJECTIVES	3
LITERATURE REVIEWS	4
MATERIALS AND METHODS	15
RESULTS AND DISCUSSION	18
CONCLUSION	45
LITERATURE CITED	46
APPENDICE	56

LIST OF TABLES

Table	Page
1 Protein extracts in protein and non-protein parts from 30 grams of the recommended (KPS1 and CN36) and mutant lines (M5-16 and M5-29) seeds	18
2 Average number of egg of <i>C. maculatus</i> on KPS1 seeds treated with distilled water and protein part extracts of four mungbean variety/line seeds	21
3 Average number of egg of <i>C. maculatus</i> on KPS1 seeds treated with distilled water and non protein part extracts of four mungbean variety/line seeds	22
4 Percent larval mortality of <i>C. maculatus</i> on KPS1 seeds treated with distilled water and protein parts extracted from four mungbean variety/line seeds	26
5 Percent larval mortality of <i>C. maculatus</i> on KPS1 seeds treated with distilled water and non-protein parts extracted from four mungbean variety/line seeds	27
6 Percent pupal mortality of <i>C. maculatus</i> on KPS1 seeds treated with distilled water and protein parts extracted from four mungbean variety/line seeds	30
7 Percent pupal mortality of <i>C. maculatus</i> on KPS1 seeds treated with distilled water and non-protein parts extracted from four mungbean variety/line seeds	31
8 Percentages of adult mortality of <i>C. maculatus</i> from KPS1 seeds treated with distilled water and protein parts extracted from four mungbean variety/line seeds	34

LIST OF TABLES (continued)

Table		Page
9	Percentages of adult mortality of <i>C. maculatus</i> from KPS1 seeds treated with distilled water and non-protein parts extracted from four mungbean variety/line seeds	35

LIST OF FIGURES

Figure	Page
1 Percent inhibition at different concentrations of four mungbean crude extracts (protein part from intact seeds) against <i>Callosobruchus maculatus</i> α -amylase	40
2 Percent inhibition at different concentrations of four mungbean crude extracts (non- protein part from intact seeds) against <i>Callosobruchus maculatus</i> α -amylase	41
3 Percent inhibition at different concentrations of four mungbean crude extracts (protein part) against barley malt α -amylase (Type VIII-A)	42
4 Percentage inhibition at different concentrations of four mungbean crude extracts (non- protein part) against barley malt α -amylase (Type VIII-A)	44

LIST OF FIGURES (continued)

Appendix Figure	Page
1 The standard curve of Bovine Serum Albumin (BSA)	59
2 Amylase specific activity ($\mu\text{mol maltose min}^{-1} \text{ mg protein}^{-1}$) in the crude extracts of cowpea weevil <i>C. maculatus</i> performed at room temperature showing pH activity profiles at pH 2-12	60
3 Amylase specific activity ($\mu\text{mol maltose min}^{-1} \text{ mg protein}^{-1}$) in the crude extracts of cowpea weevil <i>C. maculatus</i> performed at 20-80°C	61
4 Amylase specific activity ($\mu\text{mol maltose min}^{-1} \text{ mg protein}^{-1}$) in the barley malt amylase (Type VIII-A) performed at room temperature showing pH activity profile at pH 2-12	62
5 Amylase specific activity ($\mu\text{mol maltose min}^{-1} \text{ mg protein}^{-1}$) in the barley malt amylase (Type VIII-A) performed at 20-80°C	63