

# Investigation of *Avian polyomavirus* and *Psittacine beak and Feather disease virus* in parrots in Taiwan

Sheng-Chang Chung<sup>1</sup> Ching-Dong Chang<sup>1</sup> Perng-Chih Shen<sup>2</sup>

Wei Lin<sup>3</sup> Chiu-Huang Ting<sup>1</sup> Hung-Yi Wu<sup>1\*</sup>

## Abstract

This study aimed to understand the related statistics of APV and PBFDV in parrots, such as the positive percentage and the distributions of various ages, seasons, breeds and areas, in order to obtain more information on the health care of captive parrots in Taiwan today. Blood samples were collected from parrots that were household pets and from commercial aviaries in the northern (e.g. Taipei), central (e.g. Taichung) and southern (e.g. Kaohsiung) Taiwan regions randomly. Collection targets were mainly small and medium species such as *Psittacula krameri* and others such as *Psittacus erithacus*. Sixty samples were collected every month and a total of 720 samples in a year. DNA was extracted and analyzed by polymerase chain reaction (PCR), sequencing and statistical analysis. The results showed that 28.3% (204/720) were PBFDV positive, 8.6% (62/720) were APV positive in this study and they were different from the two previous studies of Taiwan and studies of other countries such as the USA and Italy; however, APV/PBFDV co-infection (12%) was similar to the two previous studies of Taiwan (10.3% and 11.04%). The main species and age most susceptible were *Psittacula krameri* (14.1%, 48.7%), Lorikeets (9.8%, 19.5%), *Aratinga solstitialis* (7.5%, 28%), *Eclectus roratus* (6.3%, 22.9%) and chicks that were under six-months-old (3.6%, 17.8%). APV and PBFDV are easily spread during the breeding season due to frequent contact which makes the proportion of positives higher from commercial aviaries (8.7%, 29.7%) than that from household pets (8.3%, 21.7%), mainly in spring (13.3%, 44.4%) and fall (13.9%, 38.9%).

---

**Keywords:** PCR, APV, PBFDV, Psittacine Birds, Taiwan

<sup>1</sup>Department of Veterinary Medicine, College of Veterinary Medicine, National Pingtung University of Science and Technology, No. 1, Shuefu Road, Neipu, Pingtung, 91201 Taiwan

<sup>2</sup>Department of Animal Science, College of Agriculture, National Pingtung University of Science and Technology, No. 1, Shuefu Road, Neipu, Pingtung, 91201 Taiwan

<sup>3</sup>Department of Biological Science & Technology, National Chiao Tung University, No 1001, Daxue Rd., East Dist., Hsinchu, 30010 Taiwan

\*Correspondence: wuh@mail.npust.edu.tw (H. Wu)

Received: May 5, 2020.

Accepted: January 22, 2021.

doi: <https://doi.org/10.14456/tjvm.2021.31>

## Introduction

Avian polyomavirus (APV) is a 4981bp double-stranded circular DNA virus of the family *Polyomaviridae*, genus *Polyomavirus*; with non-enveloped, icosahedral symmetry, about 20-50 nm in diameter and 40-55 nm of virion size (Alley et al., 2013, Arroube et al., 2009, Rott et al., 1988). *Psittacine Beak and Feather Disease Virus* (PBFDV) is a 1993bp single-stranded circular DNA virus of the family *Circoviridae*, genus *Circovirus*; the virion is about 14-26 nm, with non-enveloped and icosahedral symmetry (Harkins et al., 2014). APV and PBFDV are common feather shedding and fatal diseases in parrots which cause acute or chronic infection. The severity and sensitivity in parrots depend on the species and age of the parrot (Johns and Muller, 2007, Sanada et al., 1999, Enders et al., 1997, Crawford, 1994, Ritchie et al., 1992, Latimer et al., 1991).

APV has now been reported in the United States, Canada, Japan, Africa, Australia, central Europe and other countries. The case fatality rate in parrot chicks is 20% to 100%. The first case in Taiwan was in 1987 (Johns and Muller, 2007, Enders et al., 1997, Phalen et al., 1991, Gough, 1989, Hirai et al., 1984, McOrist et al., 1984, Pass and Perry, 1984, Dykstra and Bozeman, 1982, Bernier et al. 1981) with a mortality rate of nearly 100% in budgerigars (Krautwald et al., 1989, Kaleta et al., 1984). In the case of non-budgerigar psittacine birds, except for chicks that are at higher risk, the mortality rate is about 27% to 41%, which is usually an inapparent infection that causes them to become carriers. (Graham and Calnek, 1987, Jacobson et al., 1984). The virus can be detected in all organs with a wide range of multiple organ infection capabilities (Phalen et al., 1993).

When APV infection is widespread all over the world, analysis shows that almost all infected cases have only one genotype and one serotype (Bert et al., 2005, Hsu et al., 2006, Khan et al., 2000, Phalen et al., 1999). Parrot species are more commonly infected with APV such as *Melopsittacus undulatus*, *Electus roratus*, Conures (genus *Aratinga/Pyrrhura*) and *Psittacula krameri*; while *Psittacus erithacus*, *Derophtus accipitrinus* and Amazon parrots have fewer cases and APV also has a record of infection in non-parrot birds, such as *Pteroglossus viridis*, *Pyrenestes sanguineus*, *Chloris chloris* and *Accipitriformes* (vultures, falcons) (Johns and Muller, 2007, Rossi et al., 2005, Lafferty et al., 1999, Garcia et al., 1994, Stoll et al., 1993, Forshaw et al., 1988).

PBFDV was first found in wild parrots in Australia. Due to the popularity of the parrot trade, transmission to the United Kingdom, Germany, Portugal, the United States, Spain, Israel, New Zealand, Zimbabwe and South Africa has been reported (Heath et al., 2004, McOrist et al., 1984, Pass and Perry, 1984). Acute infection by PBFDV occurs in parrots under three years old, usually at the first feather growth (about 28-32 days old) and some cause beak deformity (Sanada et al., 1999, Crawford, 1994, Cooper et al., 1987, McOrist et al., 1984, Pass and Perry, 1984).

PBFDV affects over 40 different psittacine species (Gerlach, 1994). According to the literature published in 1990, the clinical and pathological differences in parrot species that are known to be more susceptible to

PBFDV are only due to host factors rather than the differences in viral antigens and genes (or available variability) (Ritchie et al., 1990). However, this was not been confirmed until 1993 when Raidal et al., found that different virus strains can express the same antigen in blood clotting and blood clotting inhibition experiments (Raidal et al., 1993).

Primers have been designed to amplify a fragment of the capsid protein region encoded by the ORF V1 gene of PBFDV genome to detect PBFDV in psittacine birds (Ypelaar et al., 1999). Remarkable differences have been found between mammalian polyomaviruses and APV especially in the non-coding regulatory region and in the regions encoding the large tumor (T) antigen; therefore amplification of t/T antigen region of APV genome has been used and shown conservation in all APV-positive samples as expected (Dolz et al., 2013, Ogawa et al., 2005).

Reported viral DNA positive rates in different countries vary for PBFDV from 3.5% in the USA, 8.05% in Italy, 8.59% in Taiwan and 19.7% in Costa Rica to 40.4% in Germany and 41.2% in Taiwan, and for APV, from 0.79% in Italy and 4.8% in Costa Rica to 15.2% and 16.56% in Taiwan. Meanwhile, APV/PBFDV co-infections were 3.3% in Costa Rica to 10.3% and 11.04% in Taiwan (Thongchan et al., 2015, Dolz et al., 2013, Hsu et al., 2006, Bert et al., 2005, de Kloet E and de Kloet SR, 2004, Rahaus and Wolff, 2003).

This study aimed to understand the related statistics of APV and PBFDV in parrots such as the positive percentage and the distribution of various ages, seasons, breeds and areas, in order to obtain more information on the health care of captive parrots in Taiwan today.

## Materials and Methods

**Experiment Design:** Blood samples of parrots with suspected APV or PBFDV infection were collected over a year and the distributions of various ages, seasons, breeds and areas were recorded. After extracting sample DNA and obtaining target products by polymerase chain reaction (PCR) with specific primer pairs, the samples were then sent to the Clinical Molecular and Avian Medicine Laboratory, National Pingtung University of Science and Technology for sequencing and, finally, the above data integrated for statistical analysis. The parrot ages were divided into one-week-old to less than six-months-old, six-months-old to less than one-year-old, one-year-old to less than three-years-old, and three-years-old to less than ten-years-old as four groups; each group contained 180 parrots, making a total of 720 parrots. The seasons were divided into spring (February to April), summer (May to July), fall (August to October) and winter (November to January) as four groups; each group had 180 parrots making a total of 720 parrots. Collection targets were mainly small and medium species such as *Aratinga solstitialis*, *Psittacula krameri*, *Electus roratus* and Lorikeets (genus *Eos/Lorius/Trichoglossus*); others were *Psittacus erithacus*, *Melopsittacus undulatus*, *Nymphicus hollandicus*, genus *Pionites*, genus *Agapornis* and genus *Amazona*. The areas were divided into household pets and commercial aviaries.

**Samples:** From January 2019 to December 2019 for 12 months, blood samples of parrots were collected from household pets and commercial aviaries in the northern (Taipei, Keelung, Yilan, Taoyuan, Hsinchu), central (Taichung, Changhua, Nantou) and southern (Chiayi, Tainan, Kaohsiung, Pingtung) Taiwan randomly. Sixty samples were collected every month making a total of 720 samples in a year.

**Molecular biology tests:** Bymolecular biology testing, the main purpose was to extract sample nucleic acid (DNA) and use specific primer pairs to amplify the specific fragment size of the pathogen. Finally, the pathogen could be distinguished. The detected pathogens were *Avian Polyomavirus* and *Psittacine Beak and Feather Disease Virus*.

**Nucleic acid extraction:** The collected blood paper samples were shredded and an appropriate amount of sterile water was added, followed by shaking, so that the blood on the paper was dissolved in the sterile water. Then, sterile water mixed with the blood was used as whole blood, in conjunction with the Gene Plus™ Genomic DNA Extraction Miniprep System kit

(VIOGENE, GG2002) steps to purify DNA. The purified blood DNA was to be stored at -20 °C.

**Polymerase chain reaction assay:** The purified DNA was subjected to polymerase chain reaction(PCR) with specific primer pairs of a positive control of APV and PBFDV (Table 1). The APV primer pairs were 5'-CAAGCATATGTCCCTTTATCCC-3' and 5'-CTGTTT AAGGCCTTCCAAGATG-3' which were used for the detection of the t/T antigen coding region of APV. The expected PCR product was 310 bp (Phalen, *et al.*, 1991); the PBFDV primer pairs were 5'-AACCTACAGAC GGCGAG-3' and 5'-GTCACAGTCCTCCTTGACC-3' which were used to detect the PBFDV genome. Primers were designed from the ORF V1 gene sequence. The expected PCR product was 717 bp. (Ypelaar, *et al.*, 1999).

**Purification of PCR products:** After confirming the PCR product fragments using 1.5% Agarose gel, the Agarose gel of the target fragment was cut to an appropriate size for purifying target DNA using FavorPrep™ GEL / PCR Purification Mini Kit (FAVOGEN, FAGCK001-1). The purified target DNA had to be stored at -20°C until use.

**Table 1** Specific primer pairs of APV and PBFDV.

Name	Sequence (5'-3')	Region	Position	Size	References
APV	CAAGCATATGTCCCTTTATCCC CTGTTTAAGGCCTTCCAAGATG	t/T antigen	4303-4324 4612-4591	310bp	(Phalen, <i>et al.</i> , 1991)
PBFDV	AACCTACAGACGGCGAG GTCACAGTCCTCCTTGACC	ORF V1	182-199 898-879	717bp	(Ypelaar, <i>et al.</i> , 1999)

**Recombinant plasmids and colonies:** The purified target DNA was prepared using the RBC TA Cloning Vector Kit (RBC, RC001) to make DNA recombinant plasmids. After transforming in DH5α (*Escherichia coli*, *E. coli* DH5alpha, PTA-7297) competent cells, a single bacterial colony was obtained and scaled up. Using an RBC vector-specific primer pair to confirm the target fragment, the bacteria was then mixed with 30% glycerin and the mixture frozen at -80°C. Purifying the recombinant plasmids was done with FavorPrep™ Plasmid Extraction Mini Kit (FAVORGEN, FAPDE 100) and the pure recombinant plasmid DNA obtained.

**Sequencing:** The pure recombinant plasmid DNA was sent to Purigo Biotech, Inc. for sequencing. The primers used for sequencing were M13F(-40): 5'-CAGGGTTTTCCAGTCACGAC-3' and M13R(-48): 5'-AGCGGATAACAATTTACACAGG-3' (Table 2). After sequencing, the sequences were compared with data from the NCBI(National Center for Biotechnology Information, U.S. National Library of Medicine) GenBank.

**Table 2** The sequencing primers used in this study.

Primer name	Sequence (5'-3')	Source
M13F(-40)	CAG GGT TTT CCC AGT CAC GAC	Purigo Biotech, Inc.
M13R(-48)	AGC GGA TAA CAA TTT CAC ACA GG	Purigo Biotech, Inc.


## Results

**Results of PCR:** The Agarose gel electrophoresis showed that M was the Marker, which was the exact position of the positive control for determining the results of electrophoresis; the positive control of APV was located at 310bp and the positive control of PBFDV was located at 717bp (Figure 1).

**Sequence analysis:** The sequences were amplified by PCR which, after sequencing, were compared with

data from the NCBI GenBank to determine that they were specific nucleic acid fragments of APV and PBFDV and that there was no genetic variation.

**Total positive percentage:** A total of 62 of the 720 parrots (8.6%) were APV positive and a total of 204 of the 720 parrots (28.3%) were PBFDV positive. A total of 87 of the 720 parrots (12%) were APV/PBFDV co-infections (Table 3).



**Figure 1** Agarose gel electrophoresis showing the two PCR products of APV t/T antigen and PBFDV ORF V1 gene from the positive control. Lane 1 (M): DNA Molecular Weight Marker (GeneDireX, Inc., Taiwan), lane 2(A): 310bp from the positive control of APV, lane 3(P): 717bp from the positive control of PBFDV respectively.

**Table 3** Various positive percentages of APV and PBFDV in parrots.

Items	Sub-Items	Kinds of Virus and Various positive percentages				
		APV	PBFD	APV*	PBFD*	APV/PBFDV Co-infection*
Total		8.6% (62/720)	28.3% (204/720)	-	-	12% (32/266)
Age	1 week~<6 months	14.4% (26/180)	71.1% (128/180)	42% (26/62)	63% (128/204)	-
	6 months~<1 yr	10% (18/180)	26.7% (48/180)	29% (18/62)	23% (48/204)	-
	1~<3yrs	6.7% (12/180)	10% (18/180)	19% (12/62)	9% (18/204)	-
	3~<10yrs	3.3% (6/180)	5.6% (10/180)	10% (6/62)	5% (10/204)	-
	Spring	13.3% (24/180)	44.4% (80/180)	39% (24/62)	39% (80/204)	-
Season	Summer	4.4% (8/180)	15.6% (28/180)	13% (8/62)	14% (28/204)	-
	Autumn	13.9% (25/180)	38.9% (70/180)	40% (25/62)	34% (70/204)	-
	Winter	2.8% (5/180)	14.4% (26/180)	8% (5/62)	13% (26/204)	-
	Household pets	8.3% (10/120)	21.7% (26/120)	16% (10/62)	13% (26/204)	-
Area	Commercial aviaries	8.7% (52/600)	29.7% (178/600)	84% (52/62)	87% (178/204)	-
	<i>Aratinga solstitialis</i>	7.5% (16/214)	28% (60/214)	26% (16/62)	29% (60/204)	-
	<i>Psittacula krameri</i>	14.1% (22/156)	48.7% (76/156)	35% (22/62)	37% (76/204)	-
	<i>Eclectus roratus</i>	6.3% (6/96)	22.9% (22/96)	10% (6/62)	11% (22/204)	-
Breed	Lorikeets	9.8% (8/82)	19.5% (16/82)	13% (8/62)	8% (16/204)	-
	Other	5.8% (10/172)	17.4% (30/172)	16% (10/62)	15% (30/204)	-

\*Proportion of positive samples.

**Age distribution:** Of all samples, the positive percentages of APV from the four groups which were young to old in order were 14.4% (26/180), 10% (18/180), 6.7% (12/180) and 3.3% (6/180), while of PBFDV they were 71.1%(128/180), 26.7%(48/180), 10%(18/180) and 5.6%(10/180). In the positive

samples, the proportion of APV from the four groups which were young to old in order were 42%(26/62), 29%(18/62), 19%(12/62) and 10%(6/62), while of PBFDV they were 63%(128/204), 23%(48/204), 9%(18/204) and 5%(10/204). The proportion of APV positive and PBFDV positive in all samples or positive

samples were both highest in psittacine chicks that were under six-months-old and this decreased with age (Table 3).

**Season distribution:** Of all samples, the positive percentages of APV from spring to winter in order were 13.3% (24/180), 4.4% (8/180), 13.9% (25/180) and 2.8% (5/180), while of PBFVDV they were 44.4% (80/180), 15.6% (28/180), 38.9% (70/180) and 14.4% (26/180). In the positive samples, the proportion of APV positive from spring to winter in order were 39% (24/62), 13% (8/62), 40% (25/62) and 8% (5/62), while of PBFVDV positive they were 39% (80/204), 14% (28/204), 34% (70/204) and 13% (26/204). The highest proportions of APV positive and PBFVDV positive in all samples or positive samples were both in spring and fall (Table 3).

**Area distribution:** A total of 52 of 600 parrots (8.7%) of commercial aviaries and 10 of 120 parrots (8.3%) that were household pets were APV positive. A total of 178 of 600 parrots (29.7%) of commercial aviaries and 26 of 120 parrots (21.7%) that were household pets were PBFVDV positive. In positive samples, the proportion of APV was 84% (52/62) of commercial aviaries and 16% (10/62) of household pets; the proportion of PBFVDV was 87% (178/204) of commercial aviaries and 13% (26/204) of household pets. Of all or positive samples, the highest was in commercial aviaries (Table 3).

**Breed distribution:** Of all samples, the positive percentages of APV and PBFVDV were 7.5% (16/214) and 28% (60/214) in *Aratinga solstitialis*, 14.1% (22/156) and 48.7% (76/156) in *Psittacula krameri*, 6.3% (6/96) and 22.9% (22/96) in *Eclectus roratus*, 9.8% (8/82) and 19.5% (16/82) in Lorikeets (genus *Eos/Lorius/Trichoglossus*), 5.8% and 17.4% in other breeds, showing that the highest for APV and PBFVDV positive were in *Psittacula krameri*; the second-highest for APV and PBFVDV positive were in Lorikeets and *Aratinga solstitialis* respectively. In positive samples, the proportions of APV and PBFVDV were 26% (16/62) and 29% (60/204) in *Aratinga solstitialis*, 35% (22/62) and 37% (76/204) in *Psittacula krameri*, 10% (6/62) and 11% (22/204) in *Eclectus roratus*, 13% (8/62) and 8% (16/204) in Lorikeets and 16% (10/62) and 15% (30/204) in other breeds. The highest and second-highest for APV and PBFVDV in positive samples were both in *Psittacula krameri* and *Aratinga solstitialis* respectively (Table 3).

## Discussion

The positive percentage of PBFVDV in this study is 28.3%, which is higher than that of the United States, Italy and Costa Rica but lower than that of Germany. The positive percentage of APV in this study is 8.6%, which is higher than that of Italy and Costa Rica. In terms of APV/PBFVDV co-infection, 12% of this study were higher than Costa Rica. This may be related to environment, climate, feeding methods or bird species. The main object of this study is small and medium-sized parrots, which is different from the study of Costa Rica which was mainly of medium and large-sized parrots such as *Ara macao* and *Amazona*

*autumnalis* (Dolz et al., 2013, Bert et al., 2005, De Kloet E and De Kloet SR, 2004, Rahaus and Wolff, 2003).

Compared with the previous two studies of Taiwan, the PBFVDV positive percentage was the highest in 2002-2005 at 41.2% (Hsu et al., 2006) but it dropped to 8.59% from 2010 to 2014 (Thongchan et al., 2015) and rose again in 2019 to 28.3%. The APV positive percentage decreased significantly from 15.2% (Hsu et al., 2006) and 16.56% (Thongchan et al., 2015) to 8.6%, almost by half, which might have been caused by climate change or bird migration. However, the positive percentage of APV/PBFVDV co-infection in this study is 12%, compared with previous studies which were 10.3% (Hsu et al., 2006) and 11.04% (Thongchan et al., 2015), and there is no significant difference. The main species most susceptible to being APV and PBFVDV positive in this study were *Psittacula krameri*, Lorikeets, *Aratinga solstitialis*, *Eclectus roratus*, especially in chicks that were under six-months-old, which is the same as the susceptible species described in the literature (Thongchan et al., 2015, Johne and Muller, 2007, Hsu et al., 2006, Stoll et al., 1993).

Among them, the positive percentage of APV and PBFVDV was the highest in the *Psittacula krameri*, which could be related to the higher proportion of the rising numbers in Taiwan and the crowded feeding environment. In Taiwan, the breeding season for most parrots is mainly from late winter to spring, while lorikeets mainly breed in the fall. This shows a consistency with the results of this study with the greatest APV and PBFVDV infection mainly in spring and fall. The two viruses are easily spread during the breeding season when parrots are exposed to frequent contact, especially when young birds have poor immunity and are infected with viruses from the droppings of infected adult birds and the dust of feathers or feces (Ritchie et al., 1991, Gerlach, 1986). And this is the reason why a higher proportion of APV and PBFVDV infection is from commercial aviaries rather than that from household pets.

For feeding management, the birdhouse should be kept clean and attention paid to ventilation. If the parrots raised are found to be sick, the sick birds must be isolated and the environment must be disinfected.

## Acknowledgements

Special thanks are due to Dr. Hung-Yi Wu, Dr. Ching-Dong Chang, Dr. Perng-Chih Shen and Chiu-Huang Ting for outstanding technical support.

## References

- Alley MR, Rasiah I, Lee EA, Howe L and Gartrell BD 2013. Avian polyomavirus identified in a nestling Gouldian finch (*Erythrura gouldiae*) in New Zealand. *N Z Vet J.* 61: 359-361.
- Arroube AS, Halami MY, Johne R and Dorrestein GM 2009. Mortality due to polyomavirus infection in two nightjars (*Caprimulgus europaeus*). *J Avian Med Surg.* 23: 136-140.
- Bernier G, Morin M and Marsolais G 1981. A generalized inclusion body disease in the budgerigar (*Melopsittacus undulatus*) caused by a papovavirus-like agent. *Avian Dis.* 25: 1083-1092.

- Bert E, Tomassone L, Peccati C, Navarrete MG and Sola SC 2005. Detection of Beak and Feather Disease Virus (BFDV) and Avian Polyomavirus (APV) DNA in Psittacine Birds in Italy. *J. Vet. Med. B.* 52: 64-68.
- Bozeman LH, Davis RB, Gaudry D, Lukert PD, Fletcher OJ and Dykstra MJ 1981. Characterization of a papovavirus isolated from fledgling budgerigars. *Avian Dis.* 25: 972-980.
- Cooper JE, Gschmeissner S, Parsons AJ and Coles BH 1987. Psittacine beak and feather disease. *Vet Rec.* 120: 287.
- Crawford GR 1994. Circoviridae: new viruses of pigs, parrots and chickens. *Aust Vet J.* 71: 351.
- De Kloet E and De Kloet SR 2004. Analysis of the Beak and Feather Disease Viral Genome Indicates the Existence of Several Genotypes Which Have a Complex Psittacine Host Specificity. *Archive of Virology.* 149: 2393-2412.
- Dolz G, Sheleby-Eliás J, Romero-Zuñiga JJ, Vargas-Leitón B, Gutiérrez-Espeleta G and Madriz-Ordeñana K 2013. Prevalence of Psittacine Beak and Feather Disease Virus and Avian Polyomavirus in Captivity Psittacines from Costa Rica. *Open Journal of Veterinary Medicine.* 3: 240-245.
- Dykstra MJ and Bozeman LH 1982. A light and electron microscopic examination of budgerigar fledgling disease virus in tissue and in cell culture. *Avian Pathol.* 11: 11-28.
- Enders F, Gravendyck M, Gerlach H and Kaleta EF 1997. Fatal avian polyomavirus infection during quarantine in adult wild-caught red-faced lovebirds (*Agapornis pullaria*). *Avian Dis.* 41: 496-498.
- Forshaw D, Wylie SL and Pass DA 1988. Infection with a virus resembling papovavirus in Gouldian finches (*Erythrura gouldiae*). *AUST VET J.* 65: 26-28.
- Garcia A, Latimer KS, Niagro FD, Norton TM, Campagnoli RP, Harmon BG, Howerth EW and Ritchie RW 1994. Diagnosis of polyomavirus infection in seedcrackers (*Pyrenestes* sp.) and blue bills (*Spermophaga haematina*) using DNA in situ hybridization. *Avian pathology: journal of the W.V.P.A.* 23: 525-537.
- Gerlach H 1994. Circoviridae - psittacine beak and feather disease virus. In: Ritchie, B. W., G. T. Harrison, and L. R. Harrison (eds.). *Avian Medicine: Principles and Practice*, Wingers Publishing Inc., Lake Worth, FL: 894-903.
- Gerlach H 1986. Virus. In: G. J. Harrison and L. R. Harrison (eds.). *Clinical Avian Medicine and Surgery*, WB Saunders Co Philadelphia, London: 900-902.
- Gough JF 1989. Outbreaks of budgerigar fledgling disease in three aviaries in Ontario. *The Canadian veterinary journal. La revue veterinaire canadienne.* 30: 672-674.
- Graham DL and Calnek BW 1987. Papovavirus infection in hand-fed parrots: virus isolation and pathology. *Avian Dis.* 31: 398-410.
- Harkins GW, Martin DP, Christoffels A and Varsani A 2014. Towards inferring the global movement of beak and feather disease virus. *Virology.* 450-451: 24-33.
- Heath L, Martin DP, Warburton L, Perrin M, Horsfield W, Kingsley C, Rybicki EP and Williamson AL 2004. Evidence of unique genotypes of beak and feather disease virus in southern Africa. *J Virol.* 78: 9277-9284.
- Hirai K, Nonaka H, Fukushi H, Shimakura S, Masegi T and Mizoguchi T 1984. Isolation of a papovavirus-like agent from young budgerigars with feather abnormalities. *Nihon Juigaku Zasshi.* 46: 577-582.
- Hsu CM, Ko CY and Tsai HJ 2006. Detection and sequence analysis of avian polyomavirus and psittacine beak and feather disease virus from psittacine birds in Taiwan. *AVIAN DISEASES.* 50:348-353.
- Jacobson ER, Hines SA, Quesenberry K, Mladinich C, Davis RB, Kollias GV and Olsen J 1984. Epornitic of papova-like virus-associated disease in a psittacine nursery. *J Am Vet Med Assoc.* 185: 1337-1341.
- Johne R and Muller H 2007. Polyomaviruses of birds: etiologic agents of inflammatory diseases in a tumor virus family. *J Virol.* 81: 11554-11559.
- Kaleta EF, Herbst W, Kaup FJ, Jank-Ladwig R, Marschall HJ, Drommer W and Krautwald M 1984. Viral etiology of a disease accompanied by hepatitis and feather disorders in budgerigar fledgelings (*Melopsittacus undulatus*). *Zentralblatt fur Veterinarmedizin. Reihe B. Journal of veterinary medicine. Series B.* 31: 219-224.
- Khan MS, John R, Beck I, Pawlita M, Kaleta EF and Muller H 2000. Development of a blocking enzyme-linked immunosorbent assay for the detection of avian polyomavirus-specific antibodies. *J Virol Methods.* 89: 39-48.
- Krautwald ME, Muller H and Kaleta EF 1989. Polyomavirus infection in budgerigars (*Melopsittacus undulatus*): clinical and aetiological studies. *Zentralblatt fur Veterinarmedizin. Reihe B. Journal of veterinary medicine. Series B.* 36: 459-467.
- Lafferty SL, Fudge AM, Schmidt RE, Wilson VG and Phalen DN 1999. Avian polyomavirus infection and disease in a green aracarid (*Pteroglossus viridis*). *Avian diseases.* 43: 577-585.
- Latimer KS, Rakich PM, Steffens WL, Kircher IM, Ritchie BW, Niagro FD and Lukert PD 1991. A novel DNA virus associated with feather inclusions in psittacine beak and feather disease. *Veterinary pathology.* 28: 300-304.
- McOrist S, Black DG, Pass DA, Scott PC and Marshall J 1984. Beak and feather dystrophy in wild sulphur-crested cockatoos (*Cacatua galerita*). *J Wildl Dis.* 20: 120-124.
- Ogawa H, Yamaguchi T and Fukushi H 2005. Duplex Shuttle PCR for Differential Diagnosis of Budgerigar Fledgling Disease and Psittacine Beak and Feather Disease. *Microbiology and Immunology.* 49: 227-237.
- Pass DA and Perry RA 1984. The pathology of psittacine beak and feather disease. *Aust Vet J.* 61: 69-74.
- Phalen DN, Wilson VG, Gaskin JM, Derr JN and Graham DL 1999. Genetic diversity in twenty

- variants of the avian polyomavirus. *Avian diseases.* 43: 207-218.
- Phalen DN, Wilson VG and Graham DL 1993. Organ distribution of avian polyomavirus DNA and virus-neutralizing antibody titers in healthy adult budgerigars. *AM J VET RES.* 54: 2040-2047.
- Phalen DN, Wilson VG and Graham DL 1991. Polymerase chain reaction assay for avian polyomavirus. *J Clin Microbiol.* 29: 1030-1037.
- Rahaus M and Wolff MH 2003. Psittacine Beak and Feather Disease: A First Survey of the Distribution of Beak and Feather Disease Virus Inside the Population of Captive Psittacine Birds in Germany. *J. Vet. Med. B Infect. Dis. Vet. Public Health.* 50: 368-371.
- Ritchie BW, Niagro FD, Latimer KS, Steffens WL, Pesti D, Aron L and Lukert PD 1992. Production and characterization of monoclonal antibodies to psittacine beak and feather disease virus. *J Vet Diagn Invest.* 4: 13-18.
- Ritchie BW, Niagro FD, Latimer KS, Steffens WL, Pesti D, Ancona J and Lukert PD 1991. Routes and Prevalence of Shedding of Psittacine Beak and Feather Disease Virus. *Am J Vet Res.* 52: 1804-1809.
- Ritchie BW, Niagro FD, Latimer KS, Lukert PD, Steffens WL, 3rd, Rakich PM and Pritchard N 1990. Ultrastructural, protein composition, and antigenic comparison of psittacine beak and feather disease virus purified from four genera of psittacine birds. *J. of Wildlife Diseases.* 26: 196-203.
- Rossi G, Taccini E and Tarantino C 2005. Outbreak of avian polyomavirus infection with high mortality in recently captured Crimson's seedcrackers (*Pyrenestes sanguineus*). *J. of Wildlife Diseases.* 41: 236-240.
- Rott O, Kroger M, Muller H and Hobom G 1988. The genome of budgerigar fledgling disease virus, an avian polyomavirus. *Virology.* 165: 74-86.
- Sanada Y, Sanada N and Kubo M 1999. Electron microscopical observations of psittacine beak and feather disease in an Umbrella cockatoo (*Cacatua alba*). *J Vet Med Sci.* 61: 1063-1065.
- Stoll R, Luo D, Kouwenhoven B, Hobom G and Muller H 1993. Molecular and biological characteristics of avian polyomaviruses: isolates from different species of birds indicate that avian polyomaviruses form a distinct subgenus within the polyomavirus genus. *J Gen Virol.* 74: 229-237.
- Thongchan D, Huang YL, Khatri-Chhetri R, Tsai SS and Wu HY 2015. Prevalence and Pathology of Avian Polyomavirus (APV) and Psittacine Beak and Feather Disease Virus (PBFDV) Infections in Taiwan: A Retrospective Study from 2010 to 2014. *RMUTI JOURNAL Science and Technology.* 9: 1-13.
- Ypelaar I, Bassami MR, Wilcox GE and Raidal SR 1999. A universal polymerase chain reaction for the detection of psittacine beak and feather disease virus. *Vet Microbiol.* 68: 141-148.