



วารสารคณิตศาสตร์ **Mathematical Journal** 66 (705) กันยายน – ธันวาคม 2564

โดย สมาคมคณิตศาสตร์แห่งประเทศไทย ในพระบรมราชูปถัมภ์

<http://www.mathassociation.net>

Email: MathThaiOrg@gmail.com

อัตราส่วนจำนวนเต็มของอนุกรมที่เรียงติดกัน

Integer Ratios of Some Consecutive Series

เสฐฐวุฒิ เพชรแก้ว¹ และ สุภาวดี พฤษกาพิทักษ์^{2,*}

หน่วยวิจัยพีชคณิตและการประยุกต์ สาขาวิทยาศาสตร์การคำนวณ คณะวิทยาศาสตร์

มหาวิทยาลัยสงขลานครินทร์ สงขลา 90110

Sesthawuth Petchkaew¹ and Supawadee Prugsapitak^{2,*}

Algebra and Applications Research Unit, Division of Computational Science,

Faculty of Science, Prince of Songkla University, Songkhla 90110

Email: ¹6010210276@psu.ac.th, ²supawadee.p@psu.ac.th

วันที่รับบทความ : 7 มกราคม 2564

วันที่แก้ไขบทความ : 1 พฤษภาคม 2564

วันที่ตอบรับบทความ : 3 สิงหาคม 2564

บทคัดย่อ

ในบทความนี้ ผู้ประพันธ์ได้หาเงื่อนไขที่จำเป็นและเพียงพอที่ทำให้อัตราส่วนของอนุกรมที่เรียงติดกันต่อไปนี้เป็นจำนวนเต็ม โดยอนุกรมที่สนใจได้แก่

$$B_k(n) = \frac{1}{1(1+k)} + \frac{1}{2(2+k)} + \cdots + \frac{1}{n(n+k)} \quad (n, k \in \mathbb{N})$$

$$C_{a,d}(n) = \sum_{i=1}^n a + (n-i)d \quad (a, d, n \in \mathbb{N}) \text{ และ}$$

$$D_{a,r}(n) = a + ar + \cdots + ar^{n-1} \quad (a, n \in \mathbb{N}, r \in \mathbb{Q} \setminus \{0\})$$

คำสำคัญ: อนุกรม อนุกรมที่เรียงติดกัน

* ผู้เขียนหลัก

ABSTRACT

In this article, we establish necessary and sufficient conditions for the ratio of some consecutive series to be integers. The series that we investigate in the article are as follows:

$$B_k(n) = \frac{1}{1(1+k)} + \frac{1}{2(2+k)} + \cdots + \frac{1}{n(n+k)} \quad (n, k \in \mathbb{N}),$$

$$C_{a,d}(n) = \sum_{i=1}^n a + (n-i)d \quad (a, d, n \in \mathbb{N}) \text{ and}$$

$$D_{a,r}(n) = a + ar + \cdots + ar^{n-1} \quad (a, n \in \mathbb{N}, r \in \mathbb{Q} \setminus \{0\}).$$

Keywords: Series, Consecutive series

1. Introduction

The ratios of two consecutive powers were studied by many mathematicians [1, 3]. One of the unsolved problems is to find all positive integers k and $n > 1$ such that the ratio of the power sum $\frac{1^k + 2^k + 3^k + \cdots + n^k}{1^k + 2^k + 3^k + \cdots + (n-1)^k}$ is an integer. In 2019, Loulia N. Baoulina [2] studied a ratio for consecutive alternating power sums. Namely, he gave a necessary and sufficient condition for

$$\frac{1^k - 2^k + 3^k - \cdots + (-1)^{n+1} n^k}{1^k - 2^k + 3^k - \cdots + (-1)^n (n-1)^k} \quad (n, k \in \mathbb{N} \text{ and } n > 1)$$

to be an integer. From the mentioned result, it is natural to investigate the ratios for other series. In this paper, we establish necessary and sufficient conditions for the ratio of the following consecutive series to be integers:

$$B_k(n) = \frac{1}{1(1+k)} + \frac{1}{2(2+k)} + \cdots + \frac{1}{n(n+k)} \quad (n, k \in \mathbb{N}),$$

$$C_{a,d}(n) = \sum_{i=1}^n a + (n-i)d \quad (a, d, n \in \mathbb{N}) \text{ and}$$

$$D_{a,r}(n) = a + ar + \cdots + ar^{n-1} \quad (a, n \in \mathbb{N}, r \in \mathbb{Q} \setminus \{0\}).$$

2. Main Results

Theorem 2.1 Let k and n be positive integers. Then $\frac{B_k(n+1)}{B_k(n)}$ is never an integer

for all k and n .

Proof For positive integers k and n ,

$$\begin{aligned} B_k(n) &= \frac{1}{1(1+k)} + \frac{1}{2(2+k)} + \cdots + \frac{1}{n(n+k)} \\ &= \frac{1}{k} \left(\frac{1}{1} - \frac{1}{1+k} \right) + \frac{1}{k} \left(\frac{1}{2} - \frac{1}{2+k} \right) + \cdots + \frac{1}{k} \left(\frac{1}{n} - \frac{1}{n+k} \right) \\ &= \frac{1}{k} \left(1 - \frac{1}{1+k} + \frac{1}{2} - \frac{1}{2+k} + \cdots + \frac{1}{n} - \frac{1}{n+k} \right). \end{aligned}$$

Then

$$\begin{aligned} \frac{B_k(n+1)}{B_k(n)} &= 1 + \frac{\frac{1}{n+1} - \frac{1}{n+k+1}}{1 - \frac{1}{1+k} + \frac{1}{2} - \frac{1}{2+k} + \cdots + \frac{1}{n} - \frac{1}{n+k}} \\ &= 1 + \frac{\frac{1}{n+1} - \frac{1}{n+k+1}}{1 + \frac{1}{2} + \cdots + \frac{1}{n} - \frac{1}{k+1} - \frac{1}{k+2} - \cdots - \frac{1}{n+k}}. \end{aligned}$$

For $k=1$ and $n \geq 1$,

$$\begin{aligned} \frac{2}{n+1} &< 1 + \frac{1}{n+2} \\ \frac{1}{n+1} - \frac{1}{n+2} &< 1 - \frac{1}{n+1}. \end{aligned}$$

Since $\frac{1}{n+1} - \frac{1}{n+2} > 0$, $1 < \frac{B_1(n+1)}{B_1(n)} < 2$. Thus $\frac{B_1(n+1)}{B_1(n)}$ is not an integer.

For $k=2$ and $n \geq 1$,

$$\begin{aligned} \frac{1}{n+1} + \frac{1}{n+1} + \frac{1}{n+2} &< 1 + \frac{1}{2} + \frac{1}{n+3} \\ \frac{1}{n+1} + \left(\frac{1}{3} + \frac{1}{4} + \cdots + \frac{1}{n} \right) + \frac{1}{n+1} + \frac{1}{n+2} &< 1 + \frac{1}{2} + \left(\frac{1}{3} + \frac{1}{4} + \cdots + \frac{1}{n} \right) + \frac{1}{n+3} \end{aligned}$$

$$\frac{1}{n+1} - \frac{1}{n+3} < 1 + \frac{1}{2} + \frac{1}{3} \cdots + \frac{1}{n} - \frac{1}{3} - \frac{1}{4} - \cdots - \frac{1}{n+2}$$

Since $\frac{1}{n+1} - \frac{1}{n+3} > 0$, $1 < \frac{B_2(n+1)}{B_2(n)} < 2$. Thus $\frac{B_2(n+1)}{B_2(n)}$ is not an integer.

For $n \geq 2$ and $k \geq 2$, it suffices to show that

$$\frac{1}{n+1} + \left(\frac{1}{k+1} + \frac{1}{k+2} + \cdots + \frac{1}{k+n} \right) < 1 + \left(\frac{1}{2} + \frac{1}{3} + \cdots + \frac{1}{n} \right) + \frac{1}{n+k+1}.$$

We have

$$\begin{aligned} 2n+k+1 &< n^2+nk+n+k \\ (2n+k+1)(n+k+1) &< (n+1)(n+k)(n+k+2) \\ \frac{n+k+n+1}{(n+1)(n+k)} &< \frac{n+k+2}{n+k+1} \end{aligned}$$

Thus $\frac{1}{n+1} + \frac{1}{n+k} < 1 + \frac{1}{n+k+1}$. Since $\frac{1}{k+1} + \frac{1}{k+2} + \cdots + \frac{1}{k+n-1} < \frac{1}{2} + \frac{1}{3} + \cdots + \frac{1}{n}$

and $\frac{1}{n+1} + \frac{1}{n+k} < 1 + \frac{1}{n+k+1}$, our claim holds.

Thus $0 < \frac{1}{n+1} - \frac{1}{n+k+1} < 1 + \frac{1}{2} + \cdots + \frac{1}{n} - \frac{1}{k+1} - \frac{1}{k+2} - \cdots - \frac{1}{n+k}$ and this implies that $1 < \frac{B_k(n+1)}{B_k(n)} < 2$. Therefore $\frac{B_k(n+1)}{B_k(n)}$ is not an integer. \square

We next consider arithmetic series.

Theorem 2.2 Let a, d and n be positive integers. Then $\frac{C_{a,d}(n+1)}{C_{a,d}(n)}$ is an integer if

and only if one of the following holds:

1. $n=1$ and a divides d .
2. $n=2$ and $a=d$.

Proof For positive integers a, d and n , we have

$$\begin{aligned} \frac{C_{a,d}(n+1)}{C_{a,d}(n)} &= \frac{(a+nd) + (a+(n-1)d) + (a+(n-2)d) + \cdots + (a+d) + a}{(a+(n-1)d) + (a+(n-2)d) + (a+(n-3)d) + \cdots + (a+d) + a} \\ &= 1 + \frac{a+(n)d}{\frac{n}{2}(2a+(n-1)d)} = 1 + \frac{2a+2nd}{2na+nd(n-1)}. \end{aligned}$$

For $n=1$, we have $\frac{C_{a,d}(2)}{C_{a,d}(1)} = 1 + \frac{2a+2d}{2a} = 1 + \frac{a+d}{a} = 2 + \frac{d}{a}$.

Thus $\frac{C_{a,d}(2)}{C_{a,d}(1)}$ is an integer if and only if a divides d .

For $n=2$, we have $\frac{C_{a,d}(3)}{C_{a,d}(2)} = 1 + \frac{2a+4d}{4a+2d} = 1 + \frac{a+2d}{2a+d} = 2 + \frac{d-a}{2a+d}$.

We first consider the case $d > a$. Since $2a+d > d-a > 0$, $2a+d$ divides $d-a$ if and only if $a=d$.

Now we assume that $d \leq a$. If $2a+d$ divides $d-a$ then $d-a = m(2a+d)$ for some integer $m \leq 0$. Then $a = \frac{d(1-m)}{2m+1}$. Since $m \leq 0$ and $a, d > 0$, $2m+1 > 0$. Thus $m=0$ and this implies that $d=a$. For the converse, it is easy to see that if $d=a$ then $2a+d$ divides $d-a$.

For $n \geq 3$, $2na+nd(n-1) > 2a+2nd$. Therefore $\frac{C_{a,d}(n+1)}{C_{a,d}(n)}$ is never an integer. \square

We now consider geometric series $D_{a,r}(n)$.

Theorem 3 Let $a \in \mathbb{Z} \setminus \{0\}$, $r \in \mathbb{Q} \setminus \{0\}$ and $n \in \mathbb{N}$. Then $\frac{D_{a,r}(n+1)}{D_{a,r}(n)}$ is an integer if

and only if one of the following holds:

1. $n=1$ and $r=1$.
2. n is odd and $r=-1$.
3. $n=1$ and r is an integer.
4. $n=2$ and $r=-2$.

Proof We first consider the case $r = 1$ or $r = -1$.

If $r = 1$ then $\frac{D_{a,1}(n+1)}{D_{a,1}(n)} = \frac{(n+1)a}{na} = 1 + \frac{1}{n}$ which is an integer if and only if $n = 1$.

If $r = -1$ then we consider only the case that n is odd.

Since $D_{a,-1}(n+1) = 0$, $\frac{D_{a,1}(n+1)}{D_{a,1}(n)}$ is always an integer. Thus we next consider the nontrivial case as follows.

It is easy to see that if $n = 1$ and r is an integer then

$$\frac{D_{a,r}(n+1)}{D_{a,r}(n)} = \frac{a+ar}{a} = \frac{a(1+r)}{a} = 1+r$$

is an integer.

If $n = 2$ and $r = -2$ then $\frac{D_{a,r}(n+1)}{D_{a,r}(n)} = \frac{a+ar+ar^2}{a+ar} = \frac{1+r+r^2}{1+r} = -3$ is an integer.

Now let $a \in \mathbb{Z} \setminus \{0\}$, $r \in \mathbb{Q} \setminus \{0, \pm 1\}$ and $n \in \mathbb{N}$. Suppose $\frac{D_{a,r}(n+1)}{D_{a,r}(n)}$ is an integer.

Let $r = \frac{c}{d}$ where c and d are relatively prime.

$$\frac{D_{a,r}(n+1)}{D_{a,r}(n)} = 1 + \frac{\left(\frac{c}{d}\right)^n}{1 + \frac{c}{d} + \dots + \left(\frac{c}{d}\right)^{n-1}} = 1 + \frac{c^n}{d(d^{n-1} + cd^{n-2} + \dots + c^{n-1})}$$

Since c and d are relatively prime and $\frac{D_{a,r}(n+1)}{D_{a,r}(n)}$ is an integer, $d = 1$ and this

implies that r is an integer.

For $n = 1$, $\frac{D_{a,r}(2)}{D_{a,r}(1)} = \frac{a+ar}{a} = \frac{a(1+r)}{a} = 1+r$ is an integer.

For $n = 2$, $\frac{D_{a,r}(3)}{D_{a,r}(2)} = \frac{a+ar+ar^2}{a+ar} = 1 + \frac{ar^2}{a(1+r)} = 1 + \frac{r^2}{1+r} = r + \frac{1}{r+1}$ is an integer

if and only if $r = -2$.

For $n=3$, $\frac{D_{a,r}(4)}{D_{a,r}(3)} = \frac{a+ar+ar^2+ar^3}{a+ar+ar^2} = 1 + \frac{r^3}{1+r+r^2} = r + \frac{1}{r^2+r+1}$ is an integer

if and only if $r^2+r+1 = \pm 1$. Solving $r^2+r+1 = \pm 1$ gives us $r = \pm 1$ or $r = \frac{-1 \pm \sqrt{-7}}{2}$.

Both cases give us contradictions.

For $n \geq 4$, suppose $\frac{r^n}{1+r+\dots+r^{n-1}} = m$ for some integer m .

Thus $r^n = m(1+r+\dots+r^{n-1})$. If p is a prime divisor of r , p does not divide $1+r+\dots+r^{n-1}$. Thus p divides m . Since all prime divisors of r divides m and $\gcd(r, 1+r+\dots+r^{n-1}) = 1$, we have $r^n = \pm m$. Thus $1+r+\dots+r^{n-1} = \pm 1$ and this implies that $r+\dots+r^{n-1} = 0$ or $r+\dots+r^{n-1} = -2$. If $r+\dots+r^{n-1} = 0$ then $r(1+r+r^2+\dots+r^{n-2}) = 0$. Thus $r = 0$ or $1+r+r^2+\dots+r^{n-2} = 0$. The latter case implies that r divides ± 1 and this is a contradiction. If $r+\dots+r^{n-1} = -2$ then $r = \pm 1$ or $r = \pm 2$. Since $r \neq \pm 1$ and $r+\dots+r^{n-1} = -2 < 0$, $r = -2$. Thus $r+\dots+r^{n-1} = r$. So $r^2+\dots+r^{n-1} = 0$. Therefore $r = 0$ or $1+r+\dots+r^{n-3} = 0$. Since $n \geq 4$, the latter case implies that r divides 1 which is a contradiction. \square

Acknowledgements

We sincerely thank the referees for their valuable comments.

Reference

- [1] Baoulina, I. N., Moree, P. (2016). *Forbidden Integer Ratios of Consecutive Power Sums*. In: Sander, J., Steuding, J., Steuding, R., eds. *From Arithmetic to Zeta-Functions: Number Theory in Memory of Wolfgang Schwarz*, p. 1 - 30. Springer.
- [2] Baoulina, I. N. (2019). *Integer Ratios of Consecutive Alternating Power Sums*. *The American Mathematical Monthly*, 126 (7), p. 651 - 654.
- [3] Moree, P. (2013). *Moser's Mathematical Work on The Equation $1^k + 2^k + \dots + (m-1)^k = m^k$* . *Rocky Mountain J. Math.* 43 (5), p. 1707 - 1737.