<u>Table 10</u> Average of energy balance throughout 2002 year in Lampang teak plantation.

Season/month	Rs MJ/m [*] /day	energy balance (MJ m-2 day-1)				Bowen
		Rn	LE	Н	Gs	_ ratio
dry season						
November	20.1	10.7	7.5	2.7	0.5	0.4
December	19.1	12.9	9.0	3.3	0.6	0.4
January	23.2	12.5	8.8	3.1	0.6	0.4
February	21.7	12.9	9.0	3.3	0.6	0.4
March	23.3	17.7	11.4	5.4	0.9	0.5
April	25.1	18.9	11.2	6.8	0.9	0.6
average	22.1	14.3	9.5	4.1	0.7	0.4
% of Rs	100.0	64.6	-	-	-	-
% of Rn	-	100.0	66.5	28.7	4.8	3.0
Rainy season						
May	20.0	19.7	13.4	5.9	0.4	0.4
June	21.1	20.6	16.1	4.1	0.4	0.3
July	22.5	9.6	7.5	1.9	0.2	0.3
August	14.6	8.5	6.6	1.7	0.2	0.3
September	16.1	12.0	10.4	1.4	0.2	0.1
October	14.7	11.0	9.6	1.2	0.2	0.1
average	18.2	13.6	10.6	2.7	0.3	0.2
% of Rs	100.0	74.7	-	-	-	-
% of Rn		100.0	78.1	19.9	2.0	1.8
average throughout year	20.1	13.9	10.0	3.4	0.5	0.3
% of Rs	100.0	69.2	-	-	-	-
% of Rn	-	100.0	72.2	24.5	3.3	-

Remarks: Rs = solar radiation

Gs = heat storage in soil

LE = latent heat of vaporization

Rn = net radiation

Gw = heat storage in water

H = sensible heat