



Understanding the Impacts of Climate Change on Rice Production: A Qualitative Perspective

Muhamad Khairulbahri*

*Graduate Programme of Development Studies, Bandung Institute of Technology,
Bandung 40132, West Java, Indonesia*

Received 12 November 2020; Received in revised form 17 February 2021
Accepted 22 March 2021; Available online 28 March 2022

ABSTRACT

Several quantitative studies have concluded that climate change would negatively affect the Asian rice supply. Recently, as temperatures in Asia have approached the temperature threshold for rice growth, the Asian rice supply has become vulnerable to climate change in the near future. To complement quantitative studies' projections of the negative impacts of climate change on rice supply, this paper offers a qualitative analysis using system archetypes to understand the impacts of climate change on rice production. Two structures of the system archetypes are identified including Limits to Growth and Success to Successful. Both archetypes explain that rice production is hampered by high minimum temperature, as photosynthesis output is decreased by an increasing respiration. This paper shows that using this simple tool, system archetypes, we can describe the impacts of climate change on rice production. The outputs of this study, such as the causal loop diagram and system archetypes, can be the basis for the development of a simulation model used for understanding the impacts of climate change on major crops.

Keywords: Asian rice; Climate change; Global rice production; Rice production; System archetypes; System dynamics

1. Introduction

Assessing the impacts of climate change on crop production has attracted the interest of several scholars [1-5]. Worldwide, rice is arguably the most important crop, as it is the primary food source in Asia, where more than half of the world's population is located [6-8]. Rice is also a main source of employment and economic activity, as Asia

also accounts for about 90% of the world's rice production and total rice farming areas.

Despite its importance, farmed rice is considered highly vulnerable to climate change [6-8] as elevated temperatures and changing rainfall patterns are associated with climate change. As our society grows more concerned about the elevated CO₂ levels leading to climate change, the impacts on rice

*Corresponding author: databahri@yahoo.com

should be seen as a critical issue.

Rice should be sown in areas with temperatures within its optimum temperature range [9-11]. As temperature increases beyond its optimum temperature, rice yield tends to decline [10-12]. This means that rice yield tends to increase as temperature increases, but over its temperature threshold, yield tends to decrease [9-13]. Since rice farming areas in Asia have experienced temperatures of about 33°C [6, 10-11], climate change is likely to affect global rice production.

Furthermore, other studies [7, 10-11] have explained that rice is vulnerable to climate change since the recent maximum temperature and recent minimum temperature in regions of Asia are close to and higher than the temperature thresholds of rice, respectively. [11, 14].

Existing studies [11-12] have confirmed the negative impacts of climate change on Asian rice. Unfortunately, to the best of the author's knowledge, there is no study that offers a qualitative explanation of the impacts of climate change on rice. Thus, this study uses system archetypes for understanding the negative impacts of climate change on rice. This study would help policymakers in developing countries to understand the impacts of climate change on rice, through a concise qualitative analysis.

As this study aims to provide a qualitative analysis of the impacts of climate change on rice, it uses system archetypes to explain the impacts of change.

The system archetypes firstly were named by Senge [15] in his book "*The Fifth Discipline: The Art and Practice of the Learning Organization*". At the time this paper was written, there were about 10 archetypes such as Limits to Growth, Success to Successful, Shifting Burden, and Escalation.

System archetypes are recurring or regular patterns of complex systems that give insights into the system structures [15-16]. Moreover, system archetypes have been used

to analyze multiple types of observed systems, such as the water resource system [17], agriculture [18], the Australian electricity market [19], and the construction industry [20].

This suggests that system archetypes can be applied across a diverse range of subjects, scenarios, and contexts. In other words, system archetypes are generic patterns of systems or templates that represent usual practices in our world. Furthermore, Senge [15] points out that system archetypes enable us to identify leverage points which in turn can improve system performance. Further discussion of system archetypes can be found in many studies [17-19].

Since system archetypes are generic structures of systems that describe dynamic systems, system archetypes are a useful tool for analyzing dynamic and complex systems, such as the impacts of climate change on rice. Hence, system archetypes are a suitable tool for analyzing the impacts of climate change on rice production.

2. Methods

In the first step, causal links between two influential variables are defined. If a loop of causal links self-amplifies, then the loop is called a reinforcing loop (R). Otherwise, a loop of causal links is called a balancing loop (B). For convenience, all balancing and reinforcing loops in this study will be numbered.

In the following steps, all causal links consisting of influential variables are summarized as a causal loop diagram (CLD). Afterwards, from the CLD, identified system archetypes will be identified and explained. Based on identified system archetypes, how climate change influences rice production and the proposed solution in tackling the negative impacts of climate change will be explained.

To capture a deeper understanding of the underlying structures of the impacts of climate change on rice, system archetype

structures are identified as seen in the next section.

This study used Vensim Plus® to translate a conceptual model into a CLD. Then, system archetypes were identified based on the identified CLD.

3. Results and Discussion

3.1 A causal loop diagram of rice production

Rice growth is the product of the dynamic interaction between important variables such as carbon dioxide, precipitation, and temperature [3, 10, 21]. Crops absorb CO₂ to develop organic compounds supporting rice growth [10, 21]. Besides carbon dioxide, precipitation and temperature are important for rice production. Less precipitation leads to lower yield [21] previous studies [3, 10, 21] have also pointed out that climate change negatively impacts crops, as it relates to high minimum temperatures and high maximum temperatures.

The dynamic relationships between carbon dioxide, precipitation, and temperature are shown in Fig. 1. As climate change is associated with higher temperatures [2-3, 22], connecting lines between climate change and temperature [minimum and maximum temperature] have a “+” sign.

Opposite to this, the relationship between climate change and precipitation has a “-” sign, designating it a so-called balancing loop. This is supported by existing studies [22-23] that have projected Asia will experience lower seasonal precipitation. Climate change also has connecting lines with land-use change and fossil fuel burning [22], so fossil fuel burning and land-use change each have a “+” sign with “climate change”.




Fig. 1. Relationships among CO₂, climate change, temperatures, and precipitation.

Chemical substances containing phosphorus and nitrogen are important for rice growth [21, 24]. The main sources of these elements are fertilizers and water (from reservoirs or rainfall). Photosynthesis combines chemical substances and carbon dioxide to form the assimilation pool [21, 24]. The assimilation pool is important for supporting rice biomass, such as in leaves. Fig. 2 summarizes dynamic interactions between water, photosynthesis, and fertilizer.




Fig. 2. Photosynthesis.

Compounds within the assimilation pool are also converted to form roots, stems, leaves, and rice panicle, as seen in Fig. 3 [21, 24]. Rice panicle is important in supporting rice spikelet growth. More rice panicles lead to more rice spikelets on the top of the

panicles. As rice panicles will affect the volume of rice spikelets, the number of panicles per farming area can affect rice grain per panicle [21, 24].

Fig. 3 is a combination of Figs. 1-2. So, Fig. 3 combines the negative impacts of high minimum temperatures on rice yield (B1) and the growth engines that support photosynthesis and respiration (R1-R3). While R2 leads to increasing carbon dioxide owing to climate change, R3 supports more photosynthesis after fertilizer and carbon dioxide. As reinforcing loops, such as R2 and R3, lead to more resources, reinforcing loops are known as engine growths.

It should be acknowledged that two types of rice farming are practiced: rainfed rice and irrigated rice. While irrigated rice is supported by irrigation facilities, rainfed rice depends solely on rainfall. To accommodate the two types of rice farming, a link between precipitation and chemical substances is embedded in Fig. 3 to represent rainfed rice, while another link of precipitation-irrigated water-chemical substances represents irrigated rice.

This study also accommodates two types of rice varieties: japonica and indica. Although earlier studies [25-26] have acknowledged the different sensitivities of the two rice varieties to high temperature, recent studies [14, 25, 27] have confirmed the negative impacts of high temperature on both japonica and indica rice. With this point in mind, Fig. 3 applies to all rice regardless of variety.

Besides photosynthesis, crops such as rice also need respiration. Through respiration, rice can maintain ion concentration and lipid and protein turnover [21, 27]. If the minimum temperature exceeds a critical level, it tends to induce excessive respiration [14, 27]. Hence, excess respiration tends to decrease the size of the assimilation pool, leading to lower rice biomass and lower rice yield [27].

In general, there are two mechanisms by which higher temperatures negatively influence rice yield [28-29]. The first mechanism is by decreasing spikelet sterility, thus reducing rice yield [28-29]. However, transpirational cooling (more irrigated water) can negate this negative impact [28-29]. The second mechanism is by increasing respiration, thus reducing the assimilation pool size [27]. Dynamic relationships between transpirational cooling, high temperature, and assimilation pool can be seen in Fig. 4.

Fig. 5 is a CLD that shows the relationships between climate factors, non-climate factors, and rice growth. Fig. 5 also consists of some loops – both reinforcing and balancing loops.

The first reinforcing loop (**R1**) is a reinforcing loop that explains the dynamic interaction between photosynthesis, rice biomass, and assimilation pool [photosynthesis – assimilation pool – rice biomass – leaf – photosynthesis]. This loop (**R1**) is the growth engine supporting rice biomass development and the assimilation pool.




Fig. 3. Photosynthesis and respiration.

The second reinforcing loop (R2) is the growth engine of respiration, as high minimum temperature associated with climate change tends to induce excessive respiration (respiration – CO₂ – climate change – minimum temperature – respiration). The other growth engine is the third reinforcing loop (R3), describing the continuous photosynthesis upon the interaction between water and chemical substances (photosynthesis – CO₂ – climate change – precipitation – irrigated water – chemical substances – photosynthesis).

The first balancing loop (B1) explains an important relationship between respiration and rice biomass (rice biomass – respiration– assimilation pool – rice biomass). This means that more respiration leads to less rice biomass or lower rice yield. Another balancing loop (B2) is a “clearing mechanism” where CO₂ supports photosynthesis and in turn, photosynthesis decreases CO₂ availability.

It should be noted that rice-farming areas are an important emitter of methane (CH₄), associated with climate change [7-8]. The last balancing loop (B3) shows that less precipitation leads to decreased methane. Sustainable rice practices such as manure application and intermittent irrigated water also lead to reduced methane emission [7-8]. For simplification, the latter is not shown in Figure 5.

3.2 System Archetypes

3.2.1 The limits to growth

Based on the CLD, two system archetypes are identified. The structure of the limits to growth is the first system archetype as seen in Fig. 6. As seen in Figs. 7-8, the second type of system archetype present is the success to successful.

The structure of the limits to growth enhances photosynthesis to produce an assimilation pool and rice biomass (R1), leading to higher rice yield. Unfortunately,

the growth engine is negated by a balancing loop (B1). The balancing loop (B1) induces more respiration or decreases rice biomass, owing to the negative impacts of high minimum temperature. As a high minimum temperature is associated with climate change, changing climate tends to increase respiration, threatening rice yield.




Fig. 6. The limits to growth #1

Several studies [15-16] suggest possible leverage points to cope with negative impacts detected in the limits to growth structure. Heat-tolerant rice varieties are one leverage point to minimize the negative impacts of climate change on rice production. As heat-tolerant rice varieties may negate excess respiration, heat-tolerant varieties are likely to reduce the negative impacts of high minimum temperature. Mitigation of climate change, leading to low minimum temperature, may be another leverage point.

3.2.2 The success to successful

The success to successful structure is the second archetype, seen in Figs. 7-8. In this structure, one more successful activity will consequently gain more support, so the other activity ends with less support [15-16].

Fig. 7 shows two growth engines. The first growth engine (R3), photosynthesis, increases rice biomass, leading to higher rice production. The second growth engine (R2), high minimum temperature, leads to more respiration.

Developing heat-tolerant rice varieties is making promising progress, especially after some studies [34-36] confirmed the development of heat-tolerant genes.

4. Conclusion

Asian rice production, as a representation of global rice production, is threatened by climate change. The rising temperatures associated with climate change negatively affect rice production. At the same time, the benefit of elevated CO₂ is negated by the negative impacts of high minimum temperatures on respiration, ultimately leading to lower biomass and rice production.

The application of system archetypes as seen in this study can be used to understand the impacts of climate change on rice production and identify possible solutions. With the causal loop diagram, depicting interactions between climate and non-climate factors, archetype structures can be identified and explained. These archetype structures explain the dynamic interactions between the regulation of photosynthesis and the respiration process during rice growth. The negative impacts of climate change and the role of growth engines are described to capture the system archetype structures.

Through system archetype analysis, promising solutions under a changing climate are explained. While irrigation facilities offer a short-term solution, heat-tolerant rice varieties can be seen as a sustainable, long-term solution to balance the supply and demand of rice, as well as to increase rice production under a changing climate.

A quantitative model based on the causal loop diagram can be developed further to simulate the impacts of climate change on rice production as seen in another study [37]. The quantitative model may capture the dynamics of rice growth as well as the negative impacts of climate change on rice production. Moreover, possible solutions

can be derived from the system archetype structures explained in this study.

Another possible benefit taken from this study is that the information herein can serve as a basis to qualitatively explain the negative impacts of climate change on other major crops, such as wheat and corn. In the future, possible quantitative models can be developed for other major crops.

References

- [1] Green M, Weatherhead EK. Coping with climate change uncertainty for adaptation planning: An improved criterion for decision making under uncertainty using UKCP09. *Climate Risk Management*. 2014; 1:63-75.
- [2] Lobell DB, Burke MB, Tebaldi C, Mastrandrea MD, Falcon WP, Naylor RL. Prioritizing climate change adaptation needs for food security in 2030. *Science*. 2008;319[5863]:607-10.
- [3] Lobell DB, Schlenker W, Costa-Roberts J. Climate trends and global crop production since 1980. *Science*. 2011; 333[6042]:616-20.
- [4] Lobell D, Gourdj S. The Influence of Climate Change on Global Crop Productivity. *Plant Physiology*. 2012;160(4):1686-97.
- [5] Quiroga S, Iglesias A. A comparison of the climate risks of cereal, citrus, grapevine and olive production in Spain. *Agric Syst*. 2009; 101(1-2):91-100.
- [6] Chauhan BS, Jabran K, Mahajan G, editors. *Rice Production Worldwide*. Cham, Switzerland: Springer International Publishing; 2018.
- [7] Rice and its importance to human life Gnanamanickam SS. *Biological Control of Rice Diseases*, 2009.
- [8] Maclean J, Hardy B, Hettel G. *Rice Almanac: Source book for one of the most important economic activities on earth*. IRRI; 2013.

- [9] Shekhawat K, Rathore SS, Chauhan BS. Weed Management in Dry Direct-Seeded Rice: A Review on Challenges and Opportunities for Sustainable Rice Production. *Agronomy*. 2020; 10[9]:1264.
- [10] Wassmann R, Jagadish SV, Sumfleth K, Pathak H, Howell G, Ismail A, Serraj R, Redona E, Singh RK, Heuer S. Regional vulnerability of climate change impacts on Asian rice production and scope for adaptation. *Advances in agronomy*. 2009; 102:91-133.
- [11] Welch JR, Vincent JR, Auffhammer M, Moya PF, Dobermann A, Dawe D. Rice yields in tropical/subtropical Asia exhibit large but opposing sensitivities to minimum and maximum temperatures. *Proceedings of the National Academy of Sciences*. 2010; 107[33]:14562-7.
- [12] Wang J, Hasegawa T, Li L, Lam SK, Zhang X, Liu X, Pan G. Changes in grain protein and amino acids composition of wheat and rice under short-term increased [CO₂] and temperature of canopy air in a paddy from East China. *New Phytologist*. 2019; 222[2]:726-34.
- [13] Boonwichai S, Shrestha S, Babel MS, Weesakul S, Datta A. Evaluation of climate change impacts and adaptation strategies on rainfed rice production in Songkhram River Basin, Thailand. *Science of the Total Environment*. 2019; 652:189-201.
- [14] Peng S, Huang J, Sheehy JE, Laza RC, Visperas RM, Zhong X, Centeno GS, Khush GS, Cassman KG. Rice yields decline with higher night temperature from global warming. *Proceedings of the National Academy of Sciences*. 2004; 101[27]:9971-5.
- [15] Senge PM. *The fifth discipline fieldbook: Strategies and tools for building a learning organization*. Currency; 2014.
- [16] Maani KE, Cavana RY. *Systems thinking and modelling: Understanding change and complexity*. Prentice Hall; 2000.
- [17] Mirchi A, Watkins Jr D. A systems approach to holistic total maximum daily load policy: case of Lake Allegan, Michigan. *Journal of Water Resources Planning and Management*. 2013; 139[5]:544-53
- [18] Nguyen NC, Bosch OJ. A system thinking approach to identify leverage points for sustainability: a case study in the Cat Ba Biosphere Reserve, Vietnam. *Systems Research and Behavioral Science*. 2013; 30[2]:104-15.
- [19] Newell B, Marsh DM, Sharma D. Enhancing the resilience of the Australian national electricity market: Taking a systems approach in policy development. *Ecology and Society*. 2011; 16[2].
- [20] Guo BH, Yiu TW, González VA. Identifying behaviour patterns of construction safety using system archetypes. *Accident Analysis & Prevention*. 2015; 80:125-41.
- [21] Yoshida S. *Fundamentals of rice crop science*. Int. Rice Res. Inst.; 1981.
- [22] Field CB, Barros V, Stocker TF, Dahe Q, editors. *Managing the risks of extreme events and disasters to advance climate change adaptation: special report of the intergovernmental panel on climate change*. Cambridge University Press; 2012.
- [23] Levermann A, Schewe J, Petoukhov V, Held H. Basic mechanism for abrupt monsoon transitions. *Proceedings of the National Academy of Sciences*. 2009; 106(49):20572-7.
- [24] Li T, Angeles O, Marcaida III M, Manalo E, Manalili MP, Radanielson A, Mohanty S. From ORYZA2000 to ORYZA [v3]: An improved simulation model for rice in drought and nitrogen-deficient environments. *Agricultural and forest meteorology*. 2017; 237:246-56.
- [25] Prasad, P. V. V., et al. Species, ecotype and cultivar differences in spikelet fertility

- and harvest index of rice in response to high temperature stress. *Field crops research*, 2006, 95.2-3: 398-411.
- [26] Yoshida, Shouichi; Hara, Tetsuo. Effects of air temperature and light on grain filling of an indica and a japonica rice (*Oryza sativa* L.) under controlled environmental conditions. *Soil Science and Plant Nutrition*, 1977, 23.1: 93-107.
- [27] Mohammed AR, Tarpley L. Impact of high nighttime temperature on respiration, membrane stability, antioxidant capacity, and yield of rice plants. *Crop Science*. 2009; 49[1]:313-22.
- [28] Wassmann R, Dobermann A. Climate change adaptation through rice production in regions with high poverty levels. *J Semi-Arid Tropical Agric Res*. 2007; 4(1)
- [29] Jagadish SV, Sumfleth K, Howell G, Redoña E, Wassmann R, Heuer S. Temperature effects on rice: significance and possible adaptation. Advanced technologies of rice production for coping with climate change: 'no regret' options for adaptation and mitigation and their potential uptake. 2010:19-25.
- [30] Kim HY, Lieffering M, Kobayashi K, Okada M, Mitchell MW, Gumpertz M. Effects of free-air CO₂ enrichment and nitrogen supply on the yield of temperate paddy rice crops. *Field Crops Research*. 2003; 83(3):261-70.
- [31] Yang L, Huang J, Yang H, Zhu J, Liu H, Dong G, Liu G, Han Y, Wang Y. The impact of free-air CO₂ enrichment (FACE) and N supply on yield formation of rice crops with large panicle. *Field Crops Research*. 2006; 98(2-3):141-50.
- [32] Zhang G, Sakai H, Tokida T, Usui Y, Zhu C, Nakamura H, Yoshimoto M, Fukuoka M, Kobayashi K, Hasegawa T. The effects of free-air CO₂ enrichment (FACE) on carbon and nitrogen accumulation in grains of rice (*Oryza sativa* L.). *Journal of experimental botany*. 2013; 64(11):3179-88.
- [33] Zhang Y, Tang Q, Peng S, Zou Y, Chen S, Shi W, Qin J, Laza MR. Effects of high night temperature on yield and agronomic traits of irrigated rice under field chamber system condition. *Australian Journal of Crop Science*. 2013; 7(1):7.
- [34] Kilasi NL, Singh J, Vallejos CE, Ye C, Jagadish SV, Kusolwa P, Rathinasabapathi B. Heat stress tolerance in rice (*Oryza sativa* L.): Identification of quantitative trait loci and candidate genes for seedling growth under heat stress. *Frontiers in plant science*. 2018; 9:1578.
- [35] Tenorio FA, Ye C, Redoña E, Sierra S, Laza M, Argayoso MA. Screening rice genetic resources for heat tolerance. *SABRAO Journal of Breeding and Genetics*, 2013, 45.3: 371-81.
- [36] Ye C, Tenorio FA, Argayoso MA, Laza MA, Koh HJ, Redoña ED, Jagadish KS, Gregorio GB. Identifying and confirming quantitative trait loci associated with heat tolerance at flowering stage in different rice populations. *BMC genetics*. 2015;16(1):41.
- [37] Bahri M. Integrating Statistical and System Dynamics Modelling to Analyse the Impacts of Climate Change on Rice Production in West Nusa Tenggara, Indonesia. 2017. Unpublished thesis. Victoria University of Wellington