

Effect of N₂-fixing and IAA synthesis endophytic bacteria on growth of *Vanda* under greenhouse condition

Wanwisa Inkaewpuangkham¹, Kanokwan Panjama¹, Chaiartid Inkham², Yupa Chromkaew¹ and Soraya Ruamrungsri^{1*}

¹Department of Plant and Soil Science, Faculty of Agriculture, Chiang Mai University, Chiang Mai 50200, Thailand

²Plant Nutrition and Hydroponics Research Unit, Science and Technology Research Institute, Chiang Mai University, Chiang Mai 50200, Thailand

*Corresponding author: sorayaruumrung@gmail.com

Received: December 24, 2021. Revised: February 24, 2022. Accepted: March 9, 2022.

ABSTRACT

Vanda is one of the orchid genera that are important commercial crops. It was traded in both cut flower and plant form. In 2018, the *Vanda* export of Thailand had a marketable value of about 0.13% as cut flowers and 8.9% as orchid plants. Orchid cultivation is usually used a large number of fertilizers for growth and development, which lead to high cost and environmental problems. Therefore, this research was aimed to study N₂-fixing and IAA produced by endophytic bacteria as biofertilizers under greenhouse conditions. This research was conducted in a completely randomized design (CRD), and *Vanda* plantlets from tissue culture were inoculated with isolated 3S19 endophytic bacteria compared with deionized water (as a control treatment). Each treatment had twenty replications. The results showed that there were no significantly different between isolate 3S19 inoculation plants and with sterilized deionized water in plant height and leave number at 2, 3, and 4 months after inoculation. In contrast, root length and the number of roots were significantly different at 4 months after inoculation. The data from this research is beneficial as primary data for biofertilizer application to *Vanda* orchid.

Keywords: *Vanda*, orchid, endophytic bacteria, biofertilizer

INTRODUCTION

Orchids are an important commercial crop as an ornamental plant, potted plant, medical material, and industrial material (Hinsley et al., 2018). In 2018, Thailand exported cut flower and orchid plants for about 6.81 million US dollars (Department of International Trade Promotion, 2020). *Vanda* is one of the orchid exporting products with a market value of about 0.13% as cut flower form and 8.9% as an orchid plant (De et al., 2015; Panjama, 2018). Orchid cultivation has many factors to promote quality and yield, such as water, fertilizer, greenhouse, microorganism, etc. Generally, Thai farmers use a large number of fertilizers as water-soluble fertilizer forms, i.e. 20-20-20 or 21-21-21 (balance fertilizer) for the vegetative stage, 10-52-17 during floral stimulation, and 16-21-27 for growth and development (Department of Agriculture, 2017). The above information indicated that Thai growers used large amounts of fertilizers, resulting in high costs for orchid production. In addition, using chemical fertilizer leads to chemical accumulation in the environment because chemical fertilizer is slowly disintegrating (Ray et al., 2018). Micro-organisms, including bacteria or fungi, are one of the alternative products to promote plant growth through organic matter synthesis and are eco-friendly to the environment.

Endophytic bacteria is one micro-organism that lives inside the tissue of the plant and does not

cause plant disease to the host plant (Lacava and Azevedo, 2013). It can promote host plant growth and development through many mechanisms such as plant growth promoter substance synthesis, e.g., Indole-3-acetic acid (IAA), gibberellic acid, and cytokinin substance, promoting nutrient or water absorbance, nitrogen fixation, and biological control (Lacava and Azevedo, 2013).

Therefore, this research was aimed to study N₂-fixing and IAA synthesis by endophytic bacteria to promote *Vanda* growth under greenhouse conditions. This basic information may be useful for biofertilizer production in the future.

MATERIALS AND METHODS

This research was conducted with two treatments. *Vanda* plants were inoculated with isolate 3S19 at a ratio of 1:1 (endophytic bacterial suspension: sterilized water) compared with deionized water (as control treatment). There were 20 replications per treatment.

Preparing of endophytic bacteria suspension

The isolate 3S19 of endophytic bacteria, isolated from *Vanda* 'Manuvadee' which has the capability to fix nitrogen and synthesize IAA. (Inkaewpuangkham et al., 2021a; Inkaewpuangkham et al., 2021b). The assessment of nitrogen-fixing was conducted according to Weaver and Danso (Weaver and Danso, 1994),

by acetylene reduction assay and an IAA synthesis estimating according to a modified version of the Gordon and Weber method (Gordon and Weber, 1951). Firstly, the isolate 3S19 was cultured in a nutrient agar (NA) medium for seven days. Then, a single colony of isolate 3S19 was cultured in nutrient broth (NB) medium at 25 C° and shaken at 120 rpm for three days. Finally, endophytic bacteria suspension was estimated at an optical density of 0.5 at 600 nm (Lertjantarangkool et al., 2017) (approximately 10⁸ cfu/ml) before use.

Inoculation in *Vanda* plantlet

Vanda hybrid 6 months old plantlets from tissue culture were immersed in isolate 3S19 endophytic bacteria suspension compared with sterilized deionized water for 30 minutes. Then, inoculated plantlets were transferred to the basket and cultivated under greenhouse conditions for four months. The average temperature in the greenhouse was about 28°C and 70% RH.

Data collection

The data collection were plant height, number of leaves, root length, and number of roots at 2, 3, and 4 months after inoculation. The data was collected once times per month.

Statistical analyses

The data were analyzed by T-test independent using the Sxw program (Analytical Software version 8.0) with a significant difference at 95% probability level.

RESULTS

Growth of plantlet

The height of inoculated plantlets was not significantly different from control treatments at 2, 3, and 4 months after inoculation (Table 1).

Table 1. Height (cm.) of *Vanda* at 2, 3, and 4 months after inoculation with bacterial endophytes or sterilized deionized water (control).

Treatments	Months after inoculation (month)		
	2	3	4
Inoculated <i>Vanda</i>	8.3	8.5	8.6
Control plant	7.9	8.1	8.3
T-test	ns	ns	ns

ns = not significant

The number of leaves per plant of *Vanda* increased 4 months after inoculation. However, there was no significantly different between treatments at 2, 3, and 4 months (Table 2).

Table 2. Number of leaves of *Vanda* at 2, 3, and 4 months after inoculation with bacterial endophytes or sterilized deionized water (control)

Treatments	Months after inoculation (month)		
	2	3	4
Inoculated <i>Vanda</i>	4.2	4.5	5.1
Control plant	4.2	4.7	5.3
T-test	ns	ns	ns

ns = not significant

There was no significantly different in root length of *Vanda* between treatments at 2 and 3 months after inoculation. However, there was a significantly different root length between treatments at 4 months. The inoculation treatment with isolate 3S19 showed higher root length than the control plant (Table 3.).

Table 3. Root length (cm.) of *Vanda* at 2, 3, and 4 months after inoculation with bacterial endophytes or sterilized deionized water (control)

Treatments	Months after inoculation (month)		
	2	3	4
Inoculated <i>Vanda</i>	1.1	2.0	2.9 ^a
Control plant	1.0	1.8	2.2 ^b
T-test	ns	ns	*

Data analysis by T-test

ns = not significant

* = significant difference at 95% (P< 0.05)

At 2 and 3 months after inoculation, there was no significantly different between treatments. Nevertheless, inoculated *Vanda* with isolate 3S19 had more roots than control plants at 3.8 and 3.3 roots per plant, respectively (Table 4.).

Table 4. Number of roots of *Vanda* at 2, 3, and 4 months after inoculation with bacterial endophytes or sterilized deionized water

Treatments	Months after inoculation (month)		
	2	3	4
Inoculated <i>Vanda</i>	2.8	3.5	3.8 ^a
Control plant	2.9	3.2	3.3 ^b
T-test	ns	ns	*

Data analysis by T-test

ns = not significant

* = significant difference at 95% (P< 0.05)




Figure 1. Growth of *Vanda* plantlet after 4 months inoculation with isolate 3S19 (A), with sterilized deionized water (control) (B).

DISCUSSION

Plant height and leaf number of *Vanda* plantlet during 2-3 months after inoculation were not significantly different. It might be associated with host plant and bacteria interaction. The success of plant-bacteria associate depends on several factors. The condition of the environment is one factor for endophytic bacteria colonization, including season, temperature, latitude, longitude, nutrient, soil conditions, and so on (Chiellini et al., 2014; Yang et al., 2017; Wu et al., 2021). For example, the study of Ou et al. (2019) revealed that in the spring season, Actinobacteria and Proteobacteria of endophytic bacteria were found to be the most abundant in mulberry (*Morus L.*), but in the fall season found only Proteobacteria. In addition, the host plant is the main factor for endophytic bacteria colonization, consists of the plant growth stage, plant physiology, plant tissue type: Santos et al. (2018) showed that the plant tissue type leads to the different community of endophytic bacteria in host plant, plant species that similar Engelhard et al. (2000) reported that wild rice species and traditional rice had a population of *Azoarcus sp.* higher than modern varieties of rice since the influence of plant species. Likewise, the plant defense process is associated with plant-bacteria interaction: in some genera bacteria can induce plant defensive process that results in colonization difficult by other endophytic bacteria into the host plant (Santos et al., 2018). Furthermore, micro-organism species are an important factor for entry to plants. Research by Nogueira et al. (2001) reported that plant responses involve in plant-gene expression which was found associated with bacteria and a specific bacterial species.

Root length and number of roots of inoculated *Vanda* at 4 months after inoculation showed better growth than the control treatment. It might be because of abundant endophytic bacteria in the root organ. Typically, endophytic bacteria start entering into host plant via the roots because the root plant produces exudates to interact with bacteria. After endophytic bacteria entry to the root, they can spread to colonize above ground tissue by the movement of bacteria (bacterial flagella) and plant transpiration stream (Afzal et al., 2019). So, the 3S19 endophytic bacteria might be more abundant in roots than in other organs resulting in the root zone had more IAA synthesis by endophytic bacteria than the above zone. The property of isolate 3S19 could produce indole-3-acetic acid (IAA) without and with L-tryptophan medium at 9.48 and 17.38 mg IAA/l, respectively (Inkaewpuangkham et al., 2021a). Indicating that this isolate was active to produce IAA without exogenous tryptophan. Similarly, the previous report found that orchid roots are associated with fungal and bacterial symbionts, and they can synthesize IAA into orchids (Novak et al., 2014). Furthermore, a similar result was revealed by Khalid et al.

(2004) that microorganisms isolated from plant rhizospheric soil could synthesize IAA by using exudate containing tryptophan substrate from the plant root released. Since IAA is a plant growth regulator classified in the auxin group, it is predominant for promoting root formation or root initiation and enhancing root growth (Novak et al., 2014). Thus, isolating 3S19 could effectively promote root growth. Similar Tsavkelova et al. (2007) research reported that bacteria were isolated from root terrestrial orchids and epiphytic tropical orchids. These bacteria can produce IAA and stimulate root formation of kidney bean cutting.

The above reasons indicate that most bacteria often enter through roots and spread to the above plant part, and isolate 3S19 might have an intensive bacteria population in roots than leaves. So, isolate 3S19 might affect root growth than leaf growth in this stage of *Vanda*.

CONCLUSIONS

In conclusion, *Vanda* plantlets inoculated with isolate 3S19 of endophytic bacteria at a ratio of 1:1 (endophytic bacterial suspension: sterilized water) affected the growth of *Vanda* roots in both root length and the number of roots per plant than with sterilized deionized water inoculation. However, the data from this research is basic information to develop into a biofertilizer product. In the next step, the researcher will assess aspects of this isolate, such as phosphate solubilization and siderophore synthesis.

ACKNOWLEDGMENTS

This research was supported by the Department of Plant and Soil Science, Faculty of Agriculture, Chiang Mai University, Thailand, and H.M. The King's Initiative Centre for Flower and Fruit Propagation., Chiang Mai, Thailand, for their kind support.

REFERENCES

- Afzal, I., Shinwari, Z.K., Sikandar, S., and Shahzad, S. 2019. Plant beneficial endophytic bacteria: Mechanisms, diversity, host range and genetic determinants. *Microbiological Research*. 221: 36-49. <https://doi.org/10.1016/j.micres.2019.02.001>.
- Chiellini, C., Maida, I., Emiliani, G., Mengoni, A., Mocali, S., Fabiani, A., Biffi, S., Maggini, V., Gori, L., Vannacci, A., Gallo, E., Firenzuoli, F., and Fani, R. 2014. Endophytic and rhizospheric bacterial communities isolated from the medicinal plants *Echinacea purpurea* and *Echinacea angustifolia*. *International Microbiology*. 17: 165-174. <https://doi.org/10.2436/20.1501.01.219>.
- De, L., Pathak, P., Rao, A.N., and Rajeevan, P.K. 2015. *Commercial Orchids*. De Gruyter Open Ltd, Berlin.
- Department of Agriculture. 2017. *Orchid Production Technology*. Department of Agriculture, Ministry of Agriculture and Cooperatives, Bangkok.

- Department of International Trade Promotion. 2020. Orchid Products. Department of International Trade Promotion, Ministry of Commerce, Thailand.
- Engelhard, M., Hurek, T., and Reinhold-Hurek, B. 2000. Preferential occurrence of diazotrophic endophytes, *Azoarcus* spp., in wild rice species and land races of *Oryza sativa* in comparison with modern races. *Environmental Microbiology*. 2(2): 131-141. <https://doi.org/10.1046/j.1462-2920.2000.00078.x>
- Gordon, S. A., and Weber, R. P. 1951. The colorimetric estimation of indole acetic acid. *Plant Physiology*. 26: 192-195. <https://doi.org/10.1104/pp.26.1.192>.
- Hinsley, A., Boer, H.J.DE., Fay, M.F., Gale, S.W., Gardiner, L.M., Gunasekara, R.S., Kumar, P., Masters, S., Metusala, D., Roberts, D. L., Veldman, S., Wong, S., and Phelps, J. 2018. A review of the trade in orchids and its implications for conservation. *Botanical Journal of the Linnean Society*. 186: 435-455. <https://doi.org/10.1093/botlinnean/box083>.
- Inkaewpuangkham, W., Panjama, K., Inkham, C., Chromkaew, Y., and Ruamrungsri, S. 2021a. Isolation and Selection Potential N-Fixing of Endophytic Bacteria in *Vanda* Orchid. In: Sum-im, T. (ed) The 14th Research Administration Network Conference 2021 (online). Bangkok, 5-6 Aug 2021. Strategic Wisdom and Research Institute, Srinakharinwirot University. p.87-95.
- Inkaewpuangkham, W., Panjama, K., Inkham, C., Chromkaew, Y., and Ruamrungsri, S. 2021b. Assessment of IAA synthesis by endophytic bacteria in *Vanda* (Orchidaceae) (Book of Abstracts). In: Thammasiri, K., Kongsawadworakul, P., and Swangpol, S.C. (eds) The IX International Scientific and Practical Conference on Biotechnology as an Instrument for Plant Biodiversity Conservation (Physiological, Biochemical, Embryological, Genetic and Legal Aspects) (Biotech 2021). Faculty of Science, Mahidol University, Bangkok, Thailand, 12-13 Jul 2021. Department of Plant Science, Faculty of Science, Mahidol University. p.31.
- Khalid, A., Arshad, M., and Zahir, Z.A. 2004. Screening plant growth promoting rhizobacteria for improving growth and yield of wheat. *Journal of applied microbiology*. 96: 473-480. <https://doi.org/10.1046/j.1365-2672.2003.02161.x>.
- Lacava, P.T., and Azevedo, J.L. 2013. Chapter 1 Endophytic Bacteria: A Biotechnological Potential in Agrobiological System. In: Maheshwari, D.K., Saraf, M., and Aeron, A. (ed) *Bacteria in Agrobiological: Crop Productivity*. Springer, New York. p.1-44.
- Lertjantarangkool, T., Pengnoo, A., Punsak, W., and Homhaul, W. 2017. Effectiveness of free living nitrogen fixing bacteria on seed germination of rice. In: Wattanachaiyingcharoen, D., et al. (eds) Naresum research conference 13. Phitsanulok, 20-21 July 2017. Naresum University. P.359-367.
- Nogueira, E.M., Vinagre, F., Masuda, H.P., Vargas, C., Muniz de Pádua, V.L., Da Silva, F.R., Dos Santos, R.V., Baldani, J.I., Ferreira, P.C.G., and Hemerly, A.S. 2001. Expression of sugarcane genes induced by inoculation with *Gluconacetobacter diazotrophicus* and *Herbaspirillum rubrisubalbicans*. *Genetics and Molecular Biology*. 24: 199-206. <https://doi.org/10.1590/S1415-47572001000100027>.
- Novak, S.D., Luna, L.J., and Gamage, R.N. 2014. Role of Auxin in Orchid Development. *Plant Signaling & Behavior*. 9(10): e972277-1- e972277-8. <https://doi.org/10.4161/psb.32169>.
- Ou, T., Xu, W., Wang, F., Strobel, G., Zhou, Z., Xiang, Z., Liu, J., and Xie, J. 2019. A microbiome study reveals seasonal variation in endophytic bacteria among different Mulberry cultivars. *Computational and Structural Biotechnology Journal*. 17: 1091-1100. <https://doi.org/10.1016/j.csbj.2019.07.018>.
- Panjama, K. 2018. Nitrogen uptake and assimilation in *Vanda* hybrid. Ph.D. Dissertation. Graduate school Chiang Mai University, Chiang Mai, Thailand.
- Ray, S., Singh, J., Rajput, R.S., Singh, H.B., and Singh, S. 2018. Endophytic Bacteria: an Essential Requirement of Phyto Nutrition. *Nutrition and Food Science International Journal*. (5)2: 1-5. <https://doi.org/10.19080/NFSIJ.2018.05.555657>.
- Santos, M.L.dos., Berlitz, D.L., Wiest, S.L.F., Schünemann, R., Knaak, N., and Fiuza, L.M. 2018. Benefits Associated with the Interaction of Endophytic Bacteria and Plants. *Brazilian Archives of Biology and Technology*. 61: e18160431, 1-10. <https://doi.org/10.1590/1678-4324-2018160431>.
- Tsavkelova, E.A., Cherdyntseva, T.A., Botina, S.G., and Netrusov, A.I. 2007. Bacteria associated with orchid roots and microbial production of auxin. *Microbiological Research*. 162: 69-76. <https://doi.org/10.1016/j.micres.2006.07.014>.
- Weaver, R. W., and Danso, Seth K. A. 1994. Dinitrogen fixation. In: Weaver, R.W., Angle, J.S., and Bottomley, P.S. (eds.) *Methods of Soil Analysis. Part 2: Microbiological and Biochemical Properties*. Soil Science Society of America, Inc., Madison. p. 1019-1043.
- Wu, W., Chen, W., Liu, Shiyu., Wu, Jianjun., Zhu, Yeting., Qin, Luping., and Zhu, Bo. 2021. Beneficial Relationships Between Endophytic Bacteria and Medical Plants. *Frontiers in Plant Science*. 12: 1-13. <https://doi.org/10.3389/fpls.2021.646146>.
- Yang, R., Liu, P., and Ye, W. 2017. Illumina-based analysis of endophytic bacterial diversity of tree peony (*Paeonia* Sect. *Moutan*) roots and leaves. *Brazilian Journal of Microbiology*. 48: 695-705. <https://doi.org/10.1016/j.bjm.2017.02.009>.