



On Existence and Uniqueness of Common Fixed Point in C^* -Algebra Valued Metric Spaces

Deepak Kumar^{1,*}, Rishi Dhariwal²

¹Department of Mathematics, Lovely Professional University, Phagwara 144411, India

²Research Scholar, Lovely Professional University, Phagwara 144411, India

Received 5 September 2020; Received in revised form 15 April 2021

Accepted 2 February 2022; Available online 29 June 2022

ABSTRACT

In the present article, we prove some common fixed point theorems for two pairs of weakly compatible mappings satisfying rational type contractive condition in the framework of C^* -algebra valued metric spaces. The proved results extend and generalize some of the results in the literature.

Keywords: C^* -algebra valued metric space; Common fixed points; (CLR) property; (E.A.) property; Weakly compatible maps

1. Introduction and Preliminaries

The Banach contraction principle [1] is a very useful and effective tool of fixed point theory. The principle has numerous applications in pure as well as in applied sciences. For the last few decades, many researchers have investigated several types of mapping for the existence and uniqueness of a fixed point. (For reference see [2–13]). The principle is also useful for solving different kinds of equations such as integral, differential, fraction and partial differential equations (For reference see [14–19]).

Recently, Ma et al. [20] have ex-

tended Banach contraction principle to C^* -algebra valued metric spaces by replacing the set of real numbers with the set of all positive members of unital C^* -algebra. Afterward, many researcher have proved results in the framework of C^* -algebra valued metric spaces (For reference see [21–25]).

In the present manuscript, we prove some common fixed point theorems for two pairs of weakly compatible mappings satisfying rational type contractive condition in the framework of C^* -algebra valued metric spaces. The proved results extend and generalize some of the results in the literature.

For proving the results, we use some notation and definitions given in [23, 24].

A $*$ -algebra \mathbb{A} is a complex algebra with linear involution $*$ such that for all $\alpha, \beta \in \mathbb{A}$. $(\alpha, \beta)^* = \alpha^* \beta^*$ and $\alpha^{**} = \alpha$. The pair $(\mathbb{A}, *)$ is called a unital $*$ -algebra if it contains the unity element $I_{\mathbb{A}}$. If a unital $*$ -algebra satisfy $\|\alpha^*\| = \|\alpha\|$, for all $\alpha \in \mathbb{A}$, then \mathbb{A} is called Banach $*$ -algebra. A Banach C^* -algebra satisfying $\|\alpha^* \alpha\| = \|\alpha\|^2$, for all $\alpha \in \mathbb{A}$ is called C^* -algebra. If $\alpha = \alpha^*$ and $\sigma(\alpha) = \gamma \in \mathcal{R} : \beta I_{\mathbb{A}} - \alpha$ is non-invertible, α is called the positive element of \mathbb{A} . If $\alpha \in \mathbb{A}$ is positive, we write it as $\alpha > 0_{\mathbb{A}}$. The partial ordering on \mathbb{A} can be defined as follows: $\alpha > \beta$ if and only if $\alpha - \beta > 0_{\mathbb{A}}$. Throughout the paper, by \mathbb{A} , we denote a unital C^* -algebra with the unity element $I_{\mathbb{A}}$.

Definition 1.1 ([20]). Suppose X is a non-empty set. The mapping $d : X \times X \rightarrow \mathbb{A}$ is called a C^* -algebra valued metric on X if it satisfies :

- (i) $d(p, q) \geq 0_{\mathbb{A}}$ and $d(p, q) = 0_{\mathbb{A}}$ iff $p = q$;
- (ii) $d(p, q) = d(q, p)$;
- (iii) $d(p, r) \leq d(p, q) + d(q, r)$

for all $p, q, r \in X$. Then d is called a C^* -algebra metric on X and the triplet (X, \mathbb{A}, d) is called a C^* -algebra valued metric space.

Definition 1.2 ([20]). A sequence $\{x_n\}$ in (X, \mathbb{A}, d) is said to be

- (i) convergent with respect to \mathbb{A} , if for given $\epsilon > 0$, there exists a positive integer k such that $\|d(x_n, x)\| < \epsilon$, for all $n > k$;
- (ii) a Cauchy sequence with respect to \mathbb{A} if for any $\epsilon > 0$, there exists $k \in \mathbb{N}$

such that $\|d(x_n, x_m)\| < \epsilon$, for all $n, m > k$.

The triplet (X, \mathbb{A}, d) is called a complete C^* -algebra valued metric space if every Cauchy sequence with respect to \mathbb{A} is convergent.

Definition 1.3 ([28]). Let f and g be two self-mappings of a metric space (X, d) . Then, the pair (f, g) is said to be weakly compatible if they commute at coincidence points.

Aamri and El Moutawakil [29] introduced the concept of (E.A.) property in metric space.

Definition 1.4 ([29]). Let f and g be two self-mappings of a metric space (X, d) . Then, the pair (f, g) is said to satisfy E.A. property if there exists a sequence $\{x_n\}$ in X such that

$$\lim_{n \rightarrow \infty} f x_n = \lim_{n \rightarrow \infty} g x_n = t \quad \text{for some } t \in X.$$

The (E.A.) property allows to replace the completeness requirement of the space with a more natural condition of closeness of the range.

Sintunavarat and Kumam introduced [30] the concept of (CLR) property in which there is no requirement of closeness of space.

Definition 1.5 ([30]). Let f and g be two self-mappings of a metric space (X, d) . Then, the pair (f, g) is said to satisfy (CLR_f) property if there exists a sequence $\{x_n\} \in X$ such that

$$\lim_{n \rightarrow \infty} f x_n = \lim_{n \rightarrow \infty} g x_n = f t \quad \text{for some } t \in X.$$

2. Main Results

2.1 Common fixed point theorem

Theorem 2.1. Let (X, \mathbb{A}, d) be C^* -algebra valued metric space and A, B, f and g are four self mapping on X satisfying the following conditions:

- (i) $A(X) \subseteq g(X)$ and $B(X) \subseteq f(X)$;
- (ii) for every $x, y \in X, \alpha \in \mathbb{A}$ with $\|\alpha\| \leq 1$,

$$\alpha^* \left((d(fx, Ax)d(fx, By) + d(gy, By)d(gy, Ax)) / (1 + d(fx, By) + d(gy, Ax)) \right) \alpha.$$

If one of $f(X), g(X), A(X)$ and $B(X)$ is a complete subspace of X , the pairs (A, f) and (B, g) have a coincidence point. Moreover, if the pairs (A, f) and (B, g) are weakly compatible then the mapping A, B, f and g have a unique common fixed point in X .

Proof. Let $x_0 \in X$ be an arbitrary point. From (i), we can construct a sequence $\{y_n\}$ in X as follows: $y_{2n+1} = Ax_{2n} = gx_{2n+1}$ and $y_{2n+2} = Bx_{2n+1} = fx_{2n+2}$. Define $d_n = d(y_n, y_{n+1})$. Suppose that $d_{2n} = 0$ i.e. $d(y_{2n}, y_{2n+1}) = 0$ for some n . Then $Ax_{2n} = gx_{2n+1} = Bx_{2n-1} = fx_{2n}$. Thus A and f have coincidence point. Hence the result. Now, suppose that $d_{2n} > 0$ for all $n \in \mathbb{N}$. Then, put $x = x_{2n}$ and $y = x_{2n+1}$ in condition (ii), we have

$$\alpha^* \left((d(fx_{2n}, Ax_{2n})d(fx_{2n}, Bx_{2n+1}) + d(gx_{2n+1}, Bx_{2n+1})d(gx_{2n+1}, Ax_{2n})) / (1 + d(fx_{2n}, Bx_{2n+1}) + d(gx_{2n+1}, Ax_{2n})) \right) \alpha,$$

or

$$d(y_{2n+1}, y_{2n+2}) \leq \alpha^* \left((d(y_{2n}, y_{2n+1})d(y_{2n}, y_{2n+2}) + d(y_{2n+1}, y_{2n+2})d(y_{2n+1}, y_{2n+1})) / (1 + d(y_{2n}, y_{2n+2}) + d(y_{2n+1}, y_{2n+1})) \right) \alpha,$$

or

$$d(y_{2n+1}, y_{2n+2}) \leq \alpha^* \frac{d(y_{2n}, y_{2n+1})d(y_{2n}, y_{2n+2})}{1 + d(y_{2n}, y_{2n+2})} \alpha \leq \alpha^* d(y_{2n}, y_{2n+1}) \alpha.$$

Thus, we have

$$d_{2n+1} \leq \alpha^* d_{2n} \alpha.$$

On the same argument, we can conclude that $d_{2n} \leq \alpha^*(d_{2n-1})\alpha, d_{2n-1} \leq \alpha^*(d_{2n-2})\alpha$ and so on. In general, we have

$$d_n \leq \alpha^*(d_{n-1})\alpha \quad \text{for all } n \in \mathbb{N},$$

i.e

$$d(y_n, y_{n+1}) \leq (\alpha^*)d(y_{n-1}, y_n)\alpha \leq (\alpha^*)^2 d(y_{n-2}, y_{n-1})\alpha^2 \dots \leq (\alpha^*)^n d(y_0, y_1)\alpha^n.$$

For any $p \in \mathbb{N}$ and by using triangle inequality, we get

$$d(y_{n+p}, y_n) \leq d(y_{n+p}, y_{n+p-1}) + d(y_{n+p-1}, y_{n+p-2}) + \dots + d(y_{n+1}, y_n) \leq \sum_{m=n}^{n+p-1} (\alpha^*)^m d(y_0, y_1)\alpha^m \leq \sum_{m=n}^{n+p-1} (\beta\alpha^m)^* \beta\alpha^m$$

$$\begin{aligned} &\leq \sum_{m=n}^{n+p-1} |\beta\alpha^m|^2 (1 + d(u, u) + d(u, Av)) \alpha \\ &\leq \sum_{m=n}^{n+p-1} \|\beta\alpha^m\|^2 \|I_{\mathbb{A}}\| \\ &\leq \|\beta\|^2 I_{\mathbb{A}} \sum_{m=n}^{n+p-1} (\alpha^m)^2 \\ &\rightarrow 0 \text{ as } n \rightarrow \infty, \end{aligned}$$

where $|\beta|^2 = d(y_0, y_1)$ for some $\beta \in \mathbb{A}_+$ and $I_{\mathbb{A}}$ is the unity element in \mathbb{A} . Hence, $\{y_n\}$ is a Cauchy sequence since fX is complete subspace of X . Therefore $\{y_n\}$ is contained in fX and has a limit in fX , say u . Let $v \in f^{-1}u$, then $fv = u$. Next, we shall show that $Av = u$. Assume that, $Av \neq u$. Substituting $x = v$ and $y = x_{n-1}$ in contractive condition (ii), we get

$$\begin{aligned} d(Av, Bx_{n-1}) &\leq \\ &\alpha^* \left((d(fv, Av)d(fv, Bx_{n-1}) \right. \\ &+ d(gx_{n-1}, Bx_{n-1})d(gx_{n-1}, Av)) / \\ &\left. (1 + d(fv, Bx_{n-1}) + d(gx_{n-1}, Av)) \right) \alpha \end{aligned}$$

or

$$\begin{aligned} d(Av, y_n) &\leq \\ &\alpha^* \left((d(fv, Av)d(fv, y_n) \right. \\ &+ d(y_{n-1}, y_n)d(y_{n-1}, Av)) / \\ &\left. (1 + d(fv, y_n) + d(y_{n-1}, Av)) \right) \alpha. \end{aligned}$$

Taking limit as $n \rightarrow \infty$ on both sides, we get

$$\begin{aligned} d(Av, u) &\leq \\ &\alpha^* \left((d(u, Av)d(u, u) \right. \\ &+ d(u, u)d(u, Av)) / \end{aligned}$$

Then, $\|d(Av, u)\| \leq 0$; hence $Av = u$. Thus, we have $fv = u = Av$, where v is the coincidence point of the pair (A, f) . Since $AX \subseteq gX$, $Av = u$, this implies that $u \in gX$. Let $w \in g^{-1}u$, then $gw = u$. Using same argument as above, we can easily verify that $Bw = gw = u$, i.e., w is the coincidence point of the pair (B, g) . The same result can be verified assuming gX complete instead of fX . Now, if $B(X)$ is complete, then by (i) $u \in B(X) \subseteq f(X)$. Similarly, if $A(X)$ is complete $u \in A(X) \subseteq g(X)$. Since the pair (A, f) and (B, g) are weakly compatible, therefore

$$u = Av = fv = gw = Bw,$$

then

$$\begin{aligned} gu &= gBw = Bgw = Bu, \\ fu &= fAv = Afv = Au. \end{aligned}$$

We claim that $Bu = u$. If possible, let $Bu \neq u$.

$$d(u, Bu) = d(Av, Bu) \leq$$

$$\begin{aligned} &\alpha^* \left((d(fv, Av)d(fv, Bu) \right. \\ &+ d(gu, Bu)d(gu, Av)) / \\ &\left. (1 + d(fv, Bu) + d(gu, Av)) \right) \alpha \\ &\leq 0. \end{aligned}$$

Then, $\|d(u, Bu)\| \leq 0$; hence $Bu = u$. On the same lines, we can show that $Au = u$. Thus, we get $Au = fu = gu = Bu = u$. Hence, u is a common fixed point of A, B, f and g .

Next, to prove uniqueness let z be another common fixed point different from u . i.e. $z \neq u$ of A, B, f and g .

$$\begin{aligned}
 d(z, u) &= d(Az, Bu) \leq \alpha^* \left((d(fx, Ax)d(fx, By) \right. \\
 &\quad \left. + d(gy, By)d(gy, Ax)) / \right. \\
 &+ \left. d(gu, Bu)d(gu, Az)) / \right. \\
 &\quad \left. (1 + d(fz, Bu) + d(gu, Az)) \right) \alpha \\
 &\leq \alpha^* \left((d(z, z)d(z, u) \right. \\
 &\quad \left. + (u, u)d(u, z)) / \right. \\
 &\quad \left. (1 + d(z, u) + d(u, z)) \right) \alpha \\
 &\leq 0.
 \end{aligned}$$

Then $\|d(z, u)\| \leq 0$; hence $z = u$. This implies uniqueness. \square

Corollary 2.2. *Let (X, \mathbb{A}, d) be C^* -algebra valued metric space and A and B are two self mapping on X satisfying*

$$\begin{aligned}
 \|d(Ax, By)\| \\
 \leq \|\alpha\| (\|d(x, Ax)\| + \|d(y, By)\|)
 \end{aligned}$$

for any $x, y \in X$, where $\alpha \in \mathbb{A}$ with $\|\alpha\| \leq 1$. Then A and B have a unique common fixed point in X .

Proof. Substituting $f = g = I_X$ in Theorem 2.1, one can easily verify the result. \square

2.2 Common fixed point theorem using (E.A.) property

Theorem 2.3. *Let (X, \mathbb{A}, d) be C^* -algebra valued metric space and A, B, f and g are four self mapping on X satisfying the following conditions:*

(i) $A(X) \subseteq g(X)$ and $B(X) \subseteq f(X)$;

(ii) for every $x, y \in X$, $\alpha \in \mathbb{A}$ with $\|\alpha\| \leq 1$,

$$d(Ax, By) \leq$$

(iii) The pair (A, f) and (B, g) are weakly compatible;

(iv) one of the pair (A, f) or (B, g) satisfy E.A. property.

If the range of one of the mapping $f(X)$ or $g(X)$ is a closed subspace of X , then the mapping A, B, f and g have a unique common fixed point in X .

Proof. Firstly, we assume that the pair (B, g) satisfies E.A. property. Then, by Definition 1.4, there exists a sequence $\{x_n\}$ in X such that $\lim_{n \rightarrow \infty} B(x_n) = \lim_{n \rightarrow \infty} g(x_n) = t$ for some $t \in X$.

Further, $B(X) \subseteq f(X)$, there exists a sequence $\{y_n\}$ in X such that $B(x_n) = f(y_n)$. Hence, $\lim_{n \rightarrow \infty} f(y_n) = t$. We claim that $\lim_{n \rightarrow \infty} A(y_n) = t$. Let if possible $\lim_{n \rightarrow \infty} A(x_n) = t_1 \neq t$. Then, putting $x = y_n$ and $y = x_n$ in condition (ii), we have

$$\begin{aligned}
 d(Ay_n, Bx_n) &\leq \alpha^* \left((d(fy_n, Ay_n)d(fy_n, Bx_n) \right. \\
 &\quad \left. + d(gx_n, Bx_n)d(gx_n, Ay_n)) / \right. \\
 &\quad \left. (1 + d(fy_n, Bx_n) + d(gx_n, Ay_n)) \right) \alpha.
 \end{aligned}$$

Taking norm and $\lim_{n \rightarrow \infty}$ on both sides, we get

$$\begin{aligned}
 \|d(t_1, t)\| &\leq \|\alpha\|^2 \left\| \left((d(t, t_1)d(t, t) \right. \right. \\
 &\quad \left. \left. + d(t, t)d(t, t_1)) \right) \right\| /
 \end{aligned}$$

$$\begin{aligned} & \left. \left. \left. (1 + d(t, t) + d(t, t_1)) \right) \right) \right) \\ & \text{with } \|\alpha\| \leq 1 \\ & \leq 0. \end{aligned}$$

Then $\|d(t_1, t)\| = 0$; hence $t_1 = t$ i.e., $\lim_{n \rightarrow \infty} A(y_n) = \lim_{n \rightarrow \infty} B(x_n) = t$. Now, we suppose that $f(X)$ is closed subspace of X and $fu = t$ for some $u \in X$. Subsequently, we get

$$\begin{aligned} \lim_{n \rightarrow \infty} A(y_n) &= \lim_{n \rightarrow \infty} B(x_n) = \\ \lim_{n \rightarrow \infty} g(x_n) &= \lim_{n \rightarrow \infty} f(y_n) = t = fu. \end{aligned}$$

We claim that $Au = fu$. Put $x = u$ and $y = x_n$ in condition (ii), we have

$$\begin{aligned} d(Au, Bx_n) &\leq \\ & \alpha^* \left((d(fu, Au)d(fu, Bx_n) \right. \\ & + d(gx_n, Bx_n)d(gx_n, Au)) / \\ & \left. (1 + d(fu, Bx_n) + d(gx_n, Au)) \right) \alpha. \end{aligned}$$

Taking norm and $\lim_{n \rightarrow \infty}$ on both sides, we get

$$\begin{aligned} \lim_{n \rightarrow \infty} \|d(Au, t)\| & \\ & \leq \lim_{n \rightarrow \infty} \|\alpha\|^2 \left(\left(d(t, Au)d(t, t) \right. \right. \\ & + d(t, t)d(t, Au)) / \\ & \left. \left. (1 + d(t, t) + d(t, Au)) \right) \right) \\ & \text{with } \|\alpha\| \leq 1, \end{aligned}$$

or

$$\|d(Au, t)\| \leq 0.$$

Then $\|d(Au, t)\| = 0$; hence $Au = t = fu$ i.e u is the coincidence point of the pair (A, f) .

Now the weak compatibility of the pair (A, f) implies that $Afu = fAu$ or $At = ft$.

Since $A(X) \subseteq g(X)$, there exists $v \in X$ such that $Au = gv = fu = t$. Now, we prove that v is coincidence point of pair (B, g) , i.e $Bv = gv = t$. Put $x = u$ and $y = v$ in condition (ii), we get

$$d(Au, Bv) \leq$$

$$\begin{aligned} & \alpha^* \left((d(fu, Au)d(fu, Bv) \right. \\ & + d(gv, Bv)d(gv, Au)) / \\ & \left. (1 + d(fu, Bv) + d(gv, Au)) \right) \alpha. \end{aligned}$$

Taking norm on both sides, we get

$$\begin{aligned} \|d(t, Bv)\| &\leq \\ & \|\alpha\|^2 \left(\left(d(t, t)d(t, Bv) \right. \right. \\ & + d(t, Bv)d(t, t)) / \\ & \left. \left. (1 + d(t, Bv) + d(t, t)) \right) \right) \\ & \text{with } \|\alpha\| \leq 1, \\ & \leq 0. \end{aligned}$$

Then $\|d(Bv, t)\| = 0$; hence $Bv = t$. Thus $Bv = gv = t$ and v is coincidence point of B and g .

Further, the weak compatibility of pair (B, g) implies that $Bgv = gBv$, or $Bt = gt$. Therefore, t is a common coincidence point of A, B, f and g .

Now, we prove that t is a common fixed point of A, B, f and g . Put $x = u$ and $y = t$ in condition (ii), we get

$$d(Au, Bt) \leq$$

$$\begin{aligned} & \alpha^* \left((d(fu, Au)d(fu, Bt) \right. \\ & + d(gt, Bt)d(gt, Au)) / \\ & \left. (1 + d(fu, Bt) + d(gt, Au)) \right) \alpha. \end{aligned}$$

Taking norm on both sides, we get

$$\begin{aligned} & \|d(Bt, t)\| \\ & \leq \|\alpha\|^2 \left(\|d(t, t)d(t, Bt) \right. \\ & \quad \left. + d(t, Bt)d(t, t) / \right. \\ & \quad \left. (1 + d(t, Bt) + d(t, t)) \right\| \\ & \quad \text{with } \|\alpha\| \leq 1, \\ & \leq 0. \end{aligned}$$

Then $\|d(Bt, t)\| = 0$; hence $Bt = t$. Thus, $At = Bt = ft = gt = t$.

Similar arguments arise if we assume that $g(X)$ is closed subspace of X . Similarly, the (E.A.) property of the pair (A, f) will give a similar result.

For uniqueness, let w be another common fixed point of A, B, f and g . Then, put $x = w$ and $y = t$ in condition (ii), we get

$$\begin{aligned} & d(w, t) = d(Aw, Bt) \\ & \leq \alpha^* \left((d(fw, Aw)d(fw, Bt) \right. \\ & \quad \left. + d(gt, Bt)d(gt, Aw)) / \right. \\ & \quad \left. (1 + d(fw, Bt) + d(gt, Aw)) \right) \alpha, \\ & \leq \alpha^* \left((d(w, w)d(w, t) + d(t, t)d(t, w)) / \right. \\ & \quad \left. (1 + d(w, t) + d(t, w)) \right) \alpha \\ & \leq 0. \end{aligned}$$

Then $\|d(w, t)\| \leq 0$; hence $d(w, t) = 0$ i.e $w = t$. Hence t is a unique common fixed point of A, B, f and g . \square

2.3 Common fixed point theorem using (CLR) property

Theorem 2.4. Let (X, \mathbb{A}, d) be C^* -algebra valued metric space and A, B, f and g are four self mapping on X satisfying the following conditions:

- (i) $A(X) \subseteq g(X)$ and $B(X) \subseteq f(X)$;
- (ii) for every $x, y \in X$
 $d(Ax, By) \leq \alpha^* \left((d(fx, Ax)d(fx, By) \right. \\ \left. + d(gy, By)d(gy, Ax)) / (1 + d(fx, By) + d(gy, Ax)) \right) \alpha$;

(iii) The pair (A, f) and (B, g) are weakly compatible;

(iv) one of the pair satisfy (CLR) property.

then the mapping A, B, f and g have a unique common fixed point in X .

Proof. Firstly, we suppose that the pair (B, g) satisfies (CLR_B) property. By Definition 1.5 there exists a sequence $\{x_n\}$ in X such that

$$\lim_{n \rightarrow \infty} Bx_n = \lim_{n \rightarrow \infty} gx_n = Bx = t,$$

for some $x \in X$.

Since $B(X) \subseteq f(X)$, we have $Bx = fu$, for some $u \in X$. We claim that $Au = fu = t$ (say). Put $x = u$ and $y = x_n$ in condition (ii), we get

$$\begin{aligned} & d(Au, Bx_n) \leq \\ & \alpha^* \left((d(fu, Au)d(fu, Bx_n) \right. \\ & \quad \left. + d(gx_n, Bx_n)d(gx_n, Au)) / \right. \\ & \quad \left. (1 + d(fu, Bx_n) + d(gx_n, Au)) \right) \alpha. \end{aligned}$$

Taking norm and $\lim_{n \rightarrow \infty}$ on both sides, we get

$$\begin{aligned} & \|d(Au, Bx)\| \\ & \leq \|\alpha\|^2 \left(\|d(Bx, Au)d(Bx, Bx) \right. \end{aligned}$$

$$\begin{aligned}
 &+ d(Bx, Bx)d(Bx, Au))/ \\
 &\quad \left(1 + d(Bx, Bx) + d(Bx, Au)\right) \Big\| \Big\| \\
 &\quad \text{with } \|\alpha\| \leq 1, \\
 &\leq 0.
 \end{aligned}$$

Then $\|d(Au, Bx)\| = 0$; hence $Au = fu = Bx = t$.

Since $A(X) \subseteq g(X)$, there exists $v \in X$ such that $gv = Au = fu = t$.

Now, we prove that $gv = Bv = t$ i.e., v is the coincidence point of (B, g) . Put $x = u$ and $y = v$ in condition (ii), we get

$$\begin{aligned}
 d(Au, Bv) \leq & \\
 &\alpha^* \left(d(fu, Au)d(fu, Bv) \right. \\
 &+ d(gv, Bv)d(gv, Au))/ \\
 &\quad \left. (1 + d(fu, Bv) + d(gv, Au)) \right) \alpha.
 \end{aligned}$$

Taking norm on both side, we get

$$\begin{aligned}
 \|d(Au, Bv)\| \leq & \\
 &\|\alpha\|^2 \left(\left\| d(fu, Au)d(fu, Bv) \right. \right. \\
 &+ d(gv, Bv)d(gv, Au))/ \\
 &\quad \left. \left. (1 + d(fu, Bv) + d(gv, Au)) \right\| \right) \\
 &\quad \text{with } \|\alpha\| \leq 1, \\
 &\leq 0.
 \end{aligned}$$

Then $\|d(t, Bv)\| = 0$; hence $Bv = t$ i.e $Bv = gv = t$ and v is the coincidence point of B and g .

Further, the weak compatibility of pair (B, g) implies that $Bgv = gBv$ or $Bt = gt$. Therefore, t is a common coincidence point of A, B, f and g . Now, we prove that t is common fixed point of A, B, f and g . Put $x = u$ and $y = t$ in condition (ii), we get

$$\begin{aligned}
 d(Au, Bt) \leq & \\
 &\alpha^* \left(d(fu, Au)d(fu, Bt) \right. \\
 &+ d(gt, Bt)d(gt, Au))/ \\
 &\quad \left. (1 + d(fu, Bt) + d(gt, Au)) \right) \alpha.
 \end{aligned}$$

Taking norm on both side, we get

$$\begin{aligned}
 \|d(t, Bt)\| \leq & \\
 &\|\alpha\|^2 \left(\left\| d(t, t)d(t, Bt) \right. \right. \\
 &+ d(t, Bt)d(t, t))/ \\
 &\quad \left. \left. (1 + d(t, Bt) + d(t, t)) \right\| \right) \\
 &\quad \text{with } \|\alpha\| \leq 1, \\
 &\leq 0.
 \end{aligned}$$

Then $\|d(t, Bt)\| = 0$; hence $Bt = t$. Thus $At = Bt = ft = gt = t$. i.e t is common fixed point of A, B, f and g .

For uniqueness, let w be another common fixed point of A, B, f and g . Then, put $x = w$ and $y = t$ in condition (ii), we get

$$\begin{aligned}
 d(w, t) = d(Aw, Bt) & \\
 \leq \alpha^* \left(d(fw, Aw)d(fw, Bt) \right. & \\
 + d(gt, Bt)d(gt, Aw))/ & \\
 \quad \left. (1 + d(fw, Bt) + d(gt, Aw)) \right) \alpha, & \\
 \leq \alpha^* \left(d(w, w)d(w, t) \right. & \\
 + d(t, t)d(t, w))/ & \\
 \quad \left. (1 + d(w, t) + d(t, w)) \right) \alpha & \\
 \leq 0. &
 \end{aligned}$$

Then $\|d(w, t)\| \leq 0$; hence $d(w, t) = 0$ i.e $w = t$. Hence t is a unique common fixed point of A, B, f and g . \square

Example 2.5. Let $X = [0, 1]$ and $\mathbb{A} = \mathbb{C}$. Define $d : X \times X \rightarrow \mathbb{A}$ by

$$d(x, y) = |x - y|.$$

Then, (X, \mathbb{A}, d) is C^* -algebra-valued metric space.

Define four self maps A, B, f and g on X by

$$\begin{aligned} Ax &= x, & Bx &= \frac{x}{2}, \\ gx &= 2x, & fx &= 4x \quad \forall x \in X. \end{aligned}$$

Clearly,

$$\begin{aligned} AX &= [0, 1] \subset [0, 2] = gX, \\ BX &= \left[0, \frac{1}{2}\right] \subset [0, 4] = fX. \end{aligned}$$

Also, fX is a complete subspace of X and the pair (A, f) and (B, g) are weakly compatible.

$$\begin{aligned} d(Ax, By) &= \left|x - \frac{y}{2}\right|, \\ d(fx, Ax) &= |4x - x|, \\ d(fx, By) &= \left|4x - \frac{y}{2}\right|, \\ d(gy, Ax) &= |2y - x|, \\ \text{and } d(gy, By) &= \left|2y - \frac{y}{2}\right|. \end{aligned}$$

Observe that

$$d(Ax, By) \leq$$

$$\begin{aligned} &\alpha^* \left((d(fx, Ax)d(fx, By) \right. \\ &+ d(gy, By)d(gy, Ax)) / \\ &\left. (1 + d(fx, By) + d(gy, Ax)) \right) \alpha \\ &\forall x, y \in X \text{ with } \|\alpha\| \leq 1. \end{aligned}$$

Here, 0 is the unique common fixed point of A, B, f and g .

Example 2.6. Let $X = [0, 2]$ and $\mathbb{A} = \mathbb{C}$. Define $d : X \times X \rightarrow \mathbb{A}$ by

$$d(x, y) = \begin{cases} |x| + |y| & \text{if } x \neq y \\ 0 & \text{if } x = y. \end{cases}$$

Then, (X, \mathbb{A}, d) is C^* -algebra-valued metric space.

Define four self maps A, B, f and g on X by

$$\begin{aligned} A(x) &= \begin{cases} x & \text{if } x \in [0, 1] \\ 2 & \text{if } x \in (1, 2] \end{cases}, \\ B(x) &= \begin{cases} \frac{x}{2} & \text{if } x \in [0, 1] \\ 1 & \text{if } x \in (1, 2] \end{cases}, \\ g(x) &= \begin{cases} 4x & \text{if } x \in [0, 1] \\ 5 & \text{if } x \in (1, 2] \end{cases} \end{aligned}$$

and

$$f(x) = \begin{cases} 2x & \text{if } x \in [0, 1] \\ 3 & \text{if } x \in (1, 2] \end{cases}.$$

Following cases arises

Case (i): Let $x, y \in [0, 1]$, clearly $AX \subset gX$ and $BX \subset fX$.

Now,

$$\begin{aligned} d(Ax, By) &= x + \frac{y}{2}, & d(fx, Ax) &= 3x, \\ d(fx, By) &= 2x + \frac{y}{2}, & d(gy, By) &= \frac{9y}{2} \\ \text{and } d(gy, Ax) &= x + 4y. \end{aligned}$$

Therefore,

$$\begin{aligned} &\alpha^* \left((d(fx, Ax)d(fx, By) \right. \\ &+ d(gy, By)d(gy, Ax)) / \\ &\left. (1 + d(fx, By) + d(gy, Ax)) \right) \alpha \\ &= \alpha^* \left(\left(3x(2x + \frac{y}{2}) + \frac{9y}{2}(x + 4y) \right) / \right. \end{aligned}$$

$$\begin{aligned}
 & \left(1 + 3x + \frac{9y}{2}\right)\alpha \\
 = & \|\alpha\|^2 \left(\frac{12x^2 + 12xy + 36y^2}{6x + 9y + 2}\right) \\
 \geq & \|\alpha\|^2 \left(\frac{12x^2 + 12xy + 36y^2}{6x + 9y + 3}\right) \\
 = & \|\alpha\|^2 \left(\frac{4x^2 + 4xy + 12y^2}{2x + 3y + 1}\right) \\
 = & \|\alpha\|^2 \left(\frac{((2x + 3y + 1)^2 + 3y^2 - 8xy - 6y - 4x - 1)}{(2x + 3y + 1)}\right) \\
 = & \|\alpha\|^2 \left(\frac{3y^2 - 8xy - 6y - 4x - 1}{2x + 3y + 1}\right) \\
 + & \frac{3y^2 - 8xy - 6y - 4x - 1}{2x + 3y + 1} \\
 \geq & \left(x + \frac{y}{2}\right) = d(Ax, By).
 \end{aligned}$$

Thus,

$$\begin{aligned}
 d(Ax, By) \leq & \alpha^* \left((d(fx, Ax)d(fx, By) \right. \\
 + & \left. d(gy, By)d(gy, Ax)) / \right. \\
 & \left. (1 + d(fx, By) + d(gy, Ax)) \right) \alpha \\
 \forall x, y \in [0, 1] \text{ with } & \|\alpha\| \leq 1.
 \end{aligned}$$

Case (ii): Let $x, y \in (1, 2]$, clearly $AX \subset gX$ and $BX \subset fX$.

Now,

$$\begin{aligned}
 d(Ax, By) &= 3, \quad d(fx, Ax) = 5, \\
 d(fx, By) &= 4, \quad d(gy, By) = 6 \\
 \text{and } d(gy, Ax) &= 7.
 \end{aligned}$$

Therefore,

$$\alpha^* \left((d(fx, Ax)d(fx, By)$$

$$\begin{aligned}
 + & \left. d(gy, By)d(gy, Ax)) / \right. \\
 & \left. (1 + d(fx, By) + d(gy, Ax)) \right) \alpha \\
 = & \alpha^* \left(\frac{62}{12} \right) \alpha \\
 = & \|\alpha\|^2 \left(\frac{62}{12} \right) \\
 \geq & 3 = d(Ax, By).
 \end{aligned}$$

Thus,

$$\begin{aligned}
 d(Ax, By) \leq & \alpha^* \left((d(fx, Ax)d(fx, By) \right. \\
 + & \left. d(gy, By)d(gy, Ax)) / \right. \\
 & \left. (1 + d(fx, By) + d(gy, Ax)) \right) \alpha \\
 \forall x, y \in (1, 2] \text{ with } & \|\alpha\| \leq 1.
 \end{aligned}$$

Also, fX is a complete subspace of X , the pair (A, f) and (B, g) are weakly compatible. Hence, by Theorem 2.1 the mappings A, B, f and g have a unique common fixed point. Indeed, 0 is a common unique fixed point.

Example 2.7. Let $X = [0, 2]$ and $\mathbb{A} = \mathbb{C}$. Define $d : X \times X \rightarrow \mathbb{A}$ by $d(x, y) = |x - y|$. Then, we can easily show that (X, \mathbb{A}, d) is C^* -algebra-valued metric space.

Define four self mappings A, B, f and g on X by

$$A(x) = \begin{cases} 0 & \text{if } x \in [0, 1], \\ 1 & \text{if } x \in (1, 2], \end{cases}$$

$$B(x) = \begin{cases} 0 & \text{if } x \in [0, 1], \\ \frac{1}{2} & \text{if } x \in (1, 2], \end{cases}$$

$$g(x) = \begin{cases} 3x & \text{if } x \in [0, 1], \\ 4 & \text{if } x \in (1, 2], \end{cases}$$

and

$$f(x) = \begin{cases} x & \text{if } x \in [0, 1], \\ 2 & \text{if } x \in (1, 2]. \end{cases}$$

Firstly, we show that the pair (A, f) is satisfying $(E.A.)$ property. Taking $\{x_n\}$ be a sequence in X such that $\{x_n\} = \left(\frac{1}{\sqrt{n^2+2n}}\right)$.

Then,

$$\lim_{n \rightarrow \infty} Ax_n = \lim_{n \rightarrow \infty} A\left(\frac{1}{\sqrt{n^2+2n}}\right) = \lim_{n \rightarrow \infty} (0) = 0$$

and

$$\begin{aligned} \lim_{n \rightarrow \infty} fx_n &= \lim_{n \rightarrow \infty} f\left(\frac{1}{\sqrt{n^2+2n}}\right) \\ &= \lim_{n \rightarrow \infty} \left(\frac{1}{\sqrt{n^2+2n}}\right) = 0. \end{aligned}$$

So, there exists a sequence $\{x_n\}$ in X such that $\lim_{n \rightarrow \infty} Ax_n = \lim_{n \rightarrow \infty} fx_n = 0 \in X$. Hence, the pair (A, f) satisfy $(E.A.)$ property. Similarly, we can show that the pair (B, g) satisfy $(E.A.)$ property.

Following cases arise

Case(i) : Let $x, y \in [0, 1]$, clearly $AX \subset gX$ and $BX \subset fX$.

Now,

$$\begin{aligned} d(Ax, By) &= 0, \quad d(fx, Ax) = x, \\ d(fx, By) &= x, \quad d(gy, By) = 3y \\ \text{and } d(gy, Ax) &= 3y. \end{aligned}$$

Therefore,

$$\begin{aligned} &\alpha^* \left((d(fx, Ax)d(fx, By) \right. \\ &+ \left. d(gy, By)d(gy, Ax)) / \right. \\ &\quad \left. (1 + d(fx, By) + d(gy, Ax)) \right) \alpha \\ &= \alpha^* \left(\frac{(x * x) + (3y * 3y)}{x + 3y + 1} \right) \alpha \\ &= \|\alpha\|^2 \left(\frac{x^2 + 9y^2}{x + 3y + 1} \right) \end{aligned}$$

$$\geq 0 = d(Ax, By).$$

Thus,

$$\begin{aligned} d(Ax, By) &\leq \\ &\alpha^* \left((d(fx, Ax)d(fx, By) \right. \\ &+ \left. d(gy, By)d(gy, Ax)) / \right. \\ &\quad \left. (1 + d(fx, By) + d(gy, Ax)) \right) \alpha \\ &\forall x, y \in [0, 1] \text{ with } \|\alpha\| \leq 1. \end{aligned}$$

Case (ii): Let $x, y \in (1, 2]$, clearly $AX \subset gX$ and $BX \subset fX$.

Now,

$$\begin{aligned} d(Ax, By) &= \frac{1}{2}, \quad d(fx, Ax) = 1, \\ d(fx, By) &= \frac{3}{2}, \quad d(gy, By) = \frac{7}{2} \\ \text{and } d(gy, Ax) &= 3. \end{aligned}$$

Therefore,

$$\begin{aligned} &\alpha^* \left((d(fx, Ax)d(fx, By) \right. \\ &+ \left. d(gy, By)d(gy, Ax)) / \right. \\ &\quad \left. (1 + d(fx, By) + d(gy, Ax)) \right) \alpha \\ &= \alpha^* \left(\frac{24}{11} \right) \alpha \\ &= \|\alpha\|^2 \left(\frac{24}{11} \right) \\ &\geq \frac{1}{2} = d(Ax, By). \end{aligned}$$

Thus,

$$\begin{aligned} d(Ax, By) &\leq \\ &\alpha^* \left((d(fx, Ax)d(fx, By) \right. \\ &+ \left. d(gy, By)d(gy, Ax)) / \right. \\ &\quad \left. (1 + d(fx, By) + d(gy, Ax)) \right) \alpha \end{aligned}$$

$$\forall x, y \in (1, 2] \text{ with } \|\alpha\| \leq 1.$$

Also, fX is a closed subspace of X , the pair (A, f) and (B, g) are weakly compatible. Hence, by Theorem 2.3 the mappings A, B, f and g have a unique common fixed point. Indeed, 0 is a common unique fixed point.

Example 2.8. Let $X = [0, 2]$ and $\mathbb{A} = \mathbb{C}$. Define $d : X \times X \rightarrow \mathbb{A}$ by

$$d(x, y) = \begin{cases} |x| + |y| & \text{if } x \neq y \\ 0 & \text{if } x = y \end{cases}$$

Then, (X, \mathbb{A}, d) is C^* -algebra-valued metric space.

Define four self maps A, B, f and g on X by

$$A(x) = \begin{cases} 0 & \text{if } x \in [0, 1] \\ x & \text{if } x \in (1, 2] \end{cases},$$

$$B(x) = \begin{cases} x & \text{if } x \in [0, 1] \\ \frac{3}{2} & \text{if } x \in (1, 2] \end{cases},$$

$$g(x) = \begin{cases} 3x & \text{if } x \in [0, 1] \\ 2x & \text{if } x \in (1, 2] \end{cases}$$

and

$$f(x) = \begin{cases} 4x & \text{if } x \in [0, 1] \\ 7 & \text{if } x \in (1, 2]. \end{cases}$$

Firstly, we show that the pair (A, f) is satisfying (CLR_A) property. Taking $\{x_n\}$ be a sequence in X such that $\{x_n\} = \left(\frac{1}{n^2+2n+3}\right)$. Then,

$$\begin{aligned} \lim_{n \rightarrow \infty} Ax_n &= \lim_{n \rightarrow \infty} A\left(\frac{1}{n^2+2n+3}\right) \\ &= \lim_{n \rightarrow \infty} (0) = 0 \end{aligned}$$

and

$$\begin{aligned} \lim_{n \rightarrow \infty} fx_n &= \lim_{n \rightarrow \infty} f\left(\frac{1}{n^2+2n+3}\right) \\ &= \lim_{n \rightarrow \infty} \left(\frac{4}{n^2+2n+3}\right) = 0. \end{aligned}$$

So, there exists a sequence $\{x_n\}$ in X such that $\lim_{n \rightarrow \infty} Ax_n = \lim_{n \rightarrow \infty} fx_n = A(0) = 0$ for $0 \in X$. Hence, the pair (A, f) satisfy (CLR_A) property. Similarly, we can show that the pair (B, g) satisfy (CLR_B) property.

Following cases arise

Case(i) : Let $x, y \in [0, 1]$, clearly $AX \subset gX$ and $BX \subset fX$.

Now,

$$\begin{aligned} d(Ax, By) &= y, \quad d(fx, Ax) = 4x \\ d(fx, By) &= 4x + y, \quad d(gy, By) = 4y \\ \text{and } d(gy, Ax) &= 3y. \end{aligned}$$

Therefore,

$$\begin{aligned} &\alpha^* \left((d(fx, Ax)d(fx, By) \right. \\ &+ d(gy, By)d(gy, Ax)) / \\ &\left. (1 + d(fx, By) + d(gy, Ax)) \right) \alpha \\ &= \alpha^* \left(\frac{4x(4x + y) + 4y(3y)}{1 + 4x + y + 3y} \right) \alpha \end{aligned}$$

$$= \|\alpha\|^2 \left(\frac{16x^2 + 4xy + 12y^2}{4x + 4y + 1} \right)$$

$$\geq \|\alpha\|^2 \left(\frac{16x^2 + 4xy + 12y^2}{4x + 4y + 4} \right)$$

$$= \|\alpha\|^2 \left(\frac{4x^2 + xy + 3y^2}{x + y + 1} \right)$$

$$= \|\alpha\|^2 \left((x + y + 1)^2 \right)$$

$$+ (2y^2 + 3x^2 - y - x - 1) /$$

$$\begin{aligned} & (x + y + 1) \\ = & \|\alpha\|^2 \left((x + y + 1) \right. \\ & \left. + \frac{3x^2 + 2y^2 - y - x - 1}{x + y + 1} \right) \\ \geq & y = d(Ax, By). \end{aligned}$$

Thus,

$$\begin{aligned} d(Ax, By) \leq & \alpha^* \left((d(fx, Ax)d(fx, By) \right. \\ & \left. + d(gy, By)d(gy, Ax)) / \right. \\ & \left. (1 + d(fx, By) + d(gy, Ax)) \right) \alpha \\ & \forall x, y \in [0, 1] \text{ with } \|\alpha\| \leq 1. \end{aligned}$$

Case(ii) : Let $x, y \in (1, 2]$, clearly $AX \subset gX$ and $BX \subset fX$.

Now,

$$\begin{aligned} d(Ax, By) &= x + \frac{3}{2}, \quad d(fx, Ax) = 7 + x \\ d(fx, By) &= \frac{17}{2}, \quad d(gy, By) = 2y + \frac{3}{2} \\ \text{and } d(gy, Ax) &= 2y + x. \end{aligned}$$

Therefore,

$$\begin{aligned} & \alpha^* \left((d(fx, Ax)d(fx, By) \right. \\ & \left. + d(gy, By)d(gy, Ax)) / \right. \\ & \left. (1 + d(fx, By) + d(gy, Ax)) \right) \alpha \\ = & \alpha^* \left(\frac{(\frac{17}{2}(7+x)) + (2y + \frac{3}{2})(2y+x)}{1 + 2y + x + \frac{17}{2}} \right) \alpha \\ = & \|\alpha\|^2 \left(\frac{17(7+x) + (4y+3)(2y+x)}{4y+2x+19} \right) \\ = & \|\alpha\|^2 \left(\frac{20x+6y+8y^2+4xy+119}{4y+2x+19} \right) \\ = & \|\alpha\|^2 \left(4 + \frac{12x-18y+8y^2+4xy+43}{4y+2x+19} \right) \end{aligned}$$

$$\geq x + \frac{3}{2} = d(Ax, By).$$

Thus,

$$\begin{aligned} d(Ax, By) \leq & \alpha^* \left((d(fx, Ax)d(fx, By) \right. \\ & \left. + d(gy, By)d(gy, Ax)) / \right. \\ & \left. (1 + d(fx, By) + d(gy, Ax)) \right) \alpha \\ & \forall x, y \in (1, 2] \text{ with } \|\alpha\| \leq 1. \end{aligned}$$

The pair (A, f) and (B, g) are weakly compatible. Hence, by the Theorem 2.4 the mappings A, B, f and g have a unique common fixed point. Indeed, 0 is a common unique fixed point.

Acknowledgements

The authors are thankful to the learned referees for the very careful reading and valuable suggestions.

References

- [1] S. Banach, Sur les opérations dans les ensembles abstraits et leur application aux équations intégrales, *Fundamenta Mathematicae*, 1922, 3:133-81.
- [2] Y.J. Cho and N. Petrot, Existence theorems for fixed fuzzy points with closed α -cut sets in complete metric spaces, *Communications of the Korean Mathematical Society*, 2011, 26(1):115-24.
- [3] X. Du and Z. Zhao, On Fixed Point Theorems of Mixed Monotone Operators, *Fixed Point Theory and Applications*, (2011).
- [4] W. Liguang, L. Bo and B. Ran, Stability of a Mixed Type Functional Equation on Multi-Banach Spaces: A Fixed Point Approach, *Fixed Point Theory and Applications*, (2010).

- [5] X. Li and Z. Zhao, On a fixed point theorem of mixed monotone operators and applications, *Electronic Journal of Qualitative Theory of Differential Equations*, 2012, 94:1-7.
- [6] A.F. Roldán-López-de-Hierro and W. Sintunavarat, Common fixed point theorems in fuzzy metric spaces using the CLRg property, *Fuzzy Sets and Systems*, 2016, 282:131-42.
- [7] Z. Shi-sheng, Fixed point theorems for fuzzy mappings (II), *Applied Mathematics and Mechanics*, 1986, 7(2):147-52.
- [8] W. Sintunavarat, Fixed point results in b-metric spaces approach to the existence of a solution for nonlinear integral equations, *Revista de la Real Academia de Ciencias Exactas, Físicas y Naturales. Serie A. Matemáticas*, 2016, 110(2):585-600.
- [9] W. Sintunavarat and P. Kumam, Best proximity points theorems for generalized Mizoguchi-Takahashi's contraction pairs, *Journal of Nonlinear and Convex Analysis*, 2016, 17(7):1345-61.
- [10] W. Sintunavarat, M. B. Zada and M. Sarwar, Common solution of Urysohn integral equations with the help of common fixed point results in complex valued metric spaces, *Revista de la Real Academia de Ciencias Exactas, Físicas y Naturales. Serie A. Matemáticas*, 2017, 111(2):531-45.
- [11] X. Zhang, L. Liu and Y. Wu, Fixed point theorems for the sum of three classes of mixed monotone operators and applications, *Fixed Point Theory and Applications*, (2016).
- [12] Z. Zhao, Existence and uniqueness of fixed points for some mixed monotone operators, *Nonlinear Analysis*, 2010, 73(6):1481-90.
- [13] Z. Zhao, Fixed points of τ - ϕ -convex operators and applications, *Applied Mathematics Letters*, 2010, 23(5):561-6.
- [14] X. Hao, L. Liu and Y. Wu, Positive solutions for second order differential systems with non local conditions, *Fixed Point Theory*, 2012, 13(2):507-16.
- [15] L. Liu, X. Zhang, J. Jiang and Y. Wu, The unique solution of a class of sum mixed monotone operator equations and its application to fractional boundary value problems, *Journal of Nonlinear Sciences and Applications*, 2016, 9(5):2943-58.
- [16] J. Mao, Z. Zhao and N. Xu, The existence and uniqueness of positive solutions for integral boundary value problems, *Electronic Journal of Qualitative Theory of Differential Equations*, 2010, 16:1-8.
- [17] B. Singh, V. Gupta and S. Kumar, Common Fixed Point Theorems Using the E.A. and CLR Properties in 2-Menger Spaces, *International Journal of Analysis* (2013).
- [18] T.Wang and R. Xu, Bounds for Some New Integral Inequalities With Delay on Time Scales, *Journal of Mathematical Inequalities*, 2012, 6(1):1-12.
- [19] G. Wang, H.T. Che and H.B. Chen, Feasibility-solvability theorems for generalized vector equilibrium problem in reflexive Banach spaces, *Fixed Point Theory and Applications*, 2012, 1-13.
- [20] Z. Ma, L. Jiang and H. Sun, C^* -algebra valued metric spaces and related fixed point theorems, *Fixed Point Theory and Applications*, 2014, pp. 11.
- [21] M. Asim and M. Imdad, C^* -algebra valued extended b-metric spaces and fixed point results with an application, *UPB Scientific Bulletin, Series A: Applied Mathematics and Physics*, 2020, 82(1):207-18.
- [22] M. Asim and M. Imdad, C^* -algebra valued symmetric spaces and fixed point results with an application, *Korean Journal of Mathematics*, 2010, 28(1):17-30.

- [23] K.R. Davidson, C^* -algebras by example, Fields institute monographs 6, American Mathematical Society, Providence, RI, (1996).
- [24] G.J. Murphy, C^* -algebras and operator theory, Academic Press, Boston, (1990).
- [25] S. Radenovic, P. Vetro, A. Nastasi and LT. Quan, Coupled Fixed Point Theorems in C^* -Algebra-Valued b -Metric Spaces, Scientific Publications of the State University of Novi Pazar, 2017, 9(1):81-90.
- [26] C. Vetro, On Branciaris theorem for weakly compatible mappings, Applied Mathematics Letters, 2010, 23:700-5.
- [27] Q. Xin, L. Jiang and Z. Ma, Common fixed point theorems in C^* -algebra valued metric spaces, Journal of Nonlinear Sciences and Applications, 2016, 9:4617-27.
- [28] G. Jungck, Common fixed points for non-continuous non self mappings on non-metric spaces, Far East Journal of Mathematical Sciences (FJMS), 1996, 4:199-212.
- [29] M. Aamri and D.El. Moutawakil, Some new common fixed point theorems under strict contractive conditions, Journal of Mathematical Analysis and Applications, 2002, 270:181-8.
- [30] W. Sintunavarat and P. Kumam, Common fixed point theorems for a pair of weakly compatible mappings in fuzzy metric spaces, Journal of Applied Mathematics, 2011, pp. 14.
- [31] M. Kumar, M. Imdad and M. Asim, Some Fixed Point Theorems under E.A. and (CLR) Properties on C^* -Algebra Valued Metric Spaces, Information Science Letters, (2020).