

## LIST OF FIGURES

| Figure   | Page |
|--|------|
| 1 The framework of ZSM-5 zeolite composed of well-defined pores and channels where water or guest molecules can occupy.  | 3    |
| 2 Flat hexagon lattices of a graphene sheet showing the chiral vector R and the wrapping angle $\phi$ .  | 6    |
| 3 Through slicing of graphene sheet (a) by $60^\circ$ disclination angle and roll up, one could construct the conical tip (b) of a single wall carbon nanohorn where the top region part is similar to that of the encapsulated single wall carbon nanotube (c).   | 8    |
| 4 The embedded cluster model established from the SCREEP approach composing of 3 layers: The quantum cluster model, explicit charges and surface charges   | 28   |
| 5 Cluster models for a) carbon nanohorn ( $C_{80}H_{20}$ ) and b) carbon nanotube ( $C_{80}H_{10}$ ) tips. Nine positions (site I-IX) among an analogous region (shaded area) of both models were examined as to whether they could be the possible sitting sites for the gold atom. Both of the tips share the same atom and site labeling. | 35   |
| 6 Cluster and embedded cluster models of the H-ZSM-5 zeolite. Bond distances given in the figures are in Å and bond angle is in degrees.   | 37   |
| 7 Cluster and embedded cluster models of the Li-ZSM-5 zeolite. Bond distances given in the figures are in Å and bond angle is in degrees.  | 39   |
| 8 Cluster and embedded cluster models of the H-ZSM-5/CO complex. Bond distances are in Å and bond angle is in degrees.   | 43   |
| 9 Cluster and embedded cluster models of the Li-ZSM-5/CO complex. Bond distances are in Å and bond angle is in degrees.  | 44   |
| 10 Cluster and embedded cluster models of the H-FAU/CO and Li-FAU/CO complex. Bond distances given in the figures are in Å.  | 47   |

## LIST OF FIGURES (cont'd)

| Figure  | Page |
|---|------|
| 11 Ag-ZSM-5 structures optimized at B3LYP/6-31G(d,p) using finite and embedded calculation (values in parenthesis) of a) 3T cluster and b) 5T cluster; bond distance in Å.  | 50   |
| 12 Optimized structures of the adsorption of CO on Ag-ZSM-5 zeolite at the B3LYP/6-31G(d,p) level using bare clusters. Selected bond distances (Å) are also given. Values in parentheses are from the embedded cluster model. a) 3T cluster and b) 5T cluster   | 52   |
| 13 Natural Bond Orbital (NBO) plots showing the interaction between an exchanged $\text{Ag}^+$ and an absorbed CO: a) The sigma interaction between $2\text{s}^{62.73\%}2\text{p}^{37.27\%}(\text{C})$ with $5\text{s}(\text{Ag})$ and the $\pi$ -back bonding electron transfers between $4\text{d}(\text{Ag})$ and $\pi^*(\text{CO})$ b) in xz plane and c) in yz plane | 54   |
| 14 Optimized structures of the adsorption of NO on Ag-ZSM-5 zeolite at the B3LYP/6-31G(d,p) level using bare clusters. Selected bond distances (Å) are also given. Values in parentheses are from the embedded cluster model. a) 3T cluster and b) 5T cluster   | 56   |
| 15 Natural Bond Orbital (NBO) plots showing the interaction between an exchanged $\text{Ag}^+$ and an absorbed NO: a) The sigma interaction between $2\text{s}^{10.76\%}2\text{p}^{89.21\%}(\text{N})$ and $5\text{s}(\text{Ag})$ b) The $\pi$ -back bonding electron transfer between $4\text{d}(\text{Ag})$ and $\pi^*(\text{NO})$                                      | 58   |
| 16 Interaction energy diagram for a gold atom adsorbed on the Nanohorn (SWNH) and Nanotube (SWNT) tips. Energies are in kcal/mol.   | 66   |
| 17 Molecular orbital plots for a) HOMO of SWNH tip, b) LUMO of SWNH tip, c) HOMO of SWNT tip, and d) LUMO of SWNT tip.  | 68   |
| 18 Interaction of the CO molecule with the gold atom on the top of a) Nanohorn tip and b) Nanotube tip, distances are shown in picometers and angles are shown in degrees.  | 70   |

## LIST OF FIGURES (cont'd)

| Figure   | Page |
|--|------|
| 19      Electron density difference plots for a) Au---SWNH (3D) system b) Au---SWNH (2D) system on the plane that contains C <sub>1</sub> ,C <sub>2</sub> , and Au atoms c) Au---SWNH (2D) system on the plane that cuts between C <sub>1</sub> and C <sub>2</sub> d) Au---SWNT (3D) system e) Au---SWNT (2D) system on the plane that contains C <sub>1</sub> ,C <sub>6</sub> , and Au atoms. 3D plots use the isovalue = 0.001 e/Å <sup>3</sup> . 2D plots start with isovalue = ±0.001 e/Å <sup>3</sup> and the interval = 0.01 e/Å <sup>3</sup> . The blue area represents the electron accumulation while the pink region corresponds to the electron diminishing.                    | 71   |
| 20      Electron density difference plots for a) CO---Au/SWNH (3D) system b) CO---Au/SWNH (2D) system on the plane that contains C <sub>1</sub> ,C <sub>2</sub> , and Au atoms c) CO---Au/SWNH (2D) system on the plane that cuts between C <sub>1</sub> and C <sub>2</sub> d) CO---Au/SWNT (3D) system e) CO---Au/SWNT (2D) system on the plane that contains C <sub>1</sub> ,C <sub>6</sub> , and Au atoms. 3D plots use the isovalue = 0.001 e/Å <sup>3</sup> . 2D plots start with isovalue = ±0.001 e/Å <sup>3</sup> and use the interval = 0.01 e/Å <sup>3</sup> . The blue area represents the electron accumulation while the pink region corresponds to the electron diminishing. | 73   |