

## Mosses from the Molawin-Dampalit Subwatershed, Mt. Makiling Forest Reserve, the Philippines

Ailene A. Alcala<sup>1\*</sup>, Ramil S. Alcala<sup>2</sup> and Eugene L. R. Logatoc<sup>3</sup>

<sup>1</sup>Forestry Development Center, College of Forestry and Natural Resources, University of the Philippines Los Baños, 4031 College, Laguna, Philippines

<sup>2</sup>Citicore Renewable Energy Corp., 11th F 276 Col. Boni Serrano Ave., Santolan Town Center, San Juan City, the Philippines 36577 Bangkal St. (Extension), 4030 Los Baños, Laguna, the Philippines

\*corresponding author: [aaalcala1@up.edu.ph](mailto:aaalcala1@up.edu.ph)

### ABSTRACT

Despite being important components of natural ecosystems, mosses are often not included in forest inventories, assessment, and monitoring particularly in the Philippine setting. In view of this, we aimed to document and revisit the moss flora of Mt. Makiling Forest Reserve which is regarded as one of the most biologically explored areas in the country. A collection of mosses from the Molawin-Dampalit Subwatershed of the Mt. Makiling Forest Reserve yielded a total of 27 species, one (1) subspecies, and one (1) variety in 22 genera and 10 families. These represent 12.16% of the species, 22% of the genera, and 31.25% of the families reported for the mountain. The most represented families are Neckeraceae (5 species), Calymperaceae, Pylaisiadelphaceae (3 species each). *Syrrhopodon aristifolius* Mitt. is hereby reported as new record for Luzon Island. With this, the importance and need for continued studies on the mosses is highlighted especially in underexplored portions of Mt. Makiling and should be ultimately geared towards a revised checklist of the mosses for the mountain.

**Keywords:** mosses, bryophytes, Mt. Makiling, Philippines.

### INTRODUCTION

Mount Makiling Forest Reserve and ASEAN Heritage Park (MMFR) is the subject of our recent studies which aimed to update the moss flora of the mountain and its vicinity (Alcala *et al.*, 2020). As one of the most biologically known areas in the Philippines, the mountain had been subject to numerous bryological studies in the past (*e.g.* Tan, 1982; Tixier, 1972) –resulting to 213 species, three (3) subspecies, nine (9) varieties), and one (1) forma in 96 genera and 31 families of mosses known for the area. Our recent study on mosses collected in the Greater Sipit Subwatershed yielded 36 species and one (1) subspecies, of which 10 species, one (1) genus, and one (1) family are newly recorded for the mountain (Alcala *et al.*, 2020). In addition to this, only a handful of studies have been conducted in the MMFR recently (*e.g.* Magcale-Macandog *et al.*, 2022); hence, the need to revisit and explore other areas in the mountain became apparent.

Despite being important components of natural ecosystems (Hallingbäck and Tan, 2010), bryophytes in general are not included in the recommended guidelines of the Department of Environment and

Natural Resources of the Philippines (see DENR–BMB and GIZ, 2017). The inclusion of bryophytes in forest inventories, assessment, and monitoring in the Philippines is particularly important in order to provide baseline information and monitor bryophyte richness in a particular area. An example of this is presented by Sastre-De Jesus and Tan (1995) who discussed how human activities reduced the number of mosses of Mt. Santo Tomas in Benguet from 175 species to only 80 species. With this, revisiting biologically explored areas may provide up-to-date information on the bryophyte species richness of an area; in this case for Mt. Makiling.

In this report, we present the second part of our research which focused on the collections made in the Molawin-Dampalit Subwatershed along the northeastern slope of the mountain. This covers the area along the Mariang Makiling Trail which traverses from the University of the Philippines Los Baños campus up to the summit at Peak 2 (1,090 m asl). This trail is characterized by an access road up to around 500 m asl at Agila Base and is frequently used by locals and tourists alike. Also presented here are information on the substrate, habitat, and intra-Philippine distribution of each species.

## MATERIALS AND METHODS

The methodologies followed in this research are presented in detail by Alcala *et al.* (2020). The elevation and geographical location of the plots were determined using a Global Navigation Satellite System (GNSS) receiver (Table 1; Fig. 1). Voucher specimens were examined and identified in the Department of Forest Biological Sciences, College of Forestry and Natural

Resources, University of the Philippines Los Baños and were deposited at CAHUP. Information on the intra-Philippine distribution of species were consulted primarily from Tan and Iwatsuki (1991) and the works of Azuelo *et al.* (2015), Linis (2009, 2010, 2014, 2018, 2019, 2021), Linis and Tan (2013), Lubos (2007), Shevock and Yorong (2018), Tan *et al.* (2000), Tan (1996), Tan and Shevock (2014), Tan *et al.* (2015), and Tan *et al.* (2017).

**Table 1.** Collection sites within the Molawin-Dampalit Subwatershed, Mt. Makiling.

No.	Site Description	Collection Date
1	Tropical lowland evergreen rainforest at 300 m elevation; WGS 84. Lat/Long: 14°08.52' N, 121°13.82' E	13 February 2016
2	Tropical lowland evergreen rainforest at 400 m elevation; WGS 84. Lat/Long: 14°08.16' N, 121°13.75' E	13 February 2016
3	Tropical lowland evergreen rainforest at 500 m elevation; WGS 84. Lat/Long: 14°07.77' N, 121°12.66' E	13 February 2016
4	Tropical lowland evergreen rainforest at 600 m elevation; WGS 84. Lat/Long: 14°08.03' N, 121°12.48' E	13 February 2016
5	Tropical lowland evergreen rainforest at 700 m elevation; WGS 84. Lat/Long: 14°08.31' N, 121°12.32' E	13 February 2016
6	Tropical lower montane rainforest at 800 m elevation; WGS 84. Lat/Long: 14°08.35' N, 121°11.94' E	13 February 2016
7	Tropical upper montane rainforest at 900 m elevation; WGS 84. Lat/Long: 14°08.28' N, 121°11.81' E	13 February 2016
8	Tropical upper montane rainforest at 1,000 m elevation; WGS 84. Lat/Long: 14°08.17' N, 121°11.74' E	13 February 2016

## RESULTS AND DISCUSSION

### Enumeration of Taxa

A total of 27 species, one (1) subspecies, and one (1) variety in 22 genera and 10 families were recorded in the Molawin-Dampalit Subwatershed of Mt. Makiling Forest Reserve and ASEAN Heritage Park. The list is presented in alphabetical order for ease of reference –with each species annotated by locality (see Table 1), voucher specimen number, substrate and habitat information, and intra-Philippine distribution. An asterisk (\*) indicates a new record for Mt. Makiling and vicinity. Nomenclature follows Tan and Iwatsuki (1991) and Brinda and Atwood (2022, continuously updated), adjusted following recent nomenclatural changes.

*Acroporium hermaphroditum* (Müll. Hal.) M. Fleisch. [Sematophyllaceae] – SITE 8: A.A. Alcala 91. Epiphytic in tropical upper montane rainforest. Intra-Philippine Distribution: Batan, Luzon, Mindoro, Camiguin, and Mindanao (Linis, 2009; Tan and Iwatsuki, 1991).

*Acroporium secundum* (Reinw. & Hornsch.) M. Fleisch. [Sematophyllaceae] – SITE 7: A.A. Alcala 83. SITE 8: A.A. Alcala 93. Epiphytic or epixylic in tropical upper montane rainforest. Intra-Philippine Distribution: Luzon, Mindoro, Palawan, Negros, Camiguin, and Mindanao (Linis, 2009, 2010, 2014; Tan and Iwatsuki, 1991).

*Barbella compressiramea* (Renauld & Cardot) M. Fleisch. [Meteoriaceae] – SITE 3: A.A. Alcala 56.



**Figure 1.** Collection sites in the Molawin-Dampalit Subwatershed: the tropical lowland evergreen rainforest from 300 m to 700 m asl (A-E); the tropical lower montane rainforest at 800 m asl (F); and the tropical upper montane rainforest at 900 m to 1000 m asl (G-I).

Epiphytic in tropical lowland evergreen rainforest. Intra-Philippine Distribution: Luzon (Linis, 2019; Tan and Iwatsuki, 1991).

*Caduciella mariei* (Besch.) Enroth [Neckeraceae] – SITE 1: A.A. Alcala 50. Epiphytic in tropical lowland evergreen rainforest. Intra-Philippine Distribution: Luzon, Catanduanes, Palawan, and Mindanao (Linis, 2014; Lubos, 2010 as *Pinnatella mariei*; Tan and Iwatsuki, 1991 as *Pinnatella mariei*).

*Circulifolium exiguum* (Bosch & Sande Lac.) S. Olsson, Enroth & D. Quandt [Neckeraceae] – SITE 3: A.A. Alcala 57. Epiphytic in tropical lowland evergreen rainforest. Intra-Philippine Distribution: Batan, Luzon, Catanduanes, Mindoro, and Mindanao (Linis, 2014; Tan and Iwatsuki, 1991 as *Homaliodendron exiguum*).

*Claopodium prionophyllum* (Müll. Hal.) Broth. [Leskeaceae] – SITE 5: A.A. Alcala 75. On rocks in tropical lowland evergreen rainforest. Intra-Philippine Distribution: Widespread (Tan and Iwatsuki, 1991).

*Dendro-hypnum reinwardtii* subsp. *caducifolium* (Herzog) N.E. Bell, A.E. Newton & D. Quandt

[Hypnodendraceae] – SITE 8: A.A. Alcala 86. Epiphytic in tropical upper montane rainforest. Intra-Philippine Distribution: Widespread (Tan and Iwatsuki, 1991 as *Hypnodendron reinwardtii* subsp. *caducifolium*).

*Exostratum blumei* (Nees ex Hampe) L.T. Ellis [Calymperaceae] – SITE 5: A.A. Alcala 72 & 74. Epiphytic in tropical lowland evergreen rainforest. Intra-Philippine Distribution: Batan, Luzon, Catanduanes, Negros, Cebu, Camiguin, and Mindanao (Linis, 2010; Tan and Iwatsuki, 1991).

*Fissidens braunii* (Müll. Hal.) Dozy & Molke. [Fissidentaceae] – SITE 3: A.A. Alcala 63. Epiphytic in tropical lowland evergreen rainforest. Intra-Philippine Distribution: Batan, Luzon, Lubang, and Mindanao (Tan and Iwatsuki, 1991).

*Fissidens crispulus* Brid. [Fissidentaceae] – SITE 6: A.A. Alcala 77. Epiphytic in tropical lower montane rainforest. Intra-Philippine Distribution: Widespread (Tan and Iwatsuki, 1991 as *F. zippelianus*).

*Fissidens crispulus* var. *robinsonii* (Broth.) Z. Iwats. & Z.H. Li [Fissidentaceae] – SITE 1: A.A. Alcala 49.

- SITE 3: A.A. Alcalá 64. SITE 4: A.A. Alcalá 66. On rocks or soil in tropical lowland evergreen rainforest. Intra-Philippine Distribution: Luzon, Palawan, Panay, and Mindanao (Linis, 2019 as *F. robinsonii*; Tan, 1996 *F. zippelianus* var. *robinsonii*; Tan and Iwatsuki, 1991 as *F. zippelianus* var. *robinsonii*).
- Himantocladium plumula* (Nees) M. Fleisch. [Neckeraceae] – SITE 3: A.A. Alcalá 58. Epiphytic in tropical lowland evergreen rainforest. Intra-Philippine Distribution: Luzon, Mindoro, Samar, and Mindanao (Tan and Iwatsuki, 1991).
- Homali dendron flabellatum* (Sm.) M. Fleisch. [Neckeraceae] – SITE 8: A.A. Alcalá 87. Epiphytic in tropical upper montane rainforest. Intra-Philippine Distribution: Widespread (Tan and Iwatsuki, 1991).
- Isopterygium albescens* (Hook.) A. Jaeger [Pylaisiadelphaceae] – SITE 2: A.A. Alcalá 53. SITE 4: A.A. Alcalá 67, 68, & 69. Epiphytic or on soil in tropical lowland evergreen rainforest. Intra-Philippine Distribution: Widespread (Linis, 2009 as *I. minutirameum*, 2010, 2014; Tan, 1996 as *I. minutirameum*; Tan and Iwatsuki, 1991).
- Leucobryum bowringii* Mitt. [Leucobryaceae] – SITE 7: A.A. Alcalá 79 & 80a. SITE 8: A.A. Alcalá 88, 89, & 90. Epiphytic or epixylic in tropical upper montane rainforest. Intra-Philippine Distribution: Batan, Luzon, Catanduanes, Mindoro, Sibuyan, Negros, and Panay (Linis, 2009, 2014; Tan and Iwatsuki, 1991).
- Leucobryum sanctum* (Nees ex Schwägr.) Hampe [Leucobryaceae] – SITE 7: A.A. Alcalá 80b. Epiphytic in tropical upper montane rainforest. Intra-Philippine Distribution: Widespread (Tan and Iwatsuki, 1991).
- Leucoloma molle* (Müll. Hal.) Mitt. [Dicranaceae] – SITE 7: A.A. Alcalá 84 & 85. Epiphytic in tropical upper montane rainforest. Intra-Philippine Distribution: Batan, Luzon, Catanduanes, Mindoro, Palawan, Negros, Camiguin, and Mindanao (Linis, 2009, 2010, 2014; Tan, 1996; Tan and Iwatsuki, 1991).
- Leucophanes octoblepharioides* Brid. [Calymperaceae] – SITE 3: A.A. Alcalá 59 & 60. Epiphytic in tropical lowland evergreen rainforest. Intra-Philippine Distribution: Batan, Luzon, Catanduanes, Mindoro, Palawan, Panay, Camiguin, and Mindanao (Linis, 2009, 2010, 2014; Tan, 1996; Tan and Iwatsuki, 1991).
- Neckeromnion lepineanum* (Mont.) S. Olsson, Enroth, Huttunen & D. Quandt [Neckeraceae] – SITE 2: A.A. Alcalá 52. Epiphytic in tropical lowland evergreen rainforest. Intra-Philippine Distribution: Widespread (Tan and Iwatsuki, 1991 as *Neckeropsis lepineana*).
- Pelekium bifarium* (Bosch. & Sande Lac.) M. Fleisch. [Thuidiaceae] – SITE 6: A.A. Alcalá 76. SITE 7: A.A. Alcalá 81. Epiphytic or epixylic in tropical lower and tropical upper montane rainforest. Intra-Philippine Distribution: Luzon, Mindoro, Palawan, and Panay (Linis, 2009 as *Aequatoriella bifaria*; Tan, 1996 as *Lorentzia bifaria*; Tan and Iwatsuki, 1991 as *L. bifaria*).
- Pelekium velatum* Mitt. [Thuidiaceae] – SITE 2: A.A. Alcalá 51. SITE 4: A.A. Alcalá 65. Epiphytic in tropical lowland evergreen rainforest. Intra-Philippine Distribution: Widespread (Tan and Iwatsuki, 1991 as *Lorentzia velata*).
- Pyrrhobryum spiniforme* (Hedw.) Mitt. [Rhizogoniaceae] – SITE 5: A.A. Alcalá 71. Epiphytic in tropical lowland evergreen rainforest. Intra-Philippine Distribution: Widespread (Tan and Iwatsuki, 1991).
- Rhaphidostichum luzonense* (Broth.) Broth. [Sematophyllaceae] – SITE 8: A.A. Alcalá 92. Epiphytic in tropical upper montane rainforest. Intra-Philippine Distribution: Luzon and Mindanao (Shevock and Yorong, 2018; Tan and Iwatsuki, 1991).
- Rhynchostegiella menadensis* (Sande Lac.) E.B. Bartram [Brachytheciaceae] – SITE 2: A.A. Alcalá 54 & 55. SITE 3: A.A. Alcalá 61. Epiphytic in tropical lowland evergreen rainforest. Intra-Philippine Distribution: Luzon, Palawan, and Mindanao (Shevock and Yorong, 2018; Tan, 1996; Tan and Iwatsuki, 1991).
- Rhynchostegium celebicum* (Sande Lac.) A. Jaeger [Brachytheciaceae] – SITE 6: A.A. Alcalá 78. Epiphytic in tropical lower montane rainforest. Intra-Philippine Distribution: Luzon, Mindoro, and Mindanao (Linis, 2009; Tan and Iwatsuki, 1991).
- \**Syrhropodon aristifolius* Mitt. [Calymperaceae] – SITE 5: A.A. Alcalá 73. Epiphytic in tropical lowland evergreen rainforest. Intra-Philippine Distribution: Heretofore known only from Mindoro and Mindanao (Ellis and Tan, 1999; Tan and Iwatsuki, 1991).
- Taxithelium instratum* (Brid.) Broth. [Pylaisiadelphaceae] – SITE 3: A.A. Alcalá 62. Epiphytic in tropical lowland evergreen rainforest. Intra-Philippine Distribution: Widespread (Tan and Iwatsuki, 1991).
- Taxithelium lindbergii* (A. Jaeger) Renaud & Cardot

[Pylaisiadelphaceae] – SITE 4: A.A. Alcala 70. Epiphytic in tropical lowland evergreen rainforest. Intra-Philippine Distribution: Luzon, Mindoro, Camiguin, and Mindanao (Cámara, 2011; Linis, 2019; Tan and Iwatsuki, 1991 as *Taxithelium alare*).

## DISCUSSION

Restating, a total of 27 species, one (1) subspecies, and one (1) variety in 22 genera and 10 families were

found in the Molawin-Dampalit Subwatershed of the Mt. Makiling Forest Reserve (Table 2). These represent 12.16% of the species, 22% of the genera, and 31.25% of the families reported for the mountain and its vicinity. The most represented families are Neckeraceae with five (5) species and Calymperaceae, Pylaisiadelphaceae, and Sematophyllaceae each with three (3) species. Among genera, *Acroporium*, *Fissidens*, *Leucobryum*, *Pelekium*, and *Taxithelium* are represented each with two (2) species.

**Table 2.** Summary of mosses found in the Molawin-Dampalit Subwatershed, Mt. Makiling Forest Reserve.

Families	Genera	Species and subspecies
Brachytheciaceae	two (2)	two (2) species
Calymperaceae	three (3)	three (3) species
Dicranaceae	one (1)	one (1) species
Fissidentaceae	one (1)	two (2) species and one (1) variety
Hypnodendraceae	one (1)	one (1) species and subspecies
Leskeaceae	one (1)	one (1) species
Leucobryaceae	one (1)	two (2) species
Meteoriaceae	one (1)	one (1) species
Neckeraceae	five (5)	five (5) species
Pylaisiadelphaceae	two (2)	three (3) species
Rhizogoniaceae	one (1)	one (1) species
Sematophyllaceae	two (2)	three (3) species
Thuidiaceae	one (1)	two (2) species
Total	22 genera	27 species, one (1) subspecies, and one (1) variety

Among the species recorded in the Molawin-Dampalit Subwatershed, the presence of *Syrrhopodon aristifolius* is noteworthy and is the first record of this species in Luzon Island. This Indo-Pacific species have been

previously recorded from Mindoro and Mindanao and is distinguished from other species of Calymperaceae by its linear and bristle-like leaves, poorly defined extent of the hyaline lamina, and the polystratose and usually

entire margins above leaf base (Ellis and Tan, 1999).

In light of recent studies on the moss flora of Mt. Makiling (*i.e.* Alcalá *et al.*, 2020; Magcale-Macandog *et al.*, 2022), it is apparent that further explorations on the least studied areas of the forest reserve will likely provide more information on the local moss flora and their distribution. Magcale-Macandog *et al.* (2022) found 24 species of mosses belonging to 19 families thriving in the northeastern slope of the mountain. In addition, they found out that species composition of plots sampled varied along an elevational gradient—forming three (3) clusters, of which high elevation areas (900 to 1000 m asl) form a distinct assemblage. To complement this, we include *Acroporium hermaphroditum*, *A. secundum*, *Leucobryum bowringii*, *Leucoloma molle*, and *Rhaphidostichum luzonense* as elements that characterize the high elevation areas of Mt. Makiling. Similarly, *Caduciella mariei*, *Circulifolium exiguum*, *Himantocladium plumula*, *Isopterygium albescens*, *Neckeromnion lepineanum*, and *Syrrhopodon aristifolius* are among the species found along the lower elevation of the northeastern slope of the mountain. Given these, we emphasize the need to conduct additional fieldworks within the forest reserve (*e.g.* in the Cambantoc Subwatershed and Crater Area) to gather additional data on the moss flora and its distribution which would provide essential information for a revised checklist of the mosses to be realized.

## ACKNOWLEDGEMENTS

The authors would like to express their gratitude to: the Makiling Center for Mountain Ecosystems, UPLB for granting access and permit to conduct the research in MMFR; Prof. A.S. Hadsall and Ms. M.A. San Pascual for their assistance during the herbarium work at CAHUP; and Z.D. Meneses-Adorador and J.T. Adorador for their assistance and companionship in the field. This research is financially supported by the Department of Science and Technology (DOST) of the Philippines through the ASTHRDP-NSC scholarship granted to AAA.

## REFERENCES

- Alcalá, A.A., M.L. Castillo, P.L. Malabrigo Jr., E.L.R. Logatoc, B.O. Pasion and R.S. Alcalá. 2020. Mosses from the Greater Sipit Subwatershed, Mt. Makiling Forest Reserve, the Philippines. *Journal on New Biological Reports* 9(2): 262–268.
- Azuelo, A.G., B.C. Tan, J.R. Shevock, A. Manual, A. Yorong and L.G. Sariana. 2015. Mosses new for the Mindanao Island, Republic of the Philippines. *Proceedings of the California Academy of Sciences*, Series 4, 62(4): 127–134.
- Brinda, J.C. and J.J. Atwood (eds.). 2022 [continuously updated]. *The Bryophyte Nomenclator*. Available from <https://bryonames.org>. Accessed last 13 January 2022.
- Câmara, P.E.A.S. 2011. A re-circumscription of the moss genus *Taxithelium* (Pylaisiadelphaceae) with a taxonomic revision of subgenus *Vernieri*. *Systematic Botany* 36(1): 7–21.
- Department of Environment and Natural Resources – Biodiversity Management Bureau and Deutsche Gesellschaft für Internationale Zusammenarbeit (GIZ) GmbH [DENR–BMB and GIZ]. 2017. *Manual on Biodiversity Assessment and Monitoring System for Terrestrial Ecosystems*. How-to Guidelines. Manila, Philippines. 180 pp.
- Ellis, L.T. and B.C. Tan. 1999. The moss family Calymperaceae (Musci) in the Philippines. *Bulletin of the Natural History Museum, Botany series* 29(1): 1–46.
- Hallingbäck, T. and B.C. Tan. 2010. Past and present activities and future strategy of bryophyte conservation. *Phytotaxa* 9: 266–274.
- Linis, V.C. 2009. Biogeography of Mindoro mosses. *Blumea* 54: 290–296.
- Linis, V.C. 2010. The mossflora of Camiguin Island, Philippines and their floristic relations to some adjacent islands in the archipelago. *Telopea* 12(4): 525–542.
- Linis, V.C. 2014. Biogeographical notes on the moss floras of Bicol Peninsula in Luzon and the Catanduanes Islands, the Philippines. *Philippine Journal of Science* 142: 119–133.
- Linis, V.C. 2018. A new country record and additions to the moss floras of Luzon and Mindanao islands, Philippines. *Telopea* 21:91–99.
- Linis, V.C. 2019. Notes on the diversity and floristic affinity of mosses (Musci, Bryophyta) from Zambales Mountain Range, Luzon Island, Philippines. *Phytotaxa* 388(1): 69–97.
- Linis, V.C. 2021. New Philippine moss records and additions to the Luzon moss flora. *Telopea* 24: 383–394.
- Linis, V.C. and B.C. Tan. 2013. Eight additional new records of Philippine Mosses. *Telopea* 15: 45–49.
- Lubos, L.C. 2007. New records of Philippine mosses from Mindanao Island. *Liceo Journal of Higher Education Research* 5(1): 1–8.
- Magcale-Macandog, D.B., V.G.V. Paller, N.K. Torreta, I.A.F. Lambio, A.S. Hadsall, I.E. Buot, Jr., M.D. delos Angeles, C.R. Cervancia, S.G.L. Quiñones, and J.M. Laruya. 2022. Plant diversity of Mount Makiling Forest Reserve: Implications to management and

- conservation. In: S. Ramamoorthy, I.E. Buot, Jr. and R. Chandrasekaran (eds.), *Plant Genetic Resources, Inventory, Collection and Conservation*. Springer Singapore, Singapore, pp. 97–119.
- Sastre-De Jesus, I. and B.C. Tan. 1995. Problems of bryophyte conservation in the tropics: A discussion, with case examples from Puerto Rico and the Philippines. *Caribbean Journal of Science* 31: 200–206.
- Shevock, J.R. and A.P. Yorong. 2018. New additions to the moss flora of Mindanao Island, Republic of the Philippines. *Philippine Journal of Systematic Biology* 12(1): 67–72.
- Tan, B.C. 1982. Checklist of mosses of Mt. Makiling (Luzon Island, Philippines). *Quarterly Journal of the Taiwan Museum* 35: 135–148.
- Tan, B.C. 1996. Biogeography of Palawan Mosses. *Australian Systematic Botany* 9: 193–203.
- Tan, B.C. and Z. Iwatsuki. 1991. A new annotated Philippine moss checklist. *Harvard Papers in Botany* 1(3): 1–64.
- Tan, B.C., L.C. Lubos and U. Schwarz. 2000. New and biogeographically noteworthy records of Philippine mosses from Mindanao Island. *Tropical Bryology* 18: 27–37.
- Tan, B.C. and J.R. Shevock. 2014. Noteworthy mosses and a new variety collected from the 2014 joint expedition of CAS and CMUH new to Mindanao Island of the Philippines. *Bryophyte Diversity and Evolution* 36(1): 22–30.
- Tan B.C., J.R. Shevock, F. Coritico, and V. Amoroso. 2015. Mosses new for Mindanao Island, the Philippines, III. *Bulletin of the National Museum of Science, Series B* 41(3): 91–97.
- Tan, B.C., J.R. Shevock, A.G. Azuelo and L.C. Lubos. 2017. Additions to the moss floras of Mindanao and the Philippines with a focus on the rediscovery of *Euptychium setigerum*. *Natural History Bulletin of the Siam Society* 62(1): 15–20.
- Tixier, P. 1972. Mount Maquiling bryoflora. *Gardens' Bulletin Singapore* 26: 137–153.

Received: 18 April 2021

Accepted: 11 August 2022

Published: XX XXX 2022

Published in Print: 31 December 2022