



The Number of Integers Divisible by a^k Except $a^{k+l}b$

Yanapat Tongron, Kanyaphak Paikhlaew*

*Mathematics and Applied Statistics Program, Faculty of Science and Technology,
Nakhon Ratchasima Rajabhat University, Nakhon Ratchasima 30000, Thailand*

Received 17 November 2020; Received in revised form 14 July 2021

Accepted 22 July 2021; Available online 29 September 2022

ABSTRACT

For positive integers a, b, k, l, n , let $A_{(k,l)}^{(a,b)}(n)$ be the set of all integers divisible by a^k except $a^{k+l}b$ between 1 and n . Under certain condition on these integers, we obtain the explicit formula for $|A_{(k,l)}^{(a,b)}(n)|$. We also provide bounds on $|A_{(k,l)}^{(a,b)}(n)|$ and show that these are the best bounds by examples.

Keywords: Bounds; Ceiling function; Divisibility; Floor function; Integer

1. Introduction

First, we describe the basic problem of sets in Mathematics as follows: How many integers between 1 and 100 both inclusive are divisible by 2 and 3? By the property of the floor function $\lfloor \cdot \rfloor$ defined by [1, 2].

$\lfloor x \rfloor$ is the greatest integer less than or equal to a real number x , the answer is

$$\left\lfloor \frac{100}{6} \right\rfloor = 16.$$

What happens if the question is changed from ‘and’ to ‘or’? That is “how many

integers between 1 and 100 both inclusive that are divisible by 2 or 3?”

Let

$$A = \{1 \leq n \leq 100 : 2|n\},$$

and

$$B = \{1 \leq n \leq 100 : 3|n\}.$$

We need to find $|A \cup B|$, where $|X|$ denotes the number of elements in a finite set X . This problem can be solved by the Inclusion-Exclusion Principle [3, 4] which stated that for arbitrary finite sets A and B , $|A \cup B| = |A| + |B| - |A \cap B|$. Again, we have

$$\begin{aligned}
 |A \cup B| &= \left\lfloor \frac{100}{2} \right\rfloor + \left\lfloor \frac{100}{3} \right\rfloor - \left\lfloor \frac{100}{6} \right\rfloor \\
 &= 50 + 33 - 16 \\
 &= 67.
 \end{aligned}$$

Indeed, we can always find the number of all integers divisible by an integer α or/and an integer β between 1 and a positive integer n using the above concepts.

In this work, we are interested in dealing with the similar problem on indivisibility. For positive integers a, b, k, l, n , define a finite set of positive integers $A_{(k,l)}^{(a,b)}(n)$ by

$$A_{(k,l)}^{(a,b)}(n) = \{x \leq n : a^k | x, a^{k+l} b \nmid x\}.$$

We see that $|A_{(k,l)}^{(a,b)}(n)| = 0$ if $n < a^k$ or $a^l b = 1$. Thus, only the case $n \geq a^k$ and $a^l b \neq 1$ is considered. We can find the explicit formula for $|A_{(k,l)}^{(a,b)}(n)|$ under some condition on a, b, k, l, n . Moreover, bounds on $|A_{(k,l)}^{(a,b)}(n)|$ are provided in terms of the floor function and the ceiling function $\lceil \cdot \rceil$ denoted by

$\lceil x \rceil$ is the smallest integer greater than or equal to a real number x .

Some examples to confirm that these are the best bounds are also given.

2. Some explicit formula for $|A_{(k,l)}^{(a,b)}(n)|$

In this section, we find the explicit formula for $|A_{(k,l)}^{(a,b)}(n)|$ under certain condition on a, b, k, l, n . The following

theorem is an important tool for establishing the explicit formula.

Theorem 2.1. Let a, b, k, l, n be positive integers such that $n \geq a^k$ and $a^l b \neq 1$. Then

$$\begin{aligned}
 |A_{(k,l)}^{(a,b)}(n)| - 1 + \frac{2}{a^l b} &\leq \frac{(a^l b - 1)n + a^k}{a^{k+l} b} \\
 &< |A_{(k,l)}^{(a,b)}(n)| + 1.
 \end{aligned}$$

Proof. By the property of the floor function, we have

$$a^k \left\lfloor \frac{n}{a^k} \right\rfloor \leq n < a^k \left(\left\lfloor \frac{n}{a^k} \right\rfloor + 1 \right). \tag{2.1}$$

Assume that $\lfloor n/a^k \rfloor \equiv i \pmod{a^l b}$ for some $0 \leq i < a^l b$. Write

$$\left\lfloor \frac{n}{a^k} \right\rfloor = i + ma^l b \tag{2.2}$$

for some positive integer m . Since

$$\begin{aligned}
 A_{(k,l)}^{(a,b)}(n) &= \{a^k(1), a^k(2), \dots, a^k(a^l b - 1), \\
 &\quad a^k(a^l b + 1), a^k(a^l b + 2), \dots, a^k(2a^l b - 1), \\
 &\quad a^k(2a^l b + 1), \dots, a^k(3a^l b - 1), \\
 &\quad \vdots \\
 &\quad a^k((m-1)a^l b + 1), \dots, a^k(ma^l b - 1), \\
 &\quad a^k(ma^l b + 1), \dots, a^k(ma^l b + i)\},
 \end{aligned}$$

we get

$$\begin{aligned}
 |A_{(k,l)}^{(a,b)}(n)| &= ma^l b - m + i \\
 &= (a^l b - 1)m + i.
 \end{aligned} \tag{2.3}$$

It follows from Eq. (2.1) and Eq. (2.2) together with Eq. (2.3) that

$$\begin{aligned}
 a^k(a^l b m + i) &\leq n &< a^k(a^l b m + i + 1) \\
 m + \frac{i}{a^l b} &\leq \frac{n}{a^{k+l} b} &< m + \frac{i}{a^l b} + m + \frac{1}{a^l b} \\
 (a^l b - 1)m + i - \frac{i}{a^l b} &\leq \frac{(a^l b - 1)n}{a^{k+l} b} &< (a^l b - 1)m + i - \frac{i}{a^l b} + 1 - \frac{1}{a^l b} \\
 \left| A_{(k,l)}^{(a,b)}(n) \right| - \frac{i}{a^l b} &\leq \frac{(a^l b - 1)n}{a^{k+l} b} &< \left| A_{(k,l)}^{(a,b)}(n) \right| - \frac{i}{a^l b} + 1 - \frac{1}{a^l b} \\
 \left| A_{(k,l)}^{(a,b)}(n) \right| - \frac{i}{a^l b} + \frac{1}{a^l b} &\leq \frac{(a^l b - 1)n + a^k}{a^{k+l} b} &< \left| A_{(k,l)}^{(a,b)}(n) \right| - \frac{i}{a^l b} + 1.
 \end{aligned} \tag{2.4}$$

Since

$$\begin{aligned}
 &\left| A_{(k,l)}^{(a,b)}(n) \right| - \frac{i}{a^l b} + \frac{1}{a^l b} \\
 &\geq \left| A_{(k,l)}^{(a,b)}(n) \right| - \frac{a^l b - 1}{a^l b} + \frac{1}{a^l b} \\
 &= \left| A_{(k,l)}^{(a,b)}(n) \right| - 1 + \frac{2}{a^l b},
 \end{aligned}$$

and

$$\left| A_{(k,l)}^{(a,b)}(n) \right| - \frac{i}{a^l b} + 1 \leq \left| A_{(k,l)}^{(a,b)}(n) \right| + 1,$$

the result follows from Eq. (2.4). ■

The explicit formula for $\left| A_{(k,l)}^{(a,b)}(n) \right|$ under some condition on positive integers a, b, k, l, n follows from the inequality Eq. (2.4) in the proof of Theorem 2.1 as follows:

Corollary 2.1. Let a, b, k, l, n be positive integers such that $n \geq a^k$ and $a^l b \neq 1$. Then

$$\left| A_{(k,l)}^{(a,b)}(n) \right| = \left\lfloor \frac{(a^l b - 1)n + a^k}{a^{k+l} b} \right\rfloor,$$

whenever $\lfloor n/a^k \rfloor \equiv i \pmod{a^l b}$ with $0 \leq i \leq 1$.

Proof. If $0 \leq i \leq 1$, then we have

$$\left| A_{(k,l)}^{(a,b)}(n) \right| - \frac{i}{a^l b} + \frac{1}{a^l b} \geq \left| A_{(k,l)}^{(a,b)}(n) \right|$$

and

$$\left| A_{(k,l)}^{(a,b)}(n) \right| - \frac{i}{a^l b} + 1 \leq \left| A_{(k,l)}^{(a,b)}(n) \right| + 1.$$

It follows from Eq. (2.4) that

$$\begin{aligned}
 \left| A_{(k,l)}^{(a,b)}(n) \right| &\leq \frac{(a^l b - 1)n + a^k}{a^{k+l} b} \\
 &< \left| A_{(k,l)}^{(a,b)}(n) \right| + 1.
 \end{aligned}$$

The proof is complete by the property of the floor function. ■

The next corollary follows from Corollary 2.1 and the fact that $a^l b \leq 2$ implies $\lfloor n/a^k \rfloor \equiv i \pmod{a^l b}$ such that $i = 0, 1$.

Corollary 2.2. Let a, b, k, l, n be positive integers such that $n \geq a^k$ and $a^l b \neq 1$. If $a^l b \leq 2$, then

$$\left| A_{(k,l)}^{(a,b)}(n) \right| = \left\lfloor \frac{(a^l b - 1)n + a^k}{a^{k+l} b} \right\rfloor.$$

The number of integers divisible by 2^k except 2^{k+1} between 1 and n can be established from Corollary 2.2 as the following example:

Example 2.3. Let n, k be positive integers, $a=2, b=1$ and $l=1$. Since $a^l b \leq 2$, Corollary 2.2 implies

$$\left| A_{(k,1)}^{(2,1)}(n) \right| = \left\lfloor \frac{n+2^k}{2^{k+1}} \right\rfloor.$$

Observe that for positive integers n and k , if $a=1, b=2$ and $l=1$, then we have from Corollary 2.2 that $\left| A_{(k,1)}^{(1,2)}(n) \right|$ is the number of odd integers between 1 and n , i.e.,

$$\left| A_{(k,1)}^{(1,2)}(n) \right| = \left\lfloor \frac{n+1}{2} \right\rfloor.$$

3. Bounds on $\left| A_{(k,l)}^{(a,b)}(n) \right|$

In this section, bounds on $\left| A_{(k,l)}^{(a,b)}(n) \right|$ are given by using Theorem 2.1 as follows:

Theorem 3.1. Let a, b, k, l, n be positive integers such that $n \geq a^k$ and $a^l b \neq 1$. Then

$$\left\lceil \frac{(a^l b - 1)n + a^k}{a^{k+l} b} - 1 \right\rceil \leq \left| A_{(k,l)}^{(a,b)}(n) \right| \leq \left\lfloor \frac{(a^l b - 1)n + a^k}{a^{k+l} b} + 1 - \frac{2}{a^l b} \right\rfloor.$$

Proof. From Theorem 2.1, we have

$$\left| A_{(k,l)}^{(a,b)}(n) \right| \leq \frac{(a^l b - 1)n + a^k}{a^{k+l} b} + 1 - \frac{2}{a^l b}$$

and

$$\left| A_{(k,l)}^{(a,b)}(n) \right| > \frac{(a^l b - 1)n + a^k}{a^{k+l} b} - 1.$$

These complete the proof by the property of the floor function and the ceiling function. ■

Finally, we show that Theorem 3.1 contains the perfect bounds. The following

examples show that the equality holds in Theorem 3.1 for some integer inputs.

Example 3.2. Taking $a=2, b=5, k=1, l=1$, and $n=100$; we obtain

$$\begin{aligned} A_{(k,l)}^{(a,b)}(n) &= \{1 \leq x \leq 100 : 2|x, 20 \nmid x\} \\ &= \{2, 4, 6, \dots, 16, 18, \\ &22, 24, 26, \dots, 36, 38, \\ &42, 44, 46, \dots, 56, 58, \\ &62, 64, 66, \dots, 76, 78, \\ &82, 84, 86, \dots, 96, 98\}. \end{aligned}$$

Hence, $\left| A_{(k,l)}^{(a,b)}(n) \right| = 45$. Since

$$\frac{(a^l b - 1)n + a^k}{a^{k+l} b} - 1 = \frac{(10 - 1)100 + 2}{20} - 1 = 44.1,$$

it follows that

$$\left| A_{(k,l)}^{(a,b)}(n) \right| = \left\lceil \frac{(a^l b - 1)n + a^k}{a^{k+l} b} - 1 \right\rceil.$$

Similarly, it is easy to see that

$$\left| A_{(k,l)}^{(a,b)}(n) \right| = \left\lfloor \frac{(a^l b - 1)n + a^k}{a^{k+l} b} + 1 - \frac{2}{a^l b} \right\rfloor.$$

Example 3.3. Taking $a=3, b=2, k=1, l=2$, and $n=120$; we obtain

$$\begin{aligned} A_{(k,l)}^{(a,b)}(n) &= \{1 \leq x \leq 120 : 3|x, 54 \nmid x\} \\ &= \{3, 6, 9, \dots, 24, 27, \\ &30, 33, 36, \dots, 51, 57, \\ &60, 63, 78, \dots, 81, 84, \\ &87, 90, 105, \dots, 111, 114, \\ &117, 120\}. \end{aligned}$$

Hence, $\left| A_{(k,l)}^{(a,b)}(n) \right| = 38$. Since

$$\frac{(a^l b - 1)n + a^k}{a^{k+l} b} + 1 - \frac{2}{a^l b} = \frac{(18 - 1)120 + 3}{54} + 1 - \frac{2}{18} = 38.72,$$

it follows that

$$\left| A_{(k,l)}^{(a,b)}(n) \right| = \left\lfloor \frac{(a^l b - 1)n + a^k}{a^{k+l} b} + 1 - \frac{2}{a^l b} \right\rfloor.$$

4. Conclusion

Let a, b, k, l, n be positive integers such that $n \geq a^k$ and $a^l b \neq 1$. Denote a finite set of positive integers $A_{(k,l)}^{(a,b)}(n)$ by

$$A_{(k,l)}^{(a,b)}(n) = \{x \leq n : a^k \mid x, a^{k+l} b \nmid x\}.$$

From Corollary 2.1, we have some explicit formula for $\left| A_{(k,l)}^{(a,b)}(n) \right|$ as follows:

$$\left| A_{(k,l)}^{(a,b)}(n) \right| = \left\lfloor \frac{(a^l b - 1)n + a^k}{a^{k+l} b} \right\rfloor,$$

whenever $\lfloor n/a^k \rfloor \equiv i \pmod{a^l b}$ with $0 \leq i \leq 1$. Notice that $a^l b \leq 2$ implies $\lfloor n/a^k \rfloor \equiv i \pmod{a^l b}$ such that $i = 0, 1$. The best bounds on $\left| A_{(k,l)}^{(a,b)}(n) \right|$ are provided in Theorem 3.1, i.e.,

$$\begin{aligned} \left\lfloor \frac{(a^l b - 1)n + a^k}{a^{k+l} b} - 1 \right\rfloor &\leq \left| A_{(k,l)}^{(a,b)}(n) \right| \\ &\leq \left\lfloor \frac{(a^l b - 1)n + a^k}{a^{k+l} b} + 1 - \frac{2}{a^l b} \right\rfloor \end{aligned}$$

Our future work is to generalize these results, for example, the number of integers divisible by a except ab .

References

- [1] Niven I, Zuckerman HS, Montgomery HL. An Introduction to the Theory of Numbers. Canada: John Wiley & Sons, Inc.; 1991.
- [2] Buton DM. Elementary Number Theory. New Yor: McGraw-Hill Companies, Inc.; 2007.
- [3] Farlow SJ. Finite Mathematics and its Applications. 2nd ed. Hightstown, NJ: McGraw-Hill Companies, Inc.; 1994.
- [4] Calvert J, Voxman W. Finite Mathematics with Computer- Supported Applications. Hightstown, NJ: McGraw- Hill Companies, Inc.; 1994.