

## การเพิ่มการผลิตชี-ไฟโคไซยานินและพอลิไฮดรอกซีบิวทิเรตในอาร์โทสไปรา พลาแทนซิส ที่เพาะเลี้ยงด้วยไดโอดเปล่งแสง (แอลอีดี)

ฉันทชนก ดวงศรี<sup>1</sup> วิทยุ จุญญไชยพิพัฒน์<sup>1</sup> คมสัน สัจจะสถาพร<sup>1</sup> อุเทน พรหมอริยะ<sup>1</sup>  
นาฏอหงค์ หมุดธรรม<sup>1</sup> ณัฐพงศ์ อัคริมจิรโชติ<sup>2</sup> สุชนิทธิ์ งามกาละ<sup>1</sup> และวุฒินันท์ รักษาจิตร<sup>1\*</sup>

<sup>1</sup>โปรแกรมเทคโนโลยีสุขภาพสัตว์ คณะเทคนิคการสัตวแพทย์ และ <sup>2</sup>ภาควิชาสรีรวิทยา คณะสัตวแพทยศาสตร์  
มหาวิทยาลัยเกษตรศาสตร์ กรุงเทพฯ 10900  
\*E-mail: cvtwnr@ku.ac.th

รับบทความ: 6 กุมภาพันธ์ 2565 แก้ไขบทความ: 22 กุมภาพันธ์ 2565 ยอมรับตีพิมพ์: 3 มีนาคม 2565

### บทคัดย่อ

ไซยาโนแบคทีเรีย อาร์โทสไปรา พลาแทนซิส เป็นทรัพยากรที่มีศักยภาพสูงในการผลิตสารประกอบที่มีมูลค่า เช่น ชี-ไฟโคไซยานิน (ซี-พีซี) และ พอลิไฮดรอกซีบิวทิเรต (พีเอชบี) การศึกษาครั้งนี้มีวัตถุประสงค์เพื่อประเมินผลกระทบของไดโอดเปล่งแสง (แอลอีดี) (แสงสีแดง สีเขียว สีน้ำเงิน และสีขาว) ภายใต้ความเข้มแสงต่ำ ปานกลางและสูง (LL, ML, HL; 15, 30, 60 ไมโครโมลโฟตอนต่อตารางเมตรต่อวินาที ตามลำดับ) ต่อปริมาณชีวมวล คลอโรฟิลล์-เอ ซี-พีซี และพีเอชบีของอาร์โทสไปรา พลาแทนซิส อีกทั้งประเมินแอกติวิตีการกำจัดอนุมูลอิสระของซี-พีซีของอาร์โทสไปรา พลาแทนซิส ผลการศึกษาพบว่า ค่าสูงสุดของปริมาณชีวมวล ( $2.80 \pm 0.11$  กรัมต่อลิตร) และปริมาณคลอโรฟิลล์-เอ ( $5.80 \pm 0.12$  มิลลิกรัมต่อลิตร) พบในเซลล์ที่เพาะเลี้ยงภายใต้แสงสีแดงที่มีความเข้มแสงปานกลาง ในขณะที่ค่าต่ำสุดของปริมาณชีวมวลและปริมาณคลอโรฟิลล์-เอ พบในเซลล์ที่เพาะเลี้ยงภายใต้แสงสีน้ำเงินที่มีความเข้มแสงปานกลาง ในทางกลับกัน ซี-พีซี มีปริมาณเพิ่มขึ้นเท่ากับ  $2.78 \pm 0.07$  มิลลิกรัมต่อกรัม ในเซลล์ที่เพาะเลี้ยงภายใต้แสงสีน้ำเงินที่มีความเข้มแสงสูง เมื่อเทียบกับแสงสีแดง สีเขียว และสีขาวที่มีความเข้มแสงเดียวกัน นอกจากนี้พบว่า ซี-พีซี ที่ความเข้มข้น 100 ไมโครกรัมต่อมิลลิกรัมซึ่งสกัดได้จากเซลล์ที่เพาะเลี้ยงภายใต้แสงสีน้ำเงินที่มีความเข้มแสงสูง แสดงค่าแอกติวิตีการกำจัดอนุมูลอิสระเท่ากับร้อยละ  $52.1 \pm 5.6$  นอกจากนี้พีเอชบีมีปริมาณสูงสุด ( $9.18 \pm 0.32\%$ (w/w) DW) ในเซลล์ที่เพาะเลี้ยงภายใต้แสงสีขาวที่มีความเข้มแสงสูงและขาดแหล่งอาหารไนโตรเจนในอาหารเลี้ยง เมื่อเทียบกับแสงสีแดง สีเขียว และสีน้ำเงินที่มีความเข้มแสงเดียวกัน การศึกษาครั้งนี้ให้ข้อมูลทางสรีรวิทยาที่เป็นประโยชน์เกี่ยวกับผลกระทบของสเปกตรัมแสงที่ต่างกันและความเข้มของแสงต่อการเจริญเติบโต ปริมาณรงควัตถุสังเคราะห์ด้วยแสง และพีเอชบี ซึ่งเป็นแนวทางเบื้องต้นในการปรับการผลิตสารประกอบที่มีมูลค่าสูงจากการเพาะเลี้ยงอาร์โทสไปรา พลาแทนซิส ให้เหมาะสม

คำสำคัญ: อาร์โทสไปรา พลาแทนซิส ชี-ไฟโคไซยานิน แอลอีดี พอลิไฮดรอกซีบิวทิเรต

## Enhanced Production of C-Phycocyanin and Polyhydroxybutyrate in the *Arthrospira platensis* Cultured with Light-Emitting Diodes (LEDs)

Chanchanok Duangsri<sup>1</sup>, Winai Charunchaipipat<sup>1</sup>, Khomson Satchasataporn<sup>1</sup>,  
Authen Promariya<sup>1</sup>, Nat-Anong Mudtham<sup>1</sup>, Nattaphong Akrimajirachote<sup>2</sup>,  
Suchanit Ngamkala<sup>1</sup> and Wuttinun Raksajit<sup>1\*</sup>

<sup>1</sup>Program of Animal Health Technology, Faculty of Veterinary Technology, and <sup>2</sup>Department of Physiology, Faculty of Veterinary Medicine, Kasetsart University, Bangkok 10900, Thailand

\*E-mail: cvtwnr@ku.ac.th

Received: 6 February 2022 Revised: 22 February 2022 Accepted: 3 March 2022

### Abstract

The cyanobacterium *Arthrospira platensis* is considered a highly potential resource for the production of valuable compounds such as C-phycocyanin (C-PC) and poly-3-hydroxybutyrate (PHB). This study aimed to evaluate the effects of different coloured light-emitting diodes (LEDs) (red, green, blue, and white light) under low, medium, and high light intensity (LL, ML, HL; 15, 30, 60  $\mu\text{mol photons m}^{-2}\text{s}^{-1}$ , respectively) on the biomass, chlorophyll-*a* (Chl-*a*), C-PC, and PHB contents of *A. platensis*, and further determine the free-radical scavenging activity of *A. platensis* C-PC. The results exhibited that the highest values for biomass ( $2.80 \pm 0.11 \text{ g L}^{-1}$ ) and chlorophyll-*a* (Chl-*a*) content ( $5.80 \pm 0.12 \text{ mg L}^{-1}$ ) were obtained in cells cultured under ML red light, while the lowest values for biomass and Chl-*a* content were observed in cells cultured under ML blue light. On the other hand, the C-PC value increased to  $2.78 \pm 0.07 \text{ mg g}^{-1}$  in cells cultured under HL blue light compared to red, green, and white lights with equal intensities. Besides, the purified C-PC obtained from cells cultured under HL blue light showed  $52.1 \pm 5.6\%$  of free-radical scavenging activity at a C-PC concentration of  $100 \mu\text{g mL}^{-1}$ . In addition, the maximum PHB content ( $9.18 \pm 0.32\%$ (w/w) DW) was obtained in cells cultured under HL white light with nitrogen deprivation compared to red, green, and blue lights with equal intensities. The results of the current study provide useful physiological information regarding the effects of different light spectra and light intensity on growth, photosynthetic pigments, and PHB as a prerequisite to optimize the production of high-value compounds from cultures of *A. platensis*.

**Keywords:** *Arthrospira platensis*, C-Phycocyanin, LEDs, Polyhydroxybutyrate

## Introduction

*Arthrospira platensis* is a filamentous cyanobacterium that is rich in valuable pigments like chlorophyll-*a*, carotenoids, phycobiliproteins (PBPs), and biopolymers (Lupatini *et al.*, 2017; Milia *et al.*, 2022; Nematollahi *et al.*, 2020). *A. platensis* biomass can be applied in the food sector for animal and human consumption. PBPs are major pigment-protein components of the cyanobacterial cells and can constitute up to 50% of the cell materials. PBPs are divided into four types according to their maxima light absorption wavelength: phycocyanin ( $\lambda=615\text{--}640$  nm), phycoerythrin ( $\lambda=565\text{--}575$  nm), phycoerythrocyanin ( $\lambda=560\text{--}600$  nm), and allophycocyanin ( $\lambda=650\text{--}660$  nm). Among them, phycocyanin (C-PC) is a hydrophilic, blue-coloured fluorescent protein covalently bound to chromophore tetrapyrrole groups (Minic *et al.*, 2018). It is a pigment associated with other phycobiliproteins, such as allophycocyanin and phycoerythrin. Furthermore, it is a water-soluble compound with constituents of 17% proteins (Patel *et al.*, 2005). Besides, C-PC has been shown to exhibit antioxidant, anti-inflammation, anticancer, and immunity enhancement properties (García *et al.*, 2021; Sonani *et al.*, 2017). Phycocyanin from *S. platensis* LEB-52 can enrich nutritional value and provide a natural colour to dairy products (Campos Assumpção de Amarante *et al.*, 2020). In addition, *A. platensis* have a biological system with the potential to synthesize biopolymers

such as poly-3-hydroxybutyrate (PHB) under various culture conditions (Duangsri *et al.*, 2020). The highest accumulation of PHB in *Arthrospira platensis* was observed in cells grown photoautotrophically under nitrogen deprivation with acetate supplementation (Duangsri *et al.*, 2020). Most PHB is produced and accumulated intracellularly as a carbon storage material and an energy reservoir. PHB is an attractive alternative to common thermoplastics due to its hydrophobicity, biodegradability, and biocompatibility. Furthermore, PHB nanofiber from *Spirulina platensis* has exhibited properties equal to or better than nanofiber made with commercially available PHB (Moradi *et al.*, 2021). The accumulation of photosynthetic pigments, PHB, and biomass in *A. platensis* can be achieved by optimizing the availability of growth factors such as light, temperature, pH, and nutrients. Particularly, light quality affects the growth and biochemical properties of phototrophic organisms (Khatoon *et al.*, 2018). Recently, studies have focused on investigating the effects of light intensity and light quality on microalgal growth, pigmentation, and photosynthesis (Klepacz-Smółka *et al.*, 2020; Wicaksono *et al.*, 2019). Various light sources such as sunlight, fluorescent light, and light-emitting diodes (LEDs) have been used to produce biomass and valuable biochemical components (da Fontoura Prates *et al.*, 2020). In practice, artificial illumination LEDs tend to be the preferred choice for

acclimatizing a cultivation system into an intensive one in a small area. LEDs emit a single wavelength of light and have low power consumption interpreted to low carbon emissions (Van Hieu *et al.*, 2021). Notably, LED lights produce less of an environmental impact than traditional lighting sources. Therefore, LEDs are considered an attractive artificial light for microalgal production. Even though many studies have focused on various lighting conditions during the cultivation of *Arthrospira* sp. to enhance biomass and C-PC production, little or no attention has been given to the co-production of C-PC and PHB using LEDs during cultivation. Hence, this research aimed to investigate the effects of different LEDs (white, red, green and blue light) on *A. platensis* biomass, photosynthetic pigments (chlorophyll-*a*, carotenoid, and C-PC), and PHB content, as well as further determine the *in vitro* free-radical scavenging activity of *A. platensis* C-PC.

## Research Methodology

### Cell cultivation and biomass determination

The *A. platensis* used in this study was obtained from the Institute of Food Research and Product Development, Kasetsart University (IFRPD). The cells were cultivated in 250-mL Erlenmeyer flasks (Schott Duran, Germany) containing 50 mL Zarrouk medium under photoautotrophic conditions at  $32 \pm 0.5^\circ\text{C}$ , pH 10.0. Experiments for the two-stage cul-

tivation were conducted as follows. In the first stage, *A. platensis* was cultivated under continuous illumination with white fluorescent at  $30 \mu\text{mol photons m}^{-2}\text{s}^{-1}$  for 7 days until reaching the late logarithm phase (optical density, OD~1–1.2). After that, white fluorescent was replaced with blue, green, red and white LEDs with wavelengths of 430–465 nm, 525–550 nm, 630–665 nm, and 420–665 nm, respectively. The *A. platensis* was further cultivated under three light intensities designated as low-intensity (LL,  $15 \mu\text{mol photons m}^{-2}\text{s}^{-1}$ ), medium-intensity (ML,  $30 \mu\text{mol photons m}^{-2}\text{s}^{-1}$ ), and high-intensity (HL,  $60 \mu\text{mol photons m}^{-2}\text{s}^{-1}$ ) for 5 days. The biomass concentration ( $\text{g L}^{-1}$ ) was determined spectrophotometrically according to Duangsri *et al.*, (2020). The irradiance of the LEDs was measured using a quantum meter (Biospherical Instruments, model QSL-100).

### C-phycoerythrin extraction, partial purification and determination

The *A. platensis* cells were harvested by filtration and then suspended in a 0.1 M sodium phosphate buffer (pH 7.0) at a ratio of 1:15 under darkness on ice, while cells were extracted by sonication at 20 kHz for 10 min with three repeated cycles of freezing and thawing (RFT cycles). The suspension was frozen for 3 hrs at  $-20^\circ\text{C}$  and thawed for 1 hr at  $25^\circ\text{C}$ . After 3×RFT cycles, the mixture containing C-phycoerythrin (C-PC) was centri-

fused for 30 min at 4,000×g at 4°C to remove the cell residues. Partial purification of C-PC was done using ammonium sulfate precipitation. Solid ammonium sulfate was gradually added into the beaker containing crude extracts of C-PC to obtain 40% saturation with continuous stirring for 1 hr. The solution was stored overnight at 4°C under dark conditions, and the precipitation was collected by centrifugation at 15,000×g for 15 min at 4°C. The clear supernatant was discarded, and the blue precipitate was dissolved in 0.1M PBS pH 7.0 then stored at 4°C in an amber bottle until examination (Kumar *et al.*, 2014). After that, the C-PC contents were calculated using Eq. (1) as described by Bennett and Bogorad (1973).

$$\text{C-PC (mg g}^{-1}\text{)} = (A_{620} - 0.474 \times A_{652})/5.34 \quad (1)$$

where  $A_{620}$  and  $A_{652}$  are the optical density at 620 and 652 nm, respectively, for phycocyanin.

#### **Chlorophyll-a extraction and determination**

The chlorophyll-a contents ( $\text{mg L}^{-1}$ ) were determined according to MacKinney (1941). Briefly, one milliliter of cell culture was centrifuged at 8,000×g for 10 min. The biomass was resuspended in 1 mL of 99.9% methanol. The samples were incubated under darkness overnight. The upper aqueous layer was kept and determined by a spectrophotometer at 665 nm.

#### **PHB induction and quantification**

The microalgal biomass of *A. platensis* (7 days) was harvested and washed twice with fresh nitrogen-free Zarrouk medium ( $\text{ZN}_0$ ), and inoculated into 50 mL of  $\text{ZN}_0$  followed by cultivation under blue, green, red, and white LEDs with low-intensity (LL, 15  $\mu\text{mol photons m}^{-2} \text{s}^{-1}$ ), medium-intensity (ML, 30  $\mu\text{mol photons m}^{-2} \text{s}^{-1}$ ), and high-intensity (HL, 60  $\mu\text{mol photons m}^{-2} \text{s}^{-1}$ ) for 5 days. PHB was quantified by high-performance liquid chromatography (HPLC, Waters Corp., USA) using a C18 reverse-phase column (I.D. 4.6×150 nm) with a UV detector set at 210 nm (Duangsri *et al.*, 2020). Briefly, dry cells were boiled at 100°C for 60 min with  $\text{H}_2\text{SO}_4$ . Boiled samples were filtered using a 0.45  $\mu\text{m}$  polypropylene membrane filter. The acid-hydrolyzed PHB (crotonic acid) monomer was detected using HPLC. The isocratic solvent system was run at 40% (v/v) of acetonitrile and 60%(v/v) of 0.1% acetic acid with a flow rate of 0.6  $\text{mL min}^{-1}$ . Authentic commercial PHB from Sigma-Aldrich (USA) was used as a standard which was prepared similarly to that of the cell sample. The unit of PHB content was % PHB weight per biomass dry weight (%(w/w DW)).

#### **Antioxidant activity using DPPH assay**

DPPH (2,2-diphenyl picrylhydrazyl)

free radical scavenging activity was studied using the method described by Roy and Pabbi (2022). Briefly, two milliliters of samples were mixed with 1 mL of DPPH solution (200  $\mu$ M in ethanol), incubated at room temperature in the dark for 30 min followed by measuring the absorbance of the samples at 517 nm. The DPPH reduction percentage was determined using Eq. (2), where  $A_i$  is the initial absorbance and  $A_f$  is the absorbance after incubation for 30 min.

$$\text{DPPH free-radical scavenging activity (\%)} \\ = (A_i - A_f)/A_i \times 100 \quad (2)$$


### Statistical analysis

All data obtained in this study represent the means of three independent biological replicates, and the error bars represent the standard deviation (Mean  $\pm$  SD,  $n = 3$ ). The statistical analysis was evaluated by one-way analysis of variance (ANOVA) and the significant difference ( $p < 0.05$ ) was compared by Duncan's multiple range tests using SPSS version 22 (IBM, USA).

### Results and Discussion

One of the important requirements for cyanobacteria metabolism is light. Light intensity directly affects the rate of photosynthesis. Cyanobacteria grown under various light intensities show changes in biomass produc-

tion, photosynthetic pigments, and lipid contents (Gim *et al.*, 2016; Maltsev *et al.*, 2021; McGee *et al.*, 2020). The highest biomass content ( $2.80 \pm 0.11 \text{ g L}^{-1}$ ) was obtained under ML red LED, which was significantly higher than LL red LED ( $p < 0.05$ ), and higher than ML white, ML green and ML blue LEDs (Figure 1A). Previously, it was found that the highest growth rate was obtained with red LED in batch cultures of *A. platensis* (Markou, 2014), while the lowest biomass productivity was obtained with ML blue LED. In addition, the red LED showed the most effective emission for *S. platensis* growth, followed by white, yellow and green LEDs with a similar growth pattern (Chen *et al.*, 2010). Furthermore, the levels of chlorophyll-*a* in *A. platensis* under ML intensities of all colour LEDs were significantly higher than that cells under HL and LL intensities about 0.7- to 1.5-fold, respectively ( $p < 0.05$ ). The highest chlorophyll-*a* content ( $5.80 \pm 0.12 \text{ mg L}^{-1}$ ) was obtained under ML red LED (Figure 1B). In particular, the chlorophyll-*a* levels obtained under red and white LEDs were significantly higher than those reached under green and blue illumination for ML conditions ( $p < 0.05$ ). Additionally, C-phycocyanin (C-PC) content ( $2.55 \pm 0.09 \text{ mg g}^{-1}$ ) was 1.7-fold higher under LL red LED compared to ML red LED, and significantly higher than those reached




**Figure 1** Effects of light colour and light intensity on biomass (A) chlorophyll-a (B) and C-PC (C) of *A. platensis*. Cells were cultivated under three light intensities designated as low-intensity (LL, 15  $\mu\text{mol photons m}^{-2} \text{s}^{-1}$ ), medium-intensity (ML, 30  $\mu\text{mol photons m}^{-2} \text{s}^{-1}$ ), and high-intensity (HL, 60  $\mu\text{mol photons m}^{-2} \text{s}^{-1}$ ) for 5 days. Bar graphs represent mean values ( $\pm$  SD) of three independent experiments. Different letters (a, b, or c) indicate significant differences ( $p < 0.05$ ) between the groups obtained from cells cultured in LL, ML and HL.

under green and blue illumination for LL and ML conditions ( $p < 0.05$ ) (Figure 1C). Remarkably, the highest value for C-PC increase ( $2.78 \pm 0.07 \text{ mg g}^{-1}$ ) was obtained in cells cultured under HL blue light compared to red and white lights. Chen *et al.* (2010) reported that the chlorophyll-a and C-PC contents of *S. platensis* under green and blue LEDs were significantly lower than those produced under the red, white and yellow LEDs. On the other hand, it was reported that the highest pigment production rate for *Spirulina fussiformis* was obtained under blue light (Madhyastha and Vatsala, 2007). A possible explanation is that the authors used other light sources instead of LEDs to obtain the desired colour. All results demonstrated that this cyanobacterium absorbed blue light much less efficiently than white and red light. These findings are consistent with previous investigations conducted with *A. platensis* (Markou, 2014; Wang *et al.*, 2007). Our results support the hypothesis that the blue LED creates an imbalance between photosystems I and II, with an excess of energy capture at the PSI side and a lack at the PSII side of the photosynthetic electron transport chain of *A. platensis* (Luimstra *et al.*, 2018). Moreover, cyanobacteria  $\beta$ -carotene, which absorbs blue light efficiently, is more abundant in PSI than in PSII, hence further contributing

to the photosynthesis light-harvesting capacity of PSI. Our results showed that *A. platensis* cells acclimate to blue light by enhancing the production of phycocyanin in an attempt to restore the balance between photosystems I and II. These findings corroborate the notion that a culture grown under blue light has a much lower growth rate than red and white lights, but a significantly higher phycocyanin content. Moreover, Chen *et al.* (2010) demonstrated that blue light showed a trend of more intensity accompanied with a larger *m* value which is the empirical constant in kinetic model of product formation. This meant that blue light was capable of inducing phycocyanin production at a higher light intensity as compared to lower light intensity.

PHB accumulation is induced by optimal nutrients and light intensities as well (Ansari and Fatma, 2016; Costa *et al.*, 2018; Monshupanee and Incharoensakdi, 2014). Indeed, PHB production in cyanobacteria was induced when the cells were first cultivated in a nutrient-rich medium, followed by a second step, where the medium was deprived of a nitrogen source and/or supplemented with an organic carbon source to stimulate PHB accumulation (Charunchaipat *et al.*, 2020; Duangsri *et al.*, 2020; Panda and Mallick, 2007). A similar approach was used to assess the capacity of


PHB production by *A. platensis*. No PHB was detectable in cells grown in a nutrient-rich medium, while a substantial level of PHB production was apparent under nitrogen deprivation. The highest PHB content was obtained under HL white LED ( $9.18 \pm 0.32\%$ (w/w) DW) which was significantly higher than HL red, green, and blue LEDs ( $p < 0.05$ ) (Figure 2). A previous study demonstrated that a moderately HL white intensity ( $200 \mu\text{mol photons m}^{-2}\text{s}^{-1}$ ) was optimal for the co-production of glycogen, lipid and PHB in *Synechocystis* sp. PCC6803 (Monshupanee and Incharoensakdi, 2014). Besides, Abiusi *et al.*, (2014) reported that *Tetraselmis suecica* F&M-M33 grown in red LED attained the highest protein content, while cultures grown under blue and green lights showed the highest lipid and carbohydrate content, respectively. It has been shown that the total lipid content of three microalgal species, namely *Isochrysis galbana* LB987, *Nannochloropsis oculata* CCAP849/1, and *Dunaliella salina*, increases with increasing light intensity ( $80\text{--}100 \mu\text{mol photons m}^{-2}\text{s}^{-1}$ ). This is because excessive light energy was converted into lipid (Gim *et al.*, 2016). However, a shift from low to high irradiance is known to cause an immediate increase in photosynthetic activity and growth rate (Warner and Madden, 2007), as well as induce an initial



**Figure 2** Effects of light colour and light intensity on PHB content of *A. platensis*. Cells were cultivated under three light intensities designated as LL, ML, and HL for 5 days. Bar graphs represent mean values ( $\pm$  SD) of three independent experiments. Different letters (a, b, or c) indicate significant differences ( $p < 0.05$ ) between the groups obtained from cells cultured in LL, ML and HL.

accumulation of carbohydrates. High light irradiance ( $440 \mu\text{mol photons m}^{-2}\text{s}^{-1}$ ) activated the rapid accumulation of glycogen by up to 34% of cell dry weight after 9 hrs but did not generate PHB in *Spirulina maxima* (de Philippis *et al.*, 1992). C-Phycocyanin (C-PC) is known to expand the light absorption spectra and is thought to have high antioxidant potential that can help sustain *A. platensis* growth under such extreme conditions (Mogany *et al.*, 2020; Roy and Pabbi, 2022). The antioxidant potential of C-PC was determined by measuring DPPH (2,2-diphenyl picrylhydrazyl) radical scavenging activity. The reduction in the optical density of DPPH after interaction with the antioxidant was estimated. The C-PC con-

centrations from  $10$  to  $100 \mu\text{g mL}^{-1}$  extracted from cells cultured under HL blue light were incubated with DPPH solution for 30 min. Maximum inhibition was found at  $100 \mu\text{g mL}^{-1}$  of C-PC concentration with antioxidant activity of  $52.1 \pm 5.6\%$  for *A. platensis* (Figure 3). The DPPH radical scavenging activity of ascorbic acid at  $100 \mu\text{g mL}^{-1}$  concentration was  $80.2 \pm 4.8\%$ . Comparable antioxidant activity of the C-PC concentration was also detected in *Geitlerinema* sp. TRV57 and *Geitlerinema* sp. H8DM (Patel *et al.*, 2018; Renugadevi *et al.*, 2018). The reaction capacity of antioxidants depends on their structural characteristics more specifically the hydroxyl groups available for reduction. Moreover, Pan-utai and lamtham (2019) showed similar results to our study in terms of C-PC antioxidant activity having equal crude concentration in the range of 30–60% for DPPH



**Figure 3** *In vitro* antioxidant activity of partial purified C-PC of *A. platensis*. Ascorbic acid was used as a reference antioxidant. Error bar represent  $\pm$  SD of three independent experiments.

scavenging under normal cultivation condition.

The commercialization of many natural products requires a large amount of biomass and a high percentage of the desired valuable pigments in the biomass. In this respect, the growth of *A. platensis* under LL and ML red LEDs fulfils the requirement for a higher concentration of chlorophyll-*a* and C-PC in the biomass. Although lower biomass was observed in cells cultured under HL blue LED, a high amount of C-PC was obtained compared to that continuously grown under red LED. Moreover, the use of red LEDs could result in a more economically viable production process if we consider that leftover biomass after the extraction of C-PC still contains valuable products such chlorophyll-*a* and valuable proteins.

## Conclusion

This study concluded that light-emitting diode (LEDs) radiation enhanced biomass, chlorophyll-*a*, C-phycocyanin, and PHB production in *Arthrospira platensis*. Red LED supported biomass and chlorophyll-*a*, while blue LED enhanced phycocyanin. Besides, white LED increased PHB content. The results of this study offer a promising option for the utilization of alkaliphilic cyanobacteria in the production of C-PC and PHB.

## Acknowledgments

This research was financially supported by a graduate study development scholarship from the National Research Council of Thailand as of the 2020 fiscal year to C. Duangsri and W. Raksajit, by the Capacity Building of KU Students on Internationalization Program (KUCSI 2020) to C. Duangsri, by the Graduate School Fellowship Program in Agriculture and Agroindustry from the Agricultural Research Development Agency (Public Organization) as the fiscal year 2020 to W. Charunchaipipat and W. Raksajit.

## References

- Abiusi, F., Sampietro, G., Marturano, G., Biondi, N., Rodolfi, L., D'Ottavio, M., and Tredici, M. R. (2014). Growth, photosynthetic efficiency, and biochemical composition of *Tetraselmis suecica* F&M-M33 grown with LEDs of different colors. **Bio-technology and Bioengineering** 111(5): 956–964.
- Ansari, S., and Fatma, T. (2016). Cyanobacterial polyhydroxybutyrate (PHB): Screening, optimization and characterization. **PLoS ONE** 11(6): e0158168.
- Bennett, A., and Bogorad, L. (1973). Complementary chromatic adaptation in a filamentous blue-green alga. **Journal of Cell Biology** 58(2): 419–435.
- Campos Assumpção de Amarante, M., Cava-

- Icante Braga, A.R., Sala, L., and Juliano Kalil, S. (2020). Colour stability and anti-oxidant activity of C–phycocyanin–added ice creams after *in vitro* digestion. **Food Research International** 137: 109602.
- Charunchaipipat, W., Mudtham, N.–A., Promariya, A., Duangsri, C., Satchasataporn, K., Ngamkala, S., and Raksajit, W. (2020). Effect of butyrate supplementation on the production of a high value C–phycocyanin and polyhydroxybutyrate in *Spirulina platensis*. **Journal of Research Unit on Science, Technology and Environment for Learning** 11(2): 369–379. (in Thai)
- Chen, H.–B., Wu, J.–Y., Wang, C.–F., Fu, C.–C., Shieh, C.–J., Chen, C.–I., Wang, C.–Y., and Liu, Y.–C. (2010). Modeling on chlorophyll *a* and phycocyanin production by *Spirulina platensis* under various light–emitting diodes. **Biochemical Engineering Journal** 53(1): 52–56.
- Costa, S. S., Miranda, A. L., Andrade, B. B., Assis, D. D. J., Souza, C. O., de Moraes, M. G., Costa, J. A. V., and Druzian, J. I. (2018). Influence of nitrogen on growth, biomass composition, production, and properties of polyhydroxyalkanoates (PHAs) by microalgae. **International Journal of Biological Macromolecules** 116: 552–562.
- da Fontoura Prates, D., Duarte, J. H., Vendruscolo, R. G., Wagner, R., Ballus, C. A., da Silva Oliveira, W., Godoy, H. T., Barcia, M. T., de Moraes, M. G., Radmann, E. M., and Costa, J. A. V. (2020). Role of light emitting diode (LED) wavelengths on increase of protein productivity and free amino acid profile of *Spirulina* sp. Cultures. **Bioresource Technology** 306: 123184.
- de Philippis, R., Sili, C., and Vincenzini, M. (1992). Glycogen and poly– $\beta$ –hydroxybutyrate synthesis in *Spirulina maxima*. **Journal of General Microbiology** 138(8): 1623–1628.
- Duangstri, C., Mudtham, N.–A., Incharoensakdi, A., and Raksajit, W. (2020). Enhanced polyhydroxybutyrate (PHB) accumulation in heterotrophically grown *Arthrospira platensis* under nitrogen deprivation. **Journal of Applied Phycology** 32(6): 3645–3654.
- García, A. B., Longo, E., and Bermejo, R. (2021). The application of a phycocyanin extract obtained from *Arthrospira platensis* as a blue natural colorant in beverages. **Journal of Applied Phycology** 33(5): 3059–3070.
- Gim, G. H., Ryu, J., Kim, M. J., Kim, P. I., and Kim, S. W. (2016). Effects of carbon source and light intensity on the growth and total lipid production of three microalgae under different culture conditions. **Journal of Industrial Microbiology and Biotechnology** 43(5): 605–616.
- Khattoon, H., Kok Leong, L., Abdu Rahman,

- N., Mian, S., Begum, H., Banerjee, S., and Endut, A. (2018). Effects of different light source and media on growth and production of phycobiliprotein from freshwater cyanobacteria. **Bioresource Technology** 249: 652–658.
- Klepacz–Smółka, A., Pietrzyk, D., Szelaǵ, R., Głuszcz, P., Daroch, M., Tang, J., and Ledakowicz, S. (2020). Effect of light colour and photoperiod on biomass growth and phycocyanin production by *Synechococcus* PCC 6715. **Bioresource Technology** 313: 123700.
- Kumar, D., Wattal Dhar, D., and Pabbi, S. (2014). Extraction and purification of C–phycocyanin from *Spirulina platensis* (CCC 540). **Indian Journal of Plant Physiology** 19(2): 184–188.
- Luimstra, V. M., Schuurmans, J. M., Verschuur, A. M., Hellingwerf, K. J., Huisman, J., and Matthijs, H. C. P. (2018). Blue light reduces photosynthetic efficiency of cyanobacteria through an imbalance between photosystems I and II. **Photosynthesis Research** 138(2): 177–189.
- Lupatini, A. L., Colla, L. M., Canan, C., Colla, E. (2017). Potential application of microalga *Spirulina platensis* as a protein source. **Journal of the Science of Food and Agriculture** 97(3): 724–732.
- MacKinney, G. (1941) Absorption of light by chlorophyll solutions. **Journal of Biochemical Chemistry** 140: 315–322.
- Madhyastha, H. K., and Vatsala, T. M. (2007) Pigment production in *Spirulina fussiformis* in different photophysical conditions. **Biomolecular Engineering** 24(3): 301–305.
- Maltsev, Y., Maltseva, K., Kulikovskiy, M., Maltseva, S. (2021). Influence of light conditions on microalgae growth and content of lipids, carotenoids, and fatty acid composition. **Biology** 10(10): 1060.
- Markou, G. (2014). Effect of various colors of light–emitting diodes (LEDs) on the biomass composition of *Arthrospira platensis* cultivated in semi-continuous mode. **Applied Biochemistry and Biotechnology** 172(5): 2758–2768.
- McGee, D., Archer, L., Fleming, G. T. A., Gillespie, E., and Touzet, N. (2020). Influence of spectral intensity and quality of LED lighting on photoacclimation, carbon allocation and high–value pigments in microalgae. **Photosynthesis Research** 143(1): 67–80.
- Milia, M., Corrias, F., Addis, P., Zitelli, G. C., Cicchi, B., Torzillo, G., Andreotti, V., and Angioni, A. (2022). Influence of different light sources on the biochemical composition of *Arthrospira* spp. grown in model systems. **Foods** 11(3): 399.
- Minic, S., Radomirovic, M., Savkovic, N., Radibratovic, M., Mihailovic, J., Vasovic, T.,

- Nikolic, M., Milcic, M., Stanic–Vucinic, D., and Cirkovic Velickovic, T. (2018). Covalent binding of food–derived blue pigment phycocyanobilin to bovine  $\beta$ –lactoglobulin under physiological conditions. **Food Chemistry** 269: 43–52.
- Mogany, T., Kumari, S., Swalaha, F.M., and Bux, F. (2020). An in silico structural and physiochemical analysis of C–phycocyanin of halophile *Eubalotheca* sp. **Algal Research** 51:102025.
- Monshupanee, T., and Incharoensakdi, A. (2014). Enhanced accumulation of glycogen, lipids and polyhydroxybutyrate under optimal nutrients and light intensities in the cyanobacterium *Synechocystis* sp. PCC 6803. **Journal of Applied Microbiology** 116(4): 830–838.
- Moradi, Z., Madadkar Haghjou, M., Zarei, M., Colville, L., and Raza, A. (2021). Synergy of production of value–added bioplastic, astaxanthin and phycobilin co–products and direct green 6 textile dye remediation in *Spirulina platensis*. **Chemosphere** 280: 130920.
- Nematollahi, M. A., Laird, D. W., Hughes, L. J., Raeisossadati, M., Moheimani, N. R. (2020). Effect of organic carbon source and nutrient depletion on the simultaneous production of a high value bioplastic and a specialty pigment by *Arthrospira platensis*. **Algal Research** 47: 101844.
- Panda, B., and Mallick, N. (2007). Enhanced poly– $\beta$ –hydroxybutyrate accumulation in a unicellular cyanobacterium, *Synechocystis* sp. PCC 6803. **Letters in Applied Microbiology** 44(2): 194–198.
- Pan–utai, W., and lamtham, S. (2019). Extraction, purification and antioxidant activity of phycobiliprotein from *Arthrospira platensis*. **Process Biochemistry** 82: 189–198.
- Patel, A., Mishra, S., Pawar, R., and Ghosh, P. K. (2005). Purification and characterization of C–phycocyanin from cyanobacterial species of marine and freshwater habitat. **Protein Expression and Purification** 40(2): 248–255.
- Patel, H. M., Rastogi, R. P., Trivedi, U., and Madamwar, D. (2018). Structural characterization and antioxidant potential of phycocyanin from the cyanobacterium *Geitlerinema* sp. H8DM. **Algal Research** 32: 372–383.
- Prasanna, R., Sood, A., Jaiswal, P., Nayak, S., Gupta, V., Chaudhary, V., Joshi, M., and Natarajan, C. (2010). Rediscovering cyanobacteria as valuable sources of bioactive compounds (Review). **Applied Biochemistry and Microbiology** 46(2): 119–134.
- Renugadevi, K., Valli Nachiyar, C., Sowmiya, P., and Sunkar, S. (2018). Antioxidant activity of phycocyanin pigment extracted

- from marine filamentous cyanobacteria *Geitlerinema* sp. TRV57. **Biocatalysis and Agricultural Biotechnology** 16: 237–242.
- Roy, D., and Pabbi, S. (2022). A simple improved protocol for purification of C–phycocyanin from overproducing cyanobacteria and its characterization. **Journal of Applied Phycology**. DOI: 10.1007/s10811-021-02649-z.
- Sonani, R. R., Patel, S., Bhastana, B., Jakhar, K., Chaubey, M. G., Singh, N. K., and Madamwar, D. (2017). Purification and antioxidant activity of phycocyanin from *Synechococcus* sp. R42DM isolated from industrially polluted site. **Bioresource Technology** 245: 325-331.
- van Hieu, H., Quang–Tuong, L., and Cong Doan, B. (2021). The production of high phycocyanin by applications of red light–emitting diodes (LEDs) *in vitro* algae growth on *Spirulina platensis*. **Journal of Nano–and Electronic Physics** 13(3): 1–4.
- Wang, C.–Y., Fu, C.–C., and Liu, Y. –C. (2007). Effects of using light–emitting diodes on the cultivation of *Spirulina platensis*. **Biochemical Engineering Journal** 37(1): 21–25.
- Warner, M. E., and Madden, M. L. (2007). The impact of shifts to elevated irradiance on the growth and photochemical activity of the harmful algae *Chattonella subsalsa* and *Prorocentrum minimum* from Delaware. **Harmful Algae** 6(3): 332–342.
- Wicaksono, H. A., Satyantini, W. H., and Masithah, E.D. (2019). The spectrum of light and nutrients required to increase the production of phycocyanin *Spirulina platensis*. **IOP Conference Series: Earth and Environmental Science** 236(1): 012008.