# **TABLE OF CONTENTS**

i

TABLE OF CONTENTS	i
LIST OF TABLES	ii
LIST OF FIGURES	iv
LIST OF SCHEMES	ix
INTRODUCTION	1
OBJECTIVE S	4
LITERATURE REVIEWS	5
MATERIALS AND METHODS	38
RESULTS AND DISCUSSION	65
CONCLUSION	193
LITERATURE CITED	195
APPENDIX	214
CIRRICULUM VITAE	218

## LIST OF TABLES

Table		Page
1	Teleomorph and anamorph connection of Talaromyces species	15
2	Morphological characteristics of different species of Talaromyces	
	in four sections and three series	20
3	Forty-five soil samples were collected from various locations in	46
	Thailand	
4	Taxa and selected isolates used for studying molecular	
	phylogeny	49
5	List of primer sequences of 10 microsatellite loci	50
6	Isolates of Talaromyces flavus, T. bacillsporus and Talaromyces	
	sp. 1 (KUFC 3399) from various locations, used for antagonistic test	
	against plant pathogenic fungi	53
7	Fifteen species of plant pathogenic fungi from various diseased	
	plants used for testing of antagonistic activity of Talaromyces spp.	54
8	Frequency and number of isolates of Talaromyces species from	
	various soil samples at different locations	67
9	Distribution of Talaromyces spp. from various locations using	
	different isolation methods	68
10	Percent inhibition on mycelial growth of four oomycetous plant	
	pathogenic fungi by twenty isolates of Talaromyces flavus	
	cultivated on PDA as dual culture at 28 $^{\circ}$ C for 14 days	166
11	Percent inhibition on mycelial growth of five coelomycetous plant	
	pathogenic fungi by twenty isolates of Talaromyces flavus, each	
	isolate of <i>T. bacillisporus</i> and <i>Talaromyces</i> sp.1 (KUFC 3399)	
	cultivated on PDA as dual culture test at 28 $^{\circ}$ C for 14 days	167
12	Percent inhibition on mycelial growth of four hyphomycetous plant	
	pathogenic fungi by twenty isolates of Talaromyces flavus, each	
	isolate of T. bacillisporus and Talaromyces sp.1 (KUFC 3399)	
	cultivated on PDA, as dual culture at 28 $^{\circ}$ C for 14 days	168

## LIST OF TABLES (Continued)

	Page
Percent survival of mungbeen seedlings, after seeds were treated	
with T. flavus and inoculated with Sclerotium rolfsii at 7 and 14	
days after inoculation	172
<sup>1</sup> H NMR of bacillisporins A ( <b>4</b> ), B ( <b>5</b> ), C ( <b>6</b> ), D ( <b>69</b> ), duclauxin	
(50) (DMSO, 500 MHz), and bacillisporin E (70) (CDCb, 300	
MHz)	178
<sup>13</sup> C NMR of bacillisporins A (4), B (5), C (6), D (69), duclauxin	
(50) and bacillisporin E $(70)$ (125.77MHz)	178
NMR Data of compound <b>71</b> and <b>72</b> in CDCl <sub>3</sub> ( $^{1}$ H 300 MHz, $^{13}$ C	
75 MHz)	191
<sup>1</sup> H NMR of vermistatin (75), hydroxydihydrovermistatin (76) and	
penisimplissin (74) (300 MHz, CDCb)	192
$^{13}$ C NMR of vermistatin (75), hydroxydihydrovermistatin (76) and	
penisimplissin (74) (75.47 MHz, CDCs)	192
	Percent survival of mungbeen seedlings, after seeds were treated with <i>T. flavus</i> and inoculated with <i>Sclerotium rolfsii</i> at 7 and 14 days after inoculation <sup>1</sup> H NMR of bacillisporins A (4), B (5), C (6), D (69), duclauxin (50) (DMSO, 500 MHz), and bacillisporin E (70) (CDC 300 MHz) <sup>13</sup> C NMR of bacillisporins A (4), B (5), C (6), D (69), duclauxin (50) and bacillisporin E (70) (125.77MHz) NMR Data of compound 71 and 72 in CDC ( <sup>1</sup> H 300 MHz, <sup>13</sup> C 75 MHz) <sup>1</sup> H NMR of vermistatin (75), hydroxydihydrovermistatin (76) and penisimplissin (74) (300 MHz, CDC <sup>13</sup> C NMR of vermistatin (75), hydroxydihydrovermistatin (76) and penisimplissin (74) (75.47 MHz, CDC

## LIST OF FIGURES

Figures		Page
1	Morphology characteristics of four Talaromyces species	11
2	Morphology characteristics of three Talaromyces species	12
3	A synoptic illustrations of the ascospores ornamentation of	
	Talaromyces species	15
4	Map of Thailand indicating the collection sites of the soil samples	
	from 38 provinces	45
5	Occurrence of <i>Talaromyces</i> species from various soil samples at	
	different locations	65
6	Talaromyces austrocalifornicus KUFC 3401. Obverse and reverse	
	views of colonies on different media, incubated for 7 days at	
	25°C	75
7	Talaromyces austrocalifornicus KUFC 3401. Obverse and reverse	
	views of colonies on different media, incubated for 7 days at	
	28°C	76
8	Talaromyces austrocalifornicus KUFC 3401. Obverse and reverse	
	views of colonies on different media, incubated for 14 days at	
	28°C	77
9	Photomicrographs of Talaromyces austrocalifornicus KUFC 3401	78
10	Camera lucida drawings of Talaromyces austrocalifornicus KUFC	
	3401	89
11	Talaromyces bacillisporus KUFC 3350. Obverse and reverse	
	views of colonies on different media, incubated for 7 days at	
	25°C	83
12	Photomicrographs of Talaromyces bacillisporus KUFC 3350	84
13	Camera lucida drawings of Talaromyces bacillisporus KUFC 3350	85
14	Talaromyces flavus KUFC 3530. Obverse and reverse views of	
	colonies on different media, incubated for 7 days at 25°C	90

iv

#### Page Figures 15 Talaromyces flaws KUFC 3530. Obverse and reverse views of colonies on different media, incubated for 7 days at 28°C 91 16 Variations of colony pattern of *T. flavus* on MEA at 28°C for 93 14 days 17 Variations of penicilli of Talaromyces flavus 94 18 Camera lucida drawings of *Talaromyces flavus* showing 95 penicilli, conidia, ascomatal initials, asci and ascospores 19 SEM photomicrographs showing variations in shape, size and 95 ornamentation of ascospores among different isolates of T. favus 20 *Talaromyces helicus* var. *major* KUFC 3598. Obverse and reverse views of colonies on different media, incubated for 7 days at $25^{\circ}C$ 98 21 99 Photomicrographs of *Talaromyces helicus* var. *major* KUFC 3593 22 Camera lucida drawings of *Talaromyces helicus* var. major 100 23 Talaromyces luteus KUFC 3364. Obverse and reverse views of colonies on different media, incubated for 7 days at 28°C 103 24 Photomicrographs of *Talaromyces luteus* KUFC 3364 104 25 Camera lucida drawings of *Talaromyces luteus* KUFC 3364 105 26 *Talaromyces macrosporus* KUFC 3381. Obverse and reverse views of colonies on different media, incubated for 7 days at $25^{\circ}C$ 109 27 Talaromyces macrosporus KUFC 3363. Obverse and reverse views of colonies on different media, incubated for 7 days at $28^{\circ}C$ 110 28 Variations of colony pattern showing obverse and reverse views of various isolates of *T. macosporus* on MEA at 28°C, 14 days 111

v

Figures		Page
29	Photomicrographs of Talaro myces macrosporus KUFC 3363	112
30	Camera lucida drawings of Talaromyces macrosporus KUFC	
	3363	113
31	Talaromyces indigoticus KUFC 3366. Obverse and reverse views	
	of colonies on different media, incubated for 7 days at $25^{\circ}$ C	116
32	Talaromyces indigoticus KUFC 3366. Obverse and reverse views	
	of colonies on different media, incubated for 7 days at 28°C	117
33	Talaromyces indigoticus KUFC 3366. Obverse and reverse views	
	of colonies on different media, incubated for 14 days at 37°C	118
34	Photomicrographs of Talaromyces indigoticus KUFC 3366	119
35	Camera lucida drawings of Talaromyces indigoticus KUFC 3366	120
36	Talaromyces rotundus KUFC 3359. Obverse and reverse views of	
	colonies on different media, incubated for 7 days at $25^{\circ}C$	123
37	Photomicrographs of Talaromyces rotundus KUFC 3359	124
38	Camera lucida drawings of Talaromyces rotundus	125
39	Talaromyces stipitatus KUFC 3357. Obverse and reverse views of	
	colonies on different media, incubated for 7 days at $25^{\circ}C$	129
40	Talaromyces stipitatus KUFC 3357. Obverse and reverse views of	
	colonies on different media, incubated for 7 and 14 days at $28^{\circ}C$	130
41	Photomicrographs of Talaromyces stipitatus KUFC 3357	131
42	Camera lucida drawings of Talaromyces stipitatus	132
43	Camera lucida drawings of Talaromyces stipitatus KUFC 3357	133
44	Talaromyces trachyspermus KUFC 3355. Obverse and reverse	
	views of colonies on different media, incubated for 7 days at	
	25°C	136
45	Photomicrographs of Talaromyces trachyspermus KUFC 3355	137
46	Camera lucida drawings of Talaromyces trachyspermus KUFC	
	3355	138

vi

#### Figures Page 47 Talaromyces wortmannii KUFC 3333. Obverse and reverse views 142 of colonies on different media, incubated for 7 days at 25°C 48 Talaromyces wortmannii KUFC 3333. Obverse and reverse views 143 of colonies on different media, incubated for 7 days at 28°C 49 Talaromyces wortmannii KUFC 3354. Obverse and reverse views of colonies on different media, incubated for 7 days at 25°C 143 50 Photomicrographs of *Talaromyces wortmannii* KUFC 3333 144 51 Camera lucida drawings of Talaromyces wortmannii KUFC 3333 145 52 Camera lucida drawings of *Talaromyces wortmannii* KUFC 3354 146 53 Camera lucida drawings of *Talaromyces wortmannii* 147 54 Talaromyces sp. 1 KUFC 3399. Obverse and reverse views of colonies on different media, incubated for 7 days at 25°C 151 55 Talaromyces sp. 1 KUFC 3399. Obverse and reverse views of colonies on different media, incubated for 7 days at 28°C 152 56 Photomicrographs of *Talaromyces* sp. 1 KUFC 3399 153 57 Camera lucida drawings of *Talaromyces* sp. 1 KUFC 3399 154 58 Talaromyces sp. 2 KUFC 3383. Obverse and reverse views of colonies on different media, incubated for 7 days at 25°C 157 59 Photomicrographs of *Talaromyces* sp. 2 KUFC 3383 158 50 159 Camera lucida drawings of *Talaromyces* sp. 2 KUFC 3383 61 Electrophoresis of PCR products using PM7 161 62 Phylogenetic relationships of *Talaromyces* species and related Trichocomaceae based on SSR gene 162 63 Antagonistic tests as dual cultures of different *Talaromyces flavus* isolates (left) and plant pathogenic fungi (right) on PDA 169 incubated for 14 days at 28°C

vii

Figures		Page
64	Antagonistic tests as dual cultures of different Talaromyces flavus	
	isolates (left) and plant pathogenic fungi (right) on PDA	
	incubated for 14 days at 28°C	170
65	A. Mungbean seeds + distilled water (left) and mungbean seeds +	
	Sclerotium rolfsii (right), 7 days after planting (Control), B. mungbean	
	seeds + S. rolfsii (a), mungbean seeds + Talaromyces flavus KUFC	
	3523 + S. rolfsii (b), C. mungbean seeds $+ T.$ flavus KUFC $3550 + S.$	
	roflsii (c), D. mungbean seeds + T. flavus KUFC 3528 + S. rolfsii (d), 7	
	days after planting	173
66	Mungbean seeds inoculated with Sclerotium rolfsii for 7 days;	
	the collapsed mungbean seedlings and sclerotia (circle d) are	
	shown	173
67	Structure of compounds isolated from the culture of	
	T. bacillisporus, collected from Kasetsat University, Bangkok	177
68	ORTEP view of thailandolide A (71)	190
69	ORTEP view of vermistatin (75)	190

viii

# LIST OF SCHEMES

## Schemes

## Page

1	Formation of the upper oxaphenalenone monomer	180
2	Formation of the lower oxaphenalenone monomer	181
3	Formation of the free radical of the upper oxaphenalenone	
	monomer	182
4	Formation of the free radical $(U)$ of the second oxaphenalenone	
	monomer ( <b>R</b> )	182
5	Oxidative coupling of two oxaphenalenone units	183
6	Formation of bacillosporin C	183
7	Formation of bacillosporin A and bacillosporin B	184
8	Formation of duclauxin from bacillosporin A	185