


Cite this article: Onsrissawat, P., Rodthong, S., Yahuafai, J., & Urairong, H. (2022, September).  
Hepatoprotective and antioxidant properties of *Schizophyllum commune* fruiting body. *Journal of Current  
Science and Technology*, 12(3), 615-628. DOI:



## Hepatoprotective and antioxidant properties of *Schizophyllum commune* fruiting body

Prayumat Onsrissawat<sup>1</sup>, Sureelak Rodthong<sup>3</sup>, Jantana Yahuafai<sup>2</sup>, and Hathairat Urairong<sup>1\*</sup>

<sup>1</sup>College of Agricultural Innovation and Food Technology, Rangsit University,  
Pathum Thani 12000, Thailand

<sup>2</sup>Clinical Research Section, Division of Research and Academic Support, National Cancer Institute,  
Bangkok 10400, Thailand

<sup>3</sup>School of Microbiology, Institute of Science, Suranaree University of Technology,  
Nakhon Ratchasima 30000, Thailand

\*Corresponding author; E-mail: fongpatt@yahoo.com

Received 30 April 2022; Revised 22 July 2022; Accepted 24 July 2022;  
Published online 26 December 2022

### Abstract

The split gill fungus, *Schizophyllum commune*, is an edible mushroom acknowledged for its medical importance and hepatoprotective activity. This study assessed the hepatoprotective and antioxidant activities of extraction from *S. commune*. The dried fruiting body of *S. commune* was extracted with different solvents (hexane, chloroform, ethyl acetate, and methanol) and by the hot water extraction method, followed by ethanol precipitation to provide crude polysaccharide. The extracts were evaluated for their scavenging activity, total phenolic content, and flavonoid content, with cytotoxicity determined in human hepatoma cells (HepG2). The hepatoprotective effect of *S. commune* extracts was evaluated using hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>)-induced toxicity on HepG2 cells, while protective effects and morphological changes were assessed using Hoechst 33342 staining assay. Lipid peroxidation and reduced glutathione levels induced by H<sub>2</sub>O<sub>2</sub> were studied to assess possible mechanisms of hepatoprotection by *S. commune* extracts. The five extracts showed DPPH free radical scavenging activities with IC<sub>50</sub> values of 0.57±0.02 mg/mL, 0.40±0.01 mg/mL, 0.36±0.00 mg/mL, 0.32±0.00 mg/mL and 1.00 ±0.05 mg/mL. Antioxidant activity IC<sub>50</sub> values of DPPH scavenging correlated with total phenolic and total flavonoid contents. Treatment of HepG2 cells with polysaccharide extract before H<sub>2</sub>O<sub>2</sub> exposure increased cell viability, while polysaccharide extract inhibited lipid peroxidation and increased glutathione levels against H<sub>2</sub>O<sub>2</sub>-induced damage. Results revealed that the polysaccharide extract from the fruiting body of *S. commune* had hepatoprotective potential and showed promise as a potential therapeutic agent for liver diseases.

**Keywords:** antioxidant; hydrogen peroxide; hepatoprotective; polysaccharide extract; *Schizophyllum commune*.

### 1. Introduction

Oxidative stress is an important contributor to the pathophysiology of various pathological conditions including cardiovascular dysfunctions, atherosclerosis, inflammation, carcinogenesis, reperfusion injury and neurodegenerative diseases (Aruoma, Grootveld, & Bahorun, 2006). The liver is the most significant and important organ in the

vertebrate body, involving the metabolism of food molecules, detoxification and biotransformation. (Kumar, Kumar, & Sinha, 2019). It is also the most important site of intermediary metabolism by processing both endogenous (e.g., cholesterol, fatty acids and proteins) and exogenous substances (e.g., drugs) and converting them into less-harmful chemicals by detoxification or elimination. Prolonged

exposure to exogenous compounds such as drugs, alcohol or chemicals causes liver dysfunction due to increased reactive oxygen species (ROS) in hepatic cells, leading to oxidative stress (Lister et al., 2020; Yoon, Baber, Chondhary, Kuter, & Pysopoulos, 2016). Many bioactive compounds have protective effects against oxidative stress-induced cell death (Ali et al., 2004; Jaydeokar, Bandawane, Bibave, & Patil, 2014). Mushrooms possess naturally occurring antioxidants as phenolic and polysaccharide compounds that can prevent and cure liver damage (Mau, Tsai, Tseng, & Huang, 2005; Selima, Aminul, Ugur, & Narayan, 2012). Several mushroom species including *Ganoderma lucidum* (Kao, Jesuthasan, Bishop, Glucina, & Ferguson, 2013), *Grifola frondosa* (Masuda, Ito, Konishi, & Nonbu, 2010), *Lentinula edodes* (Wu, & Hansen, 2008) and *Schizophyllum commune* contain antioxidants.

*Schizophyllum commune* or split gill mushroom is commonly found in many countries including Laos, Myanmar, India and Thailand. *S. commune* has a protein content of 16-27%, (Aletor, 1995; Longvah, & Deosthale, 1998) low-fat content, fatty acids and amino acids essential to the human diet (Longvah, & Deosthale, 1998), together with a polysaccharide known as Schizophyllan, which has a chemical structure consisting of a  $\beta$  (1-3) glucan chain linked via  $\beta$  (1-6) glucan (Zhang, Kong, Fang, Nishinari, & Phillips, 2013). Multifunctional biological properties include anti-tumor, antimicrobial, immunomodulatory and anti-inflammatory functions (Lindequist, Niedermeyer, & Julich, 2005; Joel, & Bhimba, 2013). *S. commune* methanol and ethyl acetate extracts inhibited superoxide radicals (Mirfat, Noorlidah, & Vikineswary, 2010), while the ethyl acetate fraction exhibited high inhibition against lipid peroxidation in egg yolk (Mayakrishnan et al., 2013). However, no reports are available detailing the antioxidant and associated hepatoprotective activities of *S. commune* extracted by different organics-solvents and polysaccharides.

## 2. Objectives

This study investigated the hepatoprotective activity of different organic solvents and polysaccharide extract from the fruiting body of *S. commune* against liver damage mediated by hydrogen peroxide ( $\text{H}_2\text{O}_2$ ) in human hepatocellular liver carcinoma cells (HepG2 cells) to determine its antioxidant activity.

## 3. Materials and Methods

### 3.1 Materials and chemicals

*Schizophyllum commune* (Fr.) were cultured from a certified organic farm in Chantaburi Province, Thailand. The fruiting body was dried in a hot air oven at 50 °C for 4 days, then ground into powder and stored for further extraction. Hexane, chloroform, ethyl acetate, methanol, ethanol and DMSO were purchased from RCI Labscan Co., Ltd., Thailand. Folin-Ciocalteu phenol reagent was purchased from Sigma-Aldrich, Switzerland. Standard gallic acid was purchased from Sigma-Aldrich, Germany. Standard quercetin was purchased from Sigma-Aldrich, India. Standard BSA and Coomassie reagent were purchased from Thermo Scientific, USA. Standard D-glucose was purchased from VWR Life Science, USA. HepG2 (human hepatocellular liver carcinoma cell line, ATCC) was purchased from USA. Minimum Essential Medium (MEM) was purchased from Gibco, USA. Fetal bovine serum (FBS) was purchased from Gibco, UK. Hydrogen peroxide ( $\text{H}_2\text{O}_2$ ) was purchased from Polkie Odczynniki Chemiczne S. A. MTT (thiazolyl blue tetrazolium bromide) was purchased from Sigma-Aldrich, China. Hoechst 33342 was purchased from Invitrogen, USA. 2-Thiobarbituric acid (TBA) was purchased from Sigma-Aldrich, UK. Trichloroacetic acid (TCA) was purchased from Merck, Germany. Standard glutathione was purchased from Sigma-Aldrich, USA. DTNB (5, 5'-dithiobis (2-nitrobenzoic acid) was purchased from Fluka Analytical, USA. All reagents and chemicals were of analytical grade.

### 3.2 Preparation of extracts

#### *Solvent extraction*

The dried powdered fruiting body of *S. commune* (100 g) was extracted by maceration using hexane, ethyl acetate, chloroform and methanol. Crude extracts were obtained by evaporating the solvent in a rotary evaporator and then stored at -20 °C until use. Percentage yields of the crude extracts were calculated as [weight of crude mushroom extract (g) / weight of dry mushroom (g) x 100%] (Bugrahan et al., 2017).

### 3.3 Polysaccharide extraction

The dried powdered fruiting body (100 g) was extracted with distilled water at 85 °C and centrifuged. The supernatant was concentrated under reduced pressure at 40 °C and precipitated with six volumes of 95% ethanol (Zhong et al., 2015). The

solution was freeze-dried to provide crude polysaccharides and then stored at -20 °C until use. The percentage yield of the crude extract was calculated as above.

#### 3.4 Schizophyllan analyses

The polysaccharide extract was filtered through a 0.2 µm filter and 20 µL of the filtrate was loaded into the HPLC system. (Thermo, model: Spectra Series Complete HPLC AS3000, P4000, UV1000, SN4000, USA) Separation was performed on an Agilent Eclipse Plus C18 column (100 mm × 4.6 mm, 3.5 µm) using a 0.8 mL/min flow rate at 25 °C. The mobile phase consisted of eluent A (Water) and eluent B (Methanol). A gradient program was used for elution comprising 0–5 min, 90% A; 5–10 min, 85% A. Absorbance of the sample solution was measured at 205 nm (Su, Lai, Lin, & Ng, 2016).

#### 3.5 Measurement of carbohydrate and protein contents

Total carbohydrate content of the polysaccharide was determined by the phenol–sulfuric acid colorimetric method using D-glucose as the standard (Masuko et al., 2005). Protein in the polysaccharide was also estimated by the Bradford method (Bradford, 1976) using bovine serum albumin (BSA) as the standard.

#### 3.6 Total phenolic determination

Phenolic content was determined by the Folin–Ciocalteu method following Thomas, Priyadarshini, & Michael (2012). Briefly, 10 µL of each extract was mixed with 20% Folin-Ciocalteu phenol reagent (150 µL) in 96-well plates, followed by adding 50 µL of sodium carbonate (Na<sub>2</sub>CO<sub>3</sub>) after 3 min. The mixture was incubated for 2 h in the dark and absorbance was measured at 765 nm using a microplate reader (TECAN, Infinite M200 Pro, Switzerland). Total phenolic content was expressed as gallic acid equivalent (GAE) in mg per g of extract.

#### 3.7 Total flavonoid determination

Flavonoid content of each extract was determined using an aluminum chloride colorimetric assay. (Ghasemi, Ghasemi, & Ebrahimza, 2009). Briefly, 50 µL of each extract was mixed with 150 µL of methanol, 10 µL of 10% aluminum chloride (AlCl<sub>3</sub>) and 10 µL of 1M potassium acetate, followed by 30 µL of distilled

water. Samples were mixed and kept for 30 min in the dark. Absorbance was measured at 415 nm using a microplate reader. Total flavonoid content was expressed as quercetin equivalent (QE) in mg per g of extract from a calibration curve of quercetin standard solution.

#### 3.8 DPPH free radical scavenging activity

Scavenging ability was determined by the stable radical DPPH (2, 2-diphenyl picrylhydrazyl) as described by Gerhauser et al. (2013). Various 5 µL concentrations of different extracts (hexane, chloroform, ethyl acetate, methanol and polysaccharide) were mixed with 195 µL ethanolic solution containing DPPH (0.05 mg/mL) and left for 30 min in the dark. Absorbance was measured at 515 nm using a microplate reader (TECAN, Infinite M200 Pro, Switzerland). Ethanol was used as a blank. Sample concentration that scavenged 50% of DPPH free radical (IC<sub>50</sub> value) was calculated.

#### 3.9 Cell cultures

HepG2 cells (human hepatocellular liver carcinoma cell line) were obtained from ATCC (ATCC® HB-8065™). Cells were cultured in Minimum Essential Medium (MEM), supplemented with 1% penicillin-streptomycin and 10% FBS. Cells were grown in 5% CO<sub>2</sub> at 37 °C. HepG2 showing more than 80% cell confluence was used.

#### 3.10 Cell viability

The MTT assay was performed to assess cytotoxicity following Siripong, Rassamee, Piyaviriyakul, Yahuafai, & Kanokmedhakul (2012). In brief, HepG2 cells were seeded in 96-well plates at a density of 1 × 10<sup>4</sup> cells per well and allowed to grow overnight. Then, the cells were exposed to various concentrations of each extract of *S. commune* (100–1000 µg/mL) or H<sub>2</sub>O<sub>2</sub> (100–1000 µM) for 24 h. After exposure, 5 mg/mL MTT was added to each well and the plates were further incubated for 3 h. Then, the reaction mixture was carefully taken out, and 100 µL of DMSO was added to each well and mixed gently by pipetting up and down. Absorbance of the plates was measured at 550 nm.

#### 3.11 Hepatoprotective activity assay

The cytoprotective activity of test samples on H<sub>2</sub>O<sub>2</sub>-induced cell injury was investigated. Cell densities of 1 × 10<sup>4</sup> cells per well were plated in 96-

well plates with culture medium for 24 h, then exposed to test samples for 1 h and added with H<sub>2</sub>O<sub>2</sub> 100 µM for 24 h. Following treatment, the cells were added to a 5 mg/mL MTT solution in MEM, and further incubated for 3 h. The resulting formazan was dissolved in 100 µL of DMSO. The developed color was read at 550 nm using a microplate reader. Cell viability was determined as described above and expressed as a percentage with the vehicle-treated control cells as 100% (Siripong et al., 2012).

### 3.12 Morphological changes induced by Hoechst 33342 staining

Hoechst 33342 staining was performed to observe changes in nuclear morphology. Briefly, HepG2 cells were seeded at a density of  $1 \times 10^4$  cells/mL in 6-well plates and incubated for 24 h at 37 °C. Then, the cells were treated with polysaccharide extract at final concentrations of (400, 800 and 1000 µg/mL). After 1 h, H<sub>2</sub>O<sub>2</sub> (100 µM) was added and the cells were incubated for 24h. After treatment, the HepG2 cells were stained with 5 µg/mL Hoechst 33342 for 20 min at room temperature (Phonghanpot, & Jarintanan, 2021). Finally, the cells were imaged under a fluorescence microscope (Nikon, & Yokohama, Japan).

### 3.13 Lipid peroxidation (LPO)

HepG2 cells were plated at  $1 \times 10^5$  cells per well in 6-well plates and allowed to stabilize for 24 h before treatment with 1 mL of polysaccharide extract (400-1000 µg/mL). After 1 h, H<sub>2</sub>O<sub>2</sub> (100 µM) was added and the cells were incubated for 24 h. Cells were then gently washed twice with PBS and harvested in 1.5 mL of PBS by scraping. Following centrifugation, the pellet was re-suspended in 100 µL of 25 mM Tris-HCl buffer (pH 7.4) and subjected to ultrasonication for 5 min. The protein content of the cell suspension was measured using BSA as the standard. The extent of lipid peroxidation was estimated by measuring levels of malondialdehyde (MDA) using the thiobarbituric acid reactive substances (TBARS) assay. One milliliter of the reaction mixture was mixed with 2 mL of thiobarbituric acid (15%), trichloroacetic acid (10%) and hydrochloric acid (0.25 N) at a ratio of 1:1:1. The mixture was heated in a 95 °C water bath for 40 min and cooled at room temperature for 10 min. Following centrifugation, the absorbance of the supernatant was measured at 532 nm. Positive and negative controls consisted of cells treated with

H<sub>2</sub>O<sub>2</sub> and medium alone, respectively. A standard calibration curve was prepared from 1,1,3,3-tetramethoxypropan (0.3-10 µg/mL), a commercial form of malondialdehyde (MDA). The lipid peroxidation was calculated from the standard curve. The result was expressed as LPO level (%) Chourasiya and Sharma, S. (2021).

### 3.14 Glutathione level

The glutathione level in HepG2 cells was measured following the methods of Chandra et al. (2002) and Al-Sheddi et al. (2015). In brief, HepG2 cells were plated at  $1 \times 10^5$  cells per well in 6-well plates for 24h and then exposed to polysaccharide extract (400-1000 µg/ mL) and H<sub>2</sub>O<sub>2</sub> (100 µM) for 24 h. After respective exposures, cells were centrifuged and cellular protein was precipitated with 1 mL TCA (10%) on ice for 30 min. The supernatant was taken by centrifugation at 13000 rpm for 10 min. Then, 2 mL of 0.4 M Tris buffer (pH 8.9) containing 0.02 M EDTA and 0.01 M 5, 5'-pol-dithionitrobenzoic acid (DTNB) were added to the supernatant. The absorbance of the developed yellow color was read at 412 nm after incubating for 10 min at 37 °C. The total glutathione was determined from the standard curve of reduced glutathione(GSH).The result was expressed as GSH level (%)

### 3.15 Statistical analysis

Data, presented as the mean  $\pm$  SE, were representative of at least three independent experiments. Statistical analysis was performed using SPSS version 18. Differences among means were analyzed by one-way analysis followed by a Dunnett's post hoc test to compare values between the control and treatment groups. Graphs were drawn using GraphPad Prism software version 5.

## 4. Results and Discussion

### 4.1 Quantification of total bioactive compounds

Several *in vitro* and *in vivo* studies revealed that the presence of phenolic compounds, flavonoids and polysaccharide compounds in mushroom extracts was related to important biological properties such as antioxidant, immunomodulatory, antimicrobial and anticancer activity (Ooi, & Liu, 2000). Preventive effects on chronic diseases, in particular antiallergic, antibacterial, antifungal, anti- inflammatory, antioxidative, antiviral, cytotoxic,

immunomodulating, antidepressive, antihyperlipidemic, antidiabetic, digestive, hepatoprotective, neuroprotective, nephroprotective, osteoprotective and hypotensive activities were confirmed by many studies and clinical trials (Venturella, Ferraro, Cirlincione, & Gargano, 2021). Flavonoids, phenolic acids and polysaccharides are among the most important phytochemical constituents of *S. commune*. Here, we evaluated their contents in several *S. commune*

extracts. Results of the quantitative determination of total phenolics and flavonoids from different extracts and yields of crude extracts (mg extract/g) are presented in Table 1. Extracted yields of each solvent in descending order were methanol (6.99%), ethyl acetate (1.04%), chloroform (0.78%) and hexane (0.65%) demonstrating higher yields with increasing solvent polarity. The polysaccharide extract with hot water appeared as a white powder with a yield of 8.91%.

**Table 1** Total phenolic and flavonoid contents (mg/g) in *S. commune* extracts.

Extract	Yield %w/w(	Total phenolic content) mg GAE/g of extract( a	Total flavonoid content) mg QE/g of extract( b
Hexane	0.65	16.09±0.43	3.37±0.04
Chloroform	0.78	24.29±0.24	4.53±0.26
Ethyl acetate	1.04	25.93±0.80	3.98±0.10
Methanol	6.99	16.93±0.88	3.20±0.11
Polysaccharide	8.91	11.98±0.12	2.34±0.09


Data are mean±SE of three independent experiments

<sup>a</sup> mg GAE/g gallic acid equivalent per g of extract, <sup>b</sup> mg QE/g quercetin equivalent per g of extract

Total phenolic contents of *S. commune* extracts were determined. The highest values were observed in the ethyl acetate extract (25.93±0.80 mg GAE/g) followed by chloroform (24.29±0.24 mg GAE/g) methanol (16.93±0.88 mg GAE/g), and hexane (16.09±0.43mg GAE/g). Conversely, total flavonoid contents of *S. commune* extracts exhibited a slightly different descending order as chloroform (4.53±0.26 mg QE/g of extract), ethyl acetate (3.98±0.10 mg QE/g of extract), hexane (3.37±0.04 mg QE/g of extract) and methanol (3.20±0.11 mg QE/g of extract).

Chemical compositions of *S. commune* polysaccharide extracts were analyzed by HPLC. The phenol–sulfuric acid and Bradford methods

were used to evaluate total carbohydrate content and total protein content, respectively. Results showed that the total carbohydrate content and total protein content of the polysaccharide extract were 1.57±0.07 µg/mL and 0.29 ± 0.01 µg/mL, respectively. Schizophyllan (0.38 ± 0.04 mg/g extract) was the major active compound in the polysaccharide extract of *S. commune* fruiting body but absent in the organic solvent extracts. A chromatogram of polysaccharides using the HPLC method is presented in Figure1. A and B. Our results concurred with Mirfat et al. (2010) who determined dried fruiting body phenolic contents in methanol, ethyl acetate and dichloromethane from *S. commune*.



**Figure 1** (A) HPLC chromatograms of Schizophyllan standards, (B) HPLC chromatograms of polysaccharide extract: the first peak is Schizophyllan.

#### 4.2 DPPH radical scavenging activity of *S. commune* extracts

Free radical scavenging activities of *S. commune* extracts were tested using a methanol solution of the stable free radical DPPH. This decreases significantly when exposed to radical scavengers by providing a hydrogen atom or electron as a stable diamagnetic molecule. *S. commune* extract was evaluated using gallic acid as the standard control. Table 2 shows the DPPH radical scavenging activities of all extracts of *S.*

*commune*. Results revealed that the methanolic extract exhibited the highest radical scavenging activity with an  $IC_{50}$  value of  $0.32 \pm 0.00$  mg/mL, similar to the ethyl acetate extract at  $0.36 \pm 0.00$  mg/mL and followed by the chloroform extract at  $0.40 \pm 0.01$  mg/mL, hexane at  $0.57 \pm 0.02$  mg/mL and polysaccharide extract at  $1.00 \pm 0.05$  mg/mL. This data agreed with previous studies (Mirfat et al., 2010), confirming the methanol extract of *S. commune* as the most potent antioxidant agent.

**Table 2** DPPH free radical scavenging activities of *S. commune* extracts.

Extract	$IC_{50}$ (mg/mL)
Hexane	$0.57 \pm 0.02$
Chloroform	$0.40 \pm 0.01$
Ethyl acetate	$0.36 \pm 0.00$
Methanol	$0.32 \pm 0.00$
Polysaccharide	$1.00 \pm 0.05$
Gallic acid	$0.002 \pm 0.01$


Data are mean  $\pm$  SE of three independent experiments

#### 4.3 Effects of *S. commune* extracts against $H_2O_2$ -induced cytotoxicity

Herbal medicines are globally claimed to provide relief against many liver diseases but scientific evidence is lacking. Our results suggested valid hepatoprotective activity of *S. commune* extract in HepG2 cells. Cytotoxic effects of the extracts were assessed by the MTT assay. Results showed that the hexane and methanol extracts did not have any effect ( $p > 0.01$  and  $0.001$ ) on HepG2 cell viability (Figure 2A and 2D) at concentrations lower than  $200 \mu\text{g/mL}$ , while the chloroform and

ethyl acetate extracts (Figure 2B and 2C) had no effect ( $p > 0.001$ ) on HepG2 cell viability at concentrations lower than  $400 \mu\text{g/mL}$ .

As shown in Figures 2E and 2F, the polysaccharide extract at  $100$ - $1000 \mu\text{g/mL}$  and schizophyllan, as the main active compound in the polysaccharide extract, had no effect on cell viability. The non-cytotoxic extract of more than  $80\%$  was selected to study the hepatoprotective effect against  $H_2O_2$ -induced cytotoxicity in HepG2 cells.




**Figure 2** Cytotoxicity assessment by MTT assay in HepG2 cells ( $1 \times 10^4$  cells/well) following exposure of various concentrations of hexane extract (A), chloroform extract (B), ethyl acetate extract (C), methanol extract (D), polysaccharide extract (E) and standard Schizophyllan (F) for 24 h. Values are mean  $\pm$  SE of three independent experiments. \*\* $p < 0.01$ , \*\*\* $p < 0.001$  versus the control.

As shown in Figure 3,  $H_2O_2$  ranging from 100 to 1000  $\mu M$  induced cell death in a concentration-dependent manner and also demonstrated that  $H_2O_2$  at 100  $\mu M$  could be used to induce oxidative stress-mediated cytotoxicity in a cell system. To illustrate the cytoprotective effects of non-cytotoxic extracts (hexane, chloroform, ethyl acetate, methanol, polysaccharide and schizophyllan) on the viability of HepG2 cells against cytotoxicity induced by the  $H_2O_2$ , the HepG2 cells were incubated with various concentrations of test extracts for 24 h and then with a cytotoxic concentration (100  $\mu M$ ) of  $H_2O_2$ . The  $H_2O_2$ -induced cell death and hepatoprotective effect were determined by MTT and morphological changes. The protective potential of each extract, as observed in HepG2 cells, is presented in Figure 5. A reduction in cell viability was observed in HepG2 cells following exposure to  $H_2O_2$  (100  $\mu M$ ) for 24 h. Cell viability was observed in HepG2 cells pre-exposed to hexane, chloroform, ethyl acetate and methanol extracts at 12.5, 25, 50, 100 and 200  $\mu g/mL$ . Results showed that the hexane, ethyl acetate and methanol extracts did not have protective potential for viability of HepG2 cells exposed to 100  $\mu M$   $H_2O_2$  (Figure 4A, 4B and 4C), while methanol extracts at concentrations of 100 and 200  $\mu g/mL$  showed cell viability of  $56.25 \pm 0.41$


and  $59.55 \pm 0.67\%$ , respectively ( $p > 0.01$  and  $0.001$ ) (Figure 4D). Among all the organic solvents, only these two concentrations showed slightly effective protection on HepG2 cells treated with 100  $\mu M$   $H_2O_2$ . Interestingly, the polysaccharide extract showed highly significant ( $p < 0.001$ ) protection against  $H_2O_2$ -induced cytotoxicity, expressed as percentage of cell viability at concentrations of 200, 400, 800 and 1000  $\mu g/mL$  at  $63.29 \pm 0.97$ ,  $77.75 \pm 3.56$ ,  $86.65 \pm 4.42$  and  $98.86 \pm 2.97\%$ , respectively (Figure 4E). A similar effect of standard schizophyllan at concentrations of 400, 800 and 1000  $\mu g/mL$  exhibited viability of HepG2 cells at  $69.29 \pm 1.80$ ,  $85.48 \pm 2.03$  and  $98.23 \pm 3.22\%$ , respectively ( $p < 0.001$ ) (Figure 4F). However, the polysaccharide extract was effective in protecting the viability of HepG2 cells damaged by  $H_2O_2$  at a dose of 200  $\mu g/mL$ , lower than standard schizophyllan, possibly as a result of synergistic protective effects. Alterations in the morphology of HepG2 on exposure to polysaccharide extract and  $H_2O_2$  are shown in Figure 5(A–E). Cells exposed to 100  $\mu M$  of  $H_2O_2$  showed reduced normal morphology of HepG2 cell adhesion capacity compared with the control. Most of the cells exposed to  $H_2O_2$  lost their typical morphology, showed higher amounts of cell debris and appeared smaller in size (Figure 5B). HepG2 cells exposed to

increasing concentrations of polysaccharide extract for 24 h before H<sub>2</sub>O<sub>2</sub> exposure were restored to their original morphology in a concentration-dependent manner (Figure 5C–E). Results suggested that exposure of HepG2 cells to the polysaccharide extract significantly increased viability against H<sub>2</sub>O<sub>2</sub>-induced cell death but extracts from the organic solvents did not show cytoprotective properties. Our results support previous studies that polysaccharides from mushrooms exhibit significant hepatoprotective effects against H<sub>2</sub>O<sub>2</sub>. Several studies also demonstrated that various

compounds isolated from mushrooms, such as schizophyllan, showed prominent cytoprotective potential against cell damage (Kukan et al., 2004; Soares et al., 2013). Our study affirmed for the first time the hepatoprotective activity of the polysaccharide extract of *S. commune*, with the potential to protect against cell damage induced by H<sub>2</sub>O<sub>2</sub>. Therefore, we selected the polysaccharide extract to examine morphological changes, lipid peroxidation and glutathione levels in the following experiment.




**Figure 3** Cytotoxicity assessment by MTT assay in HepG2 cells ( $1 \times 10^4$  cells/well) following exposure of various concentrations of hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>) for 24 h. Values are mean  $\pm$  SE of three independent experiments. \*\*\* $p < 0.001$  versus the control.



**Figure 4** Assessment of protective potential of compounds on cell viability of HepG2 cells by MTT assay. Cells ( $1 \times 10^4$  cells/well) were exposed to various concentrations of non-cytotoxic extracts for 24 h, and then exposed to H<sub>2</sub>O<sub>2</sub> at 100 μM for 24 h. Values are mean  $\pm$  SE of three independent experiments. \*\* $p < 0.01$ , \*\*\* $p < 0.001$  versus H<sub>2</sub>O<sub>2</sub> exposure.






**Figure 5** Morphological changes in HepG2 cells observed by contrast inverted microscopy at 20x magnification. Control group (A), H<sub>2</sub>O<sub>2</sub> 100 μM (B), polysaccharide 400 μg/ mL + H<sub>2</sub>O<sub>2</sub> 100 μM (C), polysaccharide 800 μg/mL + H<sub>2</sub>O<sub>2</sub> 100 μM (D) and polysaccharide 1000 μg/mL + H<sub>2</sub>O<sub>2</sub> 100 μM (E).

#### 4.4 Morphological changes

Cells with homogeneous chromatin distribution were considered viable, whereas the presence of chromatin condensation and/or nuclear fragmentation was indicative of hallmarks of apoptosis. The morphological changes observed in HepG2 cells pre-exposed to polysaccharide extract and then H<sub>2</sub>O<sub>2</sub> for 24 h are summarized in Figure 6. As shown in Figure 6B, H<sub>2</sub>O<sub>2</sub> at 100 μM concentration induced characteristic morphological changes, such as cell shrinkage, appearance of a

rounded fruiting body and loss of adhesion capacity compared to the control (Figure 6A). HepG2 cells exposed to 400, 800 and 1000 μg/mL concentrations of polysaccharide extract were restored to their original morphology similar to the control in a concentration-dependent manner (Figure 6C, 6D and 6E, respectively). Preservation of the normal morphology of HepG2 cells observed by fluorescence microscopy indicated the preventive role of the compounds against H<sub>2</sub>O<sub>2</sub>-induced morphological damage.




**Figure 6** Detection of nuclear morphological changes in HepG2 by Hoechst 33342 staining, observed by fluorescence microscopy at 20x magnification. Control group (A), H<sub>2</sub>O<sub>2</sub> 100 µM (B), polysaccharide 400 µg/mL + H<sub>2</sub>O<sub>2</sub> 100 µM (C), polysaccharide 800 µg/mL + H<sub>2</sub>O<sub>2</sub> 100 µM (D) and polysaccharide 1000 µg/mL + H<sub>2</sub>O<sub>2</sub> 100 µM (E).

#### 4.5 Protective effect of polysaccharide extract on lipid peroxidation

To evaluate the protective effect of the polysaccharide extract on oxidative cell damage caused by H<sub>2</sub>O<sub>2</sub>, lipid peroxidation was measured as the formation of MDA in the homogenates of HepG2 cells (Figure 7). The increase in lipid peroxidation in HepG2 cells by H<sub>2</sub>O<sub>2</sub> may be due to enhancement in hepatic MDA from the peroxidation of polyunsaturated fatty acids. In this

assay, percentage inhibition of lipid peroxidation of the polysaccharide extract significantly reduced lipid peroxidation induced by H<sub>2</sub>O<sub>2</sub> as 375.38±35.16% at 400 µg/mL, 231.54±0.94% at 800 µg/mL and 157.69±51.14% at 1000 µg/mL, respectively compared with cells treated with H<sub>2</sub>O<sub>2</sub>. This result indicated that the polysaccharide extract displayed a significant protective capability against H<sub>2</sub>O<sub>2</sub>-induced cell damage.




**Figure 7** Effects of polysaccharide extract on lipid peroxidation (LPO) level of HepG2 cells following H<sub>2</sub>O<sub>2</sub>-induced oxidative damage. Cells ( $1 \times 10^5$  cells in 6-well plates) were treated with polysaccharide extract for 24 h before H<sub>2</sub>O<sub>2</sub>-induced oxidation. Values represent mean  $\pm$  SE. \*\* $p < 0.01$  versus H<sub>2</sub>O<sub>2</sub> exposure.

#### 4.6 Protective effect of polysaccharide extract on glutathione level

To evaluate the effect of the polysaccharide extract on antioxidant defense systems, the glutathione level was measured in H<sub>2</sub>O<sub>2</sub> exposed HepG2 cells. Glutathione is a main non-enzymatic antioxidant and plays an important role in protective effects against oxidative stress. (Scharf et al., 2003). Our results showed that the glutathione level in H<sub>2</sub>O<sub>2</sub> exposed HepG2 cells significantly decreased by 67.21% ( $p < 0.001$ ) compared to the control (Figure 8). However, exposure of cells to polysaccharide extract significantly increased the glutathione level in H<sub>2</sub>O<sub>2</sub> exposed HepG2 cells in a dose-dependent manner. Increases in GSH level were  $77.76 \pm 4.36\%$  at 400

$\mu\text{g/mL}$ ,  $85.96 \pm 6.44\%$  at 800  $\mu\text{g/mL}$  and  $95.39 \pm 2.15\%$  at 1000  $\mu\text{g/mL}$ , respectively (Figure 8). Our results showed that pre-exposure of cells to the polysaccharide extract effectively restored the glutathione level. It is well known that exogenous exposure to H<sub>2</sub>O<sub>2</sub> can increase intracellular ROS generation and induce cellular oxidative damage in hepatic cells, while hepatocyte antioxidant defense mechanisms can eliminate this. Glutathione is a main non-enzymatic antioxidant, which plays an important role in the cellular defense system against oxidative stress (Scharf et al., 2003). Exposure of HepG2 cells to the polysaccharide extract quenched the intracellular destructive peroxide and increased the concentration of reduced glutathione



**Figure 8** Effects of polysaccharide extract on glutathione(GSH) level of HepG2 cells following H<sub>2</sub>O<sub>2</sub>- induced oxidative damage. Cells ( $1 \times 10^5$  cells in 6-well plates) were pre-treated with polysaccharide extract for 24 h before H<sub>2</sub>O<sub>2</sub>-induced oxidation. Values represent mean  $\pm$  SE

\* $p < 0.05$ , \*\* $p < 0.01$  versus H<sub>2</sub>O<sub>2</sub> exposure; ###  $p < 0.001$  H<sub>2</sub>O<sub>2</sub> versus the control.

## 5. Conclusions

Our results demonstrated that the scavenging activity and cytoprotective potential of the polysaccharide extract attenuated H<sub>2</sub>O<sub>2</sub>-induced cytotoxicity in HepG2 cells. The protective effect was mediated by both inhibiting lipid peroxidation and restoring glutathione levels. Findings suggested that constituents in the fruiting body of *S. commune* can protect HepG2 cells against H<sub>2</sub>O<sub>2</sub>-induced oxidative damage through antioxidant activity. Further work is necessary to find specific active compounds and elucidate the extracting mechanism involved in the hepatoprotective activity. Moreover, the animal model should be good feasibility to prove the ability of the polysaccharide extract of *S. commune* for hepatoprotective.

## 6. Acknowledgements

The authors greatly appreciated the support and help of the late Dr. Pongpun Siripong, who sadly passed away before the completion of this paper. We also acknowledge the College of Agricultural Innovation and Food Technology, Rangsit University, Pathum Thani 12000, Thailand, and the Clinical Research Section, Division of Research and Academic Support, National Cancer Institute, Bangkok 10400, Thailand and the School of Microbiology, Institute of Science Suranaree University of Technology, Nakhon Ratchasima 30000, Thailand, for facilities support.

## 7. References

Al-Sheddi, E. S., Farshori, N. N., Al-Oqail, M. M., Musarrat, J., Al-Khedhairi, A. A., &

- Siddiqui, M. A. (2016). Protective effect of *Lepidium sativum* seed extract against hydrogen peroxide-induced cytotoxicity and oxidative stress in human liver cells (HepG2). *Pharmaceutical Biology*, 54(2), 314-321. DOI: 10.3109/13880209.2015.1035795
- Aletor, V. A. (1995). Compositional studies on edible tropical species of mushrooms. *Food Chemistry*, 54(3), 265-268. DOI : 10.1016/0308-8146(95)00044-J
- Ali, M., Khan, S. A., Chang, P. S., Haque, R., Bhatia, K., & Ahmad, S. (2014). Habb-e-Asgand, polyherbal Unani formulation, protects liver and antioxidative enzymes against paracetamol induced hepatotoxicity. *Pharmaceutical Biology*, 52, 506-515. DOI: 10.3109/13880209.2013.863949
- Aruoma, O. I., Grootveld, M., & Bahorun, T. (2006). Free radicals in biology and medicine: from inflammation to biotechnology. *Biofactors*, 27(1-4), 1-3. DOI: 10.1002/biof.5520270101
- Bradford, M. M. (1976). A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Analytical biochemistry*, 72(1-2), 248-254. DOI: 10.1006/abio.1976.9999
- Bugrahan, E., Aytac, K., Abdullah, K., Seval, C., Muhamud, A., & Gokhan, S. (2017). In vitro cytotoxicity, antibacterial and antioxidant properties of various extracts from *Schizophyllum commune* fr. *Fresenius Environmental Bulletin*, 26(1), 1144-1153 .
- Chandra, D., Ramana, K. V., Wang, L., Christensen, B. N., Bhatnagar, A., & Srivastava, S. K. (2002). Inhibition of fiber cell globalization and hyperglycemia-induced lens opacification by aminopeptidase inhibitor bestatin. *Investigative Ophthalmology & Visual Science*, 43(7), 2285-2292.
- Chourasiya, R. R. & Sharma, S. (2021). Impact of dietary components and food additives on lipid peroxidation (LPO) product in *Drosophila melanogaster*. *Annals of Biochemistry and Biotechnology*, 1(1), 1001.
- Ghasemi, K., Ghasemi, Y., & Ebrahimzadeh, M. A. (2009). Antioxidant activity, phenol and flavonoid contents of 13 citrus species peels and tissues. *Pakistan Journal of Pharmaceutical Sciences*, 22(3), 277-281.
- Jaydeokar, A. V., Bandawane, D. D., Bibave, K. H., & Patil, T. V. (2014). Hepatoprotective potential of *Cassia auriculata* roots on ethanol and antitubercular drug-induced hepatotoxicity in experimental models. *Pharmaceutical Biology*, 52(3), 344-355. DOI: 10.3109/13880209.2013.837075
- Joel, E. L., & Bhimba, B. V. (2013). A secondary metabolite with antibacterial activity produced by mangrove foliar fungus *Schizophyllum commune*. *International Journal of Chemistry, Environment and Biological Science*, 1(1), 2320-4087.
- Kao, C., Jesuthasan, A. C., Bishop, K. S., Glucina, M. P., & Ferguson, L. R. (2013). Anti-cancer activities of *Ganoderma lucidum*: active ingredients and pathways. *Functional Foods Health Disease*, 3(2), 48-65. DOI :10.31989/ffhd.v3i2.65
- Kukan, M., Szatmáry, Z., Lutterová, M., Kuba, D., Vajdová, K., & Horecký, J. (2004). Effects of sizofiran on endotoxin-enhanced cold ischemia-reperfusion injury of the rat liver. *Physiological Research*, 53(4), 431-437.
- Kumar, A., Kumar, M., Sinha, M. P. (2019). Hepatoprotective efficacy edible macrofungi *Dacryopinax Spathularia* (Schwein) and *Schizophyllum commune* (Fries) against Carbon tetrachloride induced hepatotoxicity in albino Wister rats. *Journal of Applied and Natural Science*, 11(1), 62-65. DOI:10.31018/jans.v11i1.1959
- Lindequist, U., Niedermeyer, T. H. J., & Jülich, W. D. (2005). The Pharmacological Potential of Mushrooms. *Evidence-Based Complimentary and Alternative Medicine*, 2(3), 285-299. DOI : 10.1093/ecam/neh107
- Lister, I. N. E., Ginting, C. N., Girsang, E., Nataya, E. D., Azizah, A. M., & Widowati, W. (2020). Hepatoprotective properties of red betel (*Piper crocatum* Ruiz and Pav) leaves extract towards

- H<sub>2</sub>O<sub>2</sub>-induced HepG2 cells via anti-inflammatory, antinecrotic, antioxidant potency. *Saudi Pharmaceutical Journal*, 28(10), 1182-1189. DOI : 10.1016/j.jpsps.2020.08.007
- Longvah, T., & Deosthale, Y. G. (1998). Compositional and nutritional studies on edible wild mushroom from northeast India. *Food Chemistry*, 63(3), 331-334. DOI: 10.1016/S0308-8146(98)00026-0
- Masuda, Y., Ito, K., Konishi, M., & Nanba, H. (2010). A polysaccharide extracted from *Grifola frondosa* enhances the anti-tumor activity of bone marrow-derived dendritic cell-based immunotherapy against murine colon cancer. *Cancer Immunol Immunotherapy*, 59(10), 1531-1541. DOI: 10.1007/s00262-010-0880-7
- Masuko, T., Minami, A., Iwasaki, N., Majima, T., Nishimura, S.-I., & Lee, Y. C. (2005). Carbohydrate analysis by a phenol-sulfuric acid method in microplate format. *Analytical Biochemistry*, 339(1), 69-72. DOI: 10.1016/j.ab.2004.12.001
- Mau, J. L., Tsai, S. Y., Tseng, Y. H., & Huang, S. J. (2005). Antioxidant properties of hot water extracts from *Ganoderma lucidum* Murrill. *LWT - Food Science and Technology*, 38(6), 589-597. DOI:10.1016/j.lwt.2004.08.010
- Mayakrishnan, V., Abdullah, N., Hamdi, M., Abidin, Z. U., Fadzil, N. H. M., Johari, N. M. K., Abidin, N. Z., & Lumpur, K. (2013). Investigation of the Antioxidative Potential of Various Solvent Fractions from Fruiting Bodies of *Schizophyllum commune* (Fr.) Mushrooms and Characterization of Phytoconstituents. *The Journal of Agricultural Science*, 5, 58. DOI: 10.5539/jas.v5n6p58
- Mirfat, A. H. S., Noorlidah, A., & Vikineswary, S. (2010). Scavenging activity of *Schizophyllum commune* extracts and its correlation to total phenolic content. *Journal of Tropical Agriculture and Food Science*, 38(2), 231-238.
- Ooi, V. E., & Liu, F. (2000). Immunomodulation and anti-cancer activity of polysaccharide-protein complexes. *Current Medicinal Chemistry*, 7(7), 715-729. DOI: 10.2174/0929867003374705
- Phonghanpot, S., & Jarintanan, F. (2021). Antiproliferative, antibacterial, and antioxidant activities of *Bauhinia strychnifolia* Craib aqueous extracts in gut and liver perspective. *BMC Complementary Medicine and Therapies*, 21(1), 276. DOI:10.1186/s12906-021-03448-2
- Scharf, G., Prustomersky, S., Knasmüller, S., Schulte-Hermann, R., & Huber, W. W. (2003). Enhancement of glutathione and g-glutamylcysteine synthetase, the rate limiting enzyme of glutathione synthesis, by chemoprotective plant-derived food and beverage components in the human hepatoma cell line HepG2. *Nutrition and Cancer*, 45(1), 74-83. DOI: 10.1207/S15327914NC4501\_9
- Selima, K., Aminul, I., Ugur, C., & Narayan, C. (2012). Research on Mushroom as a Potential Source of Nutraceuticals :A Review on Indian Perspective. *American Journal of Experimental Agriculture*, 2(1), 47-73. DOI: 10.9734/AJEA/2012/492
- Siripong, P., Rassamee, K., Piyaviriyakul, S., Yahuafai, J., & Kanokmedhakul, K. (2012). Anti-metastatic effects on B16F10 melanoma cells of extracts and two prenylated xanthenes isolated from *Maclura amboinensis* BI roots. *The Asian Pacific Journal of Cancer Prevention*, 13(7), 3519-3528. DOI: 10.7314/apjcp.2012.13.7.3519
- Soares, A. A., de Sá-Nakanishi, A. B., Bracht, A., da Costa, S. M., Koehnlein, E. A., de Souza, C. G., & Peralta, R. M. (2013). Hepatoprotective effects of mushrooms. *Molecules*, 18(7), 7609-7630. DOI : 10.3390/molecules18077609
- Su, C. H., Lai, M. N., Lin, C. C., & Ng, L. T. (2016). Comparative characterization of physicochemical properties and bioactivities of polysaccharides from selected medicinal mushrooms. *Applied Microbiology and Biotechnology*, 100(10), 4385-4393. DOI: 10.1007/s00253-015-7260-3
- Venturella, G., Ferraro, V., Cirlincione, F., & Gargano, M. L. (2021). Medicinal Mushrooms: Bioactive Compounds, Use, and Clinical Trials. *International journal*

- of molecular sciences*, 22(2), 634. DOI: 10.3390/ijms22020634.
- Wu, X. J., & Hansen, C. (2008). Antioxidant capacity, phenolic content, and polysaccharide content of *Lentinus edodes* grown in whey permeate-based submerged culture. *Journal of Food Science*, 73(1), M1-8. DOI: 10.1111/j.1750-3841.2007.00595.x
- Yoon, E., Babar, A., Choudhary, M., Kutner, M., & Pysopoulos, N. (2016). Acetaminophen-Induced Hepatotoxicity: a Comprehensive Update. *Journal of clinical and translational hepatology*, 4(2), 131-142. DOI: 10.14218/JCTH.2015.00052
- Zhang, Y., Kong, H., Fang, Y., Nishinari, K., & Phillips, G. O. (2013). Schizophyllan: A review on its structure, properties, bioactivities and recent developments. *Bioactive Carbohydrates and Dietary Fibre*, 1(1), 53-71. DOI : 10.1016/j.bcdf.2013.01.002
- Zhong, K., Tong, L., Liu, L., Zhou, X., Liu, X., Zhang, Q., & Zhou, S. (2015). Immunoregulatory and antitumor activity of schizophyllan under ultrasonic treatment. *International Journal of Biological Macromolecules*, 80, 302-308. DOI: 10.1016/j.ijbiomac.2015.06.052