



Implant-Prosthetic Rehabilitation of Patients with Orthopedic Structures & Extreme Atrophy of the Upper Jaw During Prosthetics on Zygomatic and Angular Implants

Mihail Artunovich Avanesov¹, Lamara Georgievna Sherasova², Veronika Stanislavovna Dzoblaeva³, Raisa Kadirovna Sabanova⁴, Ibrahim Mussayevich Aushev¹, Yulia Vadimovna Maskurova^{3,5}, Ruslan Kazbekovich Yesiev⁵, Ilina Makharebekovna Shavlokhova⁵, Asya Omarovna Magdilova⁶, Semyon Alexandrovich Kuchevskiy¹, Anna Valerievna Klopova⁷, Natalya Yurievna Basova^{8*}

¹ Stavropol State Medical University, Stavropol, RUSSIA.

² Rostov State Medical University, Rostov-on-Don, RUSSIA.

³ North Ossetian State Medical Academy, Vladikavkaz, Republic of North Ossetia-Alania, RUSSIA.

⁴ Kabardino-Balkarian State University, Nalchik, Republic of Kabardino-Balkaria, RUSSIA.

⁵ North Ossetian State University named after K. L. Khetagurov, Vladikavkaz, Republic of North Ossetia-Alania, RUSSIA.

⁶ Dagestan State Medical University, Makhachkala, Republic of Dagestan, RUSSIA.

⁷ Don State Agrarian University, Persianovskiy Village, Oktyabrsky District, Rostov Region, RUSSIA.

⁸ Krasnodar Research Veterinary Institute – separate structural division of Federal State Budgetary Scientific Institution Krasnodar scientific center for animal and veterinary science, Krasnodar, RUSSIA.

*Corresponding Author (Tel: +79183500889, Email: ruslankalmykov777@yandex.ru).

Paper ID: 13A11C

Volume 13 Issue 11

Received 03 April 2022

Received in revised form 24 June 2022

Accepted 02 July 2022

Available online 15 July 2022

Keywords:

Maxillofacial pathology; Implant-prosthetic rehabilitation; Dentition defects; Maxillary atrophy; Zygomatic implants; Angular implants.

Abstract

Complex maxillofacial pathology is characterized by pronounced violations of the maxillary structure and abnormal disfigurement of the patient's face. Implant-prosthetic rehabilitation of patients is quite popular in modern reality. It is not so much about the aesthetic side of the issue, as about the physiological necessity. Extensive defects of the dentition after surgical interventions, which are most often accompanied by atrophy of the jaw tissues, inflammatory processes, neoplasms, etc. jaws require a special comprehensive approach to the rehabilitation of patients with orthopedic structures. Zygomatic implants are a rehabilitation option for patients with extreme bone atrophy in the upper jaw. The purpose of this scientific work is to analyze scientific publications describing cases of implant-prosthetic rehabilitation of patients with orthopedic structures with extreme atrophy of the upper jaw during prosthetics on zygomatic and angular implants.

Discipline: Medicine, Dentistry, Orthodontics, Surgery

©2022 INT TRANS J ENG MANAG SCI TECH.

Cite This Article:

Avanesov, M.A., Sherasova, L.G., Dzoblaeva, V.S., Sabanova, R.K., Aushev, I.M., Maskurova, Yu.V., ... Basova N.Yu. (2022). Implant-Prosthetic Rehabilitation of Patients with Orthopedic Structures & Extreme Atrophy of the Upper Jaw During Prosthetics on Zygomatic and Angular Implants. *International Transaction Journal of Engineering, Management, & Applied Sciences & Technologies*, 13(11), 13A11C, 1-10. <http://TUENGR.COM/V13/13A11C.pdf> DOI: 10.14456/ITJEMAST.2022.213

1 Introduction

Traditional removable dentures cannot always eliminate defects associated with extreme loss of jaw tissue, extensive loss of dentition. Restoration of speech, swallowing effects is possible almost in full, but chewing functions will cause the patient difficulty [1,2]. Moreover, such difficulties occur not only during the rehabilitation period, but also after the complete completion of treatment [3,4]. Most often, traditional dental implants are used, which perform the function of stabilizing the prosthesis, maintaining its resistance to horizontal loads during chewing. Cases, when such prostheses are insufficient, are associated with extreme tissue loss during surgical intervention, resection of extensive areas of bone tissue [5].

Zygomatic and angular prostheses are used to restore the upper jaw, the method consists in installing the implant at a special angle, which is determined by the presence of bone tissue, excluding bone block transplantation and lateralization of the nervus mandibularis [6-8].

2 Literary review

The installation of prostheses on zygomatic and angular implants proved to be effective. Complications and risks for the patient are minimized, provided that all the recommendations of the specialist are followed. Inflammatory processes, fistulas, ulceration of tissue occur only through the fault of the patient himself [9-11].

Delayed prosthetics after tooth extraction leads to a number of structural changes in the bone tissue: bone density decreases, the spongy layer begins to decrease, atrophy, osteopenia occurs, and then osteoporosis of the jaw [12]. These phenomena do not always depend on the use of prostheses, genetic predisposition and other factors affecting the patient's health also affect here.

The advantages of zygomatic implants are the recovery time and economic feasibility. Such implants are much cheaper than bone grafting, they allow you to combine implantation and resection of tissues to be removed in one operation [13].

Patients with traumatic lesions of the maxillofacial region, age-related patients, oncopatients have problems with fixing prosthetic structures, both removable and non-removable [14].

To solve this problem, protocols of transculular, angular implantation and intraoperative prosthetics are used [15].

Modern orthopedic planning of the protocol of operations makes it possible to abandon, among other things, the use of angular abutments, which leads not only to simplification and cheaper construction but also to increase its reliability. Implants with the possibility of intercortical and polycortical fixation are also used to achieve the tasks set [16-18]. Due to the fact that transculary implantation protocols are performed, as a rule, in the practice of maxillofacial surgeons, this technology, unfortunately, is limited in outpatient dental practice [7,14,19].

3 Method

The main direction in angular implantology is the rehabilitation of the upper jaw using implants of various types in combination with angular implants.

Information about the technology of installing zygomatic and angular implants has been available since 1989 and is actively developing in many countries. In the Russian periodical scientific press, information about the use of implant-prosthetic technology for zygomatic and angular implants has been found since 2006. The effectiveness and complexity of this protocol and the impossibility of using such a technique in outpatient settings are noted.

Observations of more than 3,000 patients were carried out in 11 private and public clinics of the Russian Federation. The age of the patients ranged from 13-92 years. Indications for prosthetics in 80 percent of cases – bone atrophy, two-tenths of a percent accounted for oncological diseases, the remaining cases are associated with injuries.

Prosthetics on zygomatic implants can improve the quality of life of patients with complex cases. Zygomatic and angular implants exclude the use of bone grafting with a long period of tissue stabilization for subsequent implantation of implants and installation of prostheses on implants [20].

Rods of 30-60 mm in size are screwed into the subglacial (zygomatic) bone tissue, which is little susceptible to destructive changes [21]. Today, two types of systems are used: straight and beveled platforms [22]. The edge of the beveled rod is installed parallel to the gum, which does not require the use of additional structures, therefore, the maximum simplification of the fixation of the orthoprosthetic system occurs. Rods differ in their properties depending on the manufacturer. Some have a smooth structure and antibacterial coating, while others, on the contrary, are porous and have micro-sections [23-25]. The choice of a rod for implantation into the bones of the upper jaw depends on the protocol of orthodontic treatment and indications for use (Figure 1).




Figure 1: Installation of a titanium rod in the zygomatic bone

Implantation of the rod into the bone zygomatic tissue is not a special protocol of dental prosthetics. There are three ways to use this technique in the complete absence of the dentition of the upper jaw [26]:

- install 4 roots – 2 front and 2 sides;
- install 6 roots, all can be zygomatic (figure 2);
- install 8-14 roots at an angle in the lateral sections.




Figure 2: X-ray image with the demonstration of implanted zygomatic implants (6 roots)

Figure 3 schematically shows some options for implanting titanium rods into the zygomatic tissue and bone tissue of the upper jaw.



Figure 3: Options for implanting implants in the bone tissue of the upper jaw

This technique is applicable for extreme atrophy of the upper jaw, it does not require bone grafting and a long period of rehabilitation with loss of aesthetic appearance of the patient. A prosthesis can be installed on zygomatic implants immediately, or 3-4 days after implantation. Such implants can participate in the restoration of the masticatory group of the dentition [27].

The installation of prostheses immediately after surgery eliminates the loosening of the implants, they can serve the patient until the end of his life. It only requires the replacement of the orthopedic systems themselves as they wear out.

Indications for the installation of zygomatic and angular implants are:

- complete or partial loss of upper jaw teeth;
- extreme atrophy of the upper jaw due to resection of degenerative tissues;
- periodontal disease (advanced stage);
- periodontitis (advanced stage);
- osteopenia;
- osteoporosis;
- oncological diseases followed by radiation therapy.

This methodology is characterized by the following positive characteristics, which make it attractive in maxillofacial surgery:

- accelerated treatment protocol – from implantation to prosthetics takes from one to seven days;
- there are practically no contraindications;

- applicable even in cases of extreme maxillary atrophy: absence of the alveolar ridge, complete or partial absence of the maxillary bone due to cancer or injury;
- no bone buildup is required, titanium roots are implanted into the dense zygomatic bone at a certain angle;
- a prosthesis is immediately installed, which looks like a natural dentition (Figure 4);
- the materials from which the prosthesis is made (metal-plastic or ceramic composite) do not create excessive load on the implants during physiological movements of the jaw, including chewing;
- reliable fixation of implants in the zygomatic bone;
- a high probability of survival due to the absence of inflammatory processes in the zygomatic bone;
- durability of titanium implants.



Figure 4: Orthodontic prosthesis installed on zygomatic and angular implants

Nevertheless, as mentioned above, this technique is practically not applicable in outpatient settings [28]. Such orthodontic treatment can only be performed in a hospital by a maxillofacial surgeon.

The preparatory and operational period includes a number of stages [29]:

- a complete examination of the patient for contraindications to surgery;
- computed tomography of the jaw;
- 3D modeling of a copy of the jaw;
- using a 3D printer to print out a full-size jaw as a stand for an upcoming operation;
- installation of dummy rods in a 3D model for precise positioning of implants in the zygomatic bone of the patient;
- the installation of zygomatic implants takes place at an angle of 30-60 °, which requires jewelry precision in the area of the nasal sinuses in order to avoid injury to them;

- simultaneous installation of a prosthesis with a metal beam that stabilizes the pressure on the implants during chewing is possible;
- the operation is performed under general anesthesia.

The use of angular implants reduces the time of dental rehabilitation of patients, minimizes injuries from surgical intervention, tissue regeneration occurs much faster, and the implantability of such implants is ninety-eight percent. The quality of regeneration also depends on the patient and their compliance with hygienic standards of oral care. It is necessary to pay attention to patients with loss of motor functions of the limbs who cannot take care of the oral cavity on their own, the exclusion of complications in the postoperative period depends on this [30,31].

4 Result and Discussion

Studies in the field of the installation of zygomatic implants, which are published and available to the scientific community, are almost unambiguous in their conclusions: this technique is effective, and gives an opportunity for a high-quality life to patients with neglected or hopeless cases of upper jaw deformity.

The use of angular, trans-musculoskeletal and polycortical implants, both according to the original protocol and in a combination of protocols, reduces the time of dental and maxillofacial rehabilitation and reduces the traumatism of operations. The possibility of installing temporary screw prostheses in the oral cavity is realized – direct intraoperative prosthetics within 0 – 72 hours after surgery.

5 Conclusion

Thus, the protocols of angular, trans-musculoskeletal and polycortical implantation expand the possibilities of dental and maxillofacial rehabilitation and are one of the most actively developing areas in pre-prosthetic reconstructive surgery. Modern orthopedic planning of such an operation protocol makes it possible to abandon the use of cement fixation and angular abutments, which leads not only to simplification and cheaper construction but also to an increase in its reliability. To master the protocols of transculular, angular and polycortical implantation, training of specialists is required within the framework of a new platform of interdisciplinary cooperation, that is, the formation of practical skills in a team of specialists.

Implantation of artificial roots into the zygomatic bone practically leveled the statistics of hopeless dental cases, improving the quality of life of patients with extreme maxillary atrophy. This type of surgery is impossible without the participation of a maxillofacial surgeon and an anesthesiologist, the type of clinical research assumes the presence of high-tech modern equipment in the clinic. The cost of zygomatic implants is quite high, but the material from which they are made, and the method of their implantation, guarantees their lifelong use with periodic replacement of the external dental prosthesis as it wears out.

6 Availability of Data and Material

Data can be made available by contacting the corresponding author.

7 References

- [1] Remizova A A, Dzgoeva M G, Tingaeva Y I, Hubulov S A, Gutnov V M, Bitarov P A, et al. Tissue Dental Status and Features of Periodontal Microcirculation in Patients with New COVID-19 Coronavirus Infection. *Pharmacophore*. 2021;12(2):6-13. <https://doi.org/10.51847/5JIbnUbHkT>
- [2] Ayivi R, Ibrahim S, Colleran H, Silva R, Williams L, Galanakis C, Fidan H, Tomovska J and Siddiqui SA. COVID-19: human immune response and the influence of food ingredients and active compounds. *Bioactive Compounds in Health and Disease*. 2021;4(6):100
- [3] Galabueva AI, Biragova AK, Kotsoyeva GA, Borukayeva ZK, Yesiev RK, Dzgoeva ZG, et al. Optimization of Modern Methods of Treating Chronic Generalized Periodontitis of Mild Severity. *Pharmacophore* 2020;11(1):47-51
- [4] Dmitry Anatolyevich Domenyuk, Vladimir Alexandrovich Zelensky, Igor Vladimirovich Rzhepakovsky, Oksana Ivanovna Anfinogenova, Application of Laboratory and X-Ray General Studies on Early Diagnostics of Metabolic Disturbances of Bone Tissue in Children with Autoimmune Diabetes Mellitus, *EntomolApplSci Lett*. 2018; 5(4):1-12
- [5] Sadyrin E, Swain M, Mitrin B, Rzhepakovsky I, Nikolaev A, Irkha V, Yogina D, Lyanguzov N, Maksyukov S, Aizikovich S. Characterization of Enamel and Dentine about a White Spot Lesion: Mechanical Properties, Mineral Density, Microstructure and Molecular Composition. *Nanomaterials (Basel)*. 2020;10(9):1889. DOI: 10.3390/nano10091889
- [6] El Fattah H, Zaghloul A, Pedemonte E, Escuin T. Pre-prosthetic surgical alterations in maxillectomy to enhance the prosthetic prognoses as part of rehabilitation of oral cancer patient. *Med Oral Patol Oral Cir Bucal*. 2012 Mar 1;17(2):e262-70. doi: 10.4317/medoral.17482.
- [7] Ali Redha, Ali & Siddiqui, Shahida & Ibrahim, Salam. Advanced extraction techniques for Berberis species phytochemicals: A review. *International Journal of Food Science & Technology*. 2021:1-12, doi:10.1111/ijfs.15315
- [8] Rauf A, Abu-Izneid T, Olatunde A, Ahmed Khalil A, Alhumaydhi FA, Tufail T, Shariati MA, Rebezov M, Almarhoon ZM, Mabkhot YN, Alsayari A, Rengasamy KRR. COVID-19 Pandemic: Epidemiology, Etiology, Conventional and Non-Conventional Therapies. *International Journal of Environmental Research and Public Health*. 2020; 17(21):8155. <https://doi.org/10.3390/ijerph17218155>
- [9] Rzhepakovsky IV, Areshidze DA, Avanesyan SS, Grimm WD, Filatova NV, Kalinin AV, Kochergin SG, Kozlova MA, Kurchenko VP, Sizonenko MN, Terentiev AA, Timchenko LD, Trigub MM, Nagdalian AA, Piskov SI. Phytochemical Characterization, Antioxidant Activity, and Cytotoxicity of Methanolic Leaf Extract of *Chlorophytum Comosum* (Green Type) (Thunb.) Jacq. *Molecules*. 2022; 27(3):762. <https://doi.org/10.3390/molecules27030762>
- [10] Siddiqui SA, Ali Redha A, Snoeck ER, Singh S, Simal-Gandara J, Ibrahim SA, Jafari SM. Anti-Depressant Properties of Crocin Molecules in Saffron. *Molecules*. 2022; 27(7):2076. <https://doi.org/10.3390/molecules27072076>
- [11] Ranjha MMAN, Shafique B, Rehman A, Mehmood A, Ali A, Zahra SM, Roobab U, Singh A, Ibrahim SA and Siddiqui SA (2022) Biocompatible Nanomaterials in Food Science, Technology, and Nutrient Drug Delivery: Recent Developments and Applications. *Front. Nutr.* 8:778155. doi: 10.3389/fnut.2021.778155
- [12] Maisigov, J. B., Kuznetsova, G. V., Magomedov, A. M., Adzhigova, F. Z., Magomedova, A. S., Burdukova, S. A., Mishvelov, A. E. and Povetkin, S. N. (2021) "Anthropometric Analysis of Digital Models of the Dentition Using 3D Technologies in Orthodontics", *Journal of Pharmaceutical Research International*, 33(40A), pp. 101-105. doi: 10.9734/jpri/2021/v33i40A32225

- [13] Davó R, Bankauskas S, Laurincikas R, Koçyigit ID, Mate Sanchez de Val JE. Clinical Performance of Zygomatic Implants—Retrospective Multicenter Study. *Journal of Clinical Medicine*. 2020; 9(2):480. <https://doi.org/10.3390/jcm9020480>
- [14] Eldzharov, A. V., Niazyan, D. A., Esiev, R. K., Toboev, G. V., Uzdenova, J. K., Shabanova, Z. M., Abakarov, Z. A., Abrekov, A. B., Babatkhanov, A. S. and Mishvelov, A. E. (2021) “Clinical and Immunological Characteristics of Patients with Odontogenic Maxillary Sinusitis”, *Journal of Pharmaceutical Research International*, 33(50A), pp. 184-195. doi: 10.9734/jpri/2021/v33i50A33394
- [15] Feher B, Gruber R, Gahleitner A, Celar A, Necsea PL, Ulm C, Kuchler U. Angular changes in implants placed in the anterior maxillae of adults: a cephalometric pilot study. *Clin Oral Investig*. 2021 Mar;25(3):1375-1381. doi: 10.1007/s00784-020-03445-8.
- [16] Bledzhyants GA, Mishvelov AE, Nuzhnaya KV, Anfinogenova OI, Isakova JA, Melkonyan RS, et al. The effectiveness of the medical decision-making support system "electronic clinical pharmacologist" in the management of patients therapeutic profile. *Pharmacophore*. 2019;10(2): 76-81
- [17] Shahida Anusha Siddiqui, Nur Alim Bahmid, Ahmed Taha, Ibrahim Khalifa, Sipper Khan, Hadis Rostamabadi & Seid Mahdi Jafari (2022) Recent advances in food applications of phenolic-loaded micro/nanodelivery systems, *Critical Reviews in Food Science and Nutrition*, DOI: 10.1080/10408398.2022.2056870
- [18] Gutnova, T. S., Kompantsev, D. V., Gvozdenko, A. A., Kramarenko, V. N., & Blinov, A. V. VITAMIN D NANOCAPSULATION. *IZVESTIYA VYSSHIKH UCHEBNYKH ZAVEDENII KHIMIYA KHIMICHESKAYA TEKHNLOGIYA*, 20216 64(5), 98-105. <https://doi.org/10.6060/ivkkt.20216405.6399>
- [19] Kotsoeva GA, Esiev RK, Toboev GV, Zakaeva RS, Kulova AA, Tsokova LV, et al. Phytoadaptogenic Cocktail Use "Biorithm-E" In The Complex Treatment Of Odontogenic Inflammatory Diseases Of The Maxillofacial Region. *Ann. Dent. Spec*. 2021;9(2):52-57. <https://doi.org/10.51847/PyVv83OTGt>
- [20] Amato F, Polara G, Spedicato GA. Immediate Loading of Fixed Partial Dental Prostheses on Extra-Short and Short Implants in Patients with Severe Atrophy of the Posterior Maxilla or Mandible: An Up-to-4-year Clinical Study. *Int J Oral Maxillofac Implants*. 2020 May/Jun;35(3):607-615. doi: 10.11607/jomi.7943.
- [21] Yalçın M, Can S, Akbaş M, Dergin G, Garip H, Aydil BA, Varol A. Retrospective Analysis of Zygomatic Implants for Maxillary Prosthetic Rehabilitation. *Int J Oral Maxillofac Implants*. 2020 Jul/Aug;35(4):750-756. doi: 10.11607/jomi.8196.
- [22] Orsaeva AT, Tamrieva LA, Mischvelov AE, Osadchiy SS, Osipchuk GV, Poveikin SN, Simonov AN. Digital clinic “smart ward. *Pharmacophore*. 2020;11(1):142-146
- [23] Aparicio C, Manresa C, Francisco K, Claros P, Alánde J, González-Martín O, Albrektsson T. Zygomatic implants: indications, techniques and outcomes, and the zygomatic success code. *Periodontol* 2000. 2014 Oct;66(1):41-58. doi: 10.1111/prd.12038.
- [24] Rawal S, Balshi T, Jivraj S. Restoration of Zygomatic Implants. *Atlas Oral Maxillofac Surg Clin North Am*. 2021 Sep;29(2):291-299. doi: 10.1016/j.cxom.2021.04.006.
- [25] Mishununa VV, Chapanov MM, Gakaeva KI, Tsoroeva MB, Kazanova SA, Gorlova MI, et al. Computed Quantum Chemical Modeling of the Effect of Nanosilver on Coronavirus COVID-19. *Pharmacophore*. 2021;12(2):14-21
- [26] Zaninovich M. Clinical guidelines for rehabilitation of the severely atrophic maxilla using extended-length subcrestal angulated implants and modified trans-sinus nasal protocol: A case report. *Int J Oral*

- [27] Mocanu RG, Florica LI, Preoteasa CT, Meghea MD, Preoteasa E. Periimplant Bone Resorption at the Level of Tilted Implants in SKY Fast & Fixed Restorations. J Med Life. 2020 Jul-Sep;13(3):336-341. doi: 10.25122/jml-2020-0131.
- [28] Tovlahanova TJH et al. Study of the Effect of the Image Scanning Speed and the Type of Conductive Coating on the Quality of Sem-Micrographs of Oxide Nano Materials for Medical Use. Ann Med Health Sci Res. 2021;11:S3:60-64
- [29] Yusupova MI, Mantikova KA, Kodzokova MA, Mishvelov AE, Paschenko AI, Ashurova ZAK, et al. Study Of The Possibilities Of Using Augmented Reality In Dentistry. Ann. Dent. Spec. 2021;9(2):17-21. Available:<https://doi.org/10.51847/BG1ZAqXRc>
- [30] Zotti Francesca, Dalessandri Domenico, Salgarello Stefano, Piancino Maria Grazia, Bonetti, Stefano & Visconti, Luca & Paganelli, Corrado. Usefulness of an app in improving oral hygiene compliance in adolescent orthodontic patients. The Angle orthodontist. 2015; 86. 10.2319/010915-19.1.
- [31] Remizova A A, Sakaeva Z U, Dzgoeva Z G, Rayushkin I I, Tingaeva Y I, Povetkin S N, et al. The Role Of Oral Hygiene In The Effectiveness Of Prosthetics On Dental Implants. Ann Dent Spec. 2021;9(1):39-46. <https://doi.org/10.51847/HuTuWdD0mB>
-




Mihail Artunovich Avanesov is a student of Stavropol State Medical University, Stavropol, Russia




Lamara Georgievna Sherasova is a student of Rostov State Medical University, Rostov-on-Don, Russia



Veronika Stanislavovna Dzoblaeva is a student of North Ossetian State Medical Academy, Vladikavkaz, Republic of North Ossetia-Alania, Russia



Raisa Kadirovna Sabanova is a lecturer of Kabardino-Balkarian State University, Nalchik, Republic of Kabardino-Balkaria, Russia



Ibrahim Mussayevich Aushev is a student of Stavropol State Medical University, Stavropol, Russia




Yulia Vadimovna Maskurova is a student of North Ossetian State Medical Academy, Student of North Ossetian State University named after K. L. Khetagurov, Vladikavkaz, Republic of North Ossetia-Alania, Russia



Ruslan Kazbekovich Yesiev is a student of North Ossetian State University named after K. L. Khetagurov, Vladikavkaz, Republic of North Ossetia-Alania, Russia



Ilina Makharbekovna Shavlokhova is a student of North Ossetian State University named after K. L. Khetagurov, Vladikavkaz, Republic of North Ossetia-Alania, Russia



Asya Omarovna Magdilova is a student of Dagestan State Medical University, Makhachkala, Republic of Dagestan, Russia



Semyon Alexandrovich Kuchevskiy is a student of Stavropol State Medical University, Stavropol, Russia



Dr. Anna Valerievna Klopova is an associate Professor, Department of Food Technology and Commodity Science, Don State Agrarian University, Persianovskiy Village, Oktyabrsky District, Rostov Region, Russia



Dr. Natalya Yurievna Basova is head of the Department of Therapy and Obstetrics, Doctor of Veterinary Sciences, Leading Researcher, Krasnodar Research Veterinary Institute – separate structural division of Federal State Budgetary Scientific Institution Krasnodar scientific center for animal and veterinary science, Krasnodar, Russia