

Original article

Comparative efficacy of topical dimethicone and permethrin for the treatment of head lice infestation in students

Behroz Davari ^a, Somayeh Barik - Abi^a, Hassan Nasirian^{b,c,*}
Amir Hossein Zahirnia ^a, Younes Mohammadi^d, Aref Salehzadeh ^a

^aDepartment of Medical Entomology, School of Medicine, Hamadan University of Medical Sciences, Hamadan, Islamic Republic of Iran

^bDepartment of Medical Entomology and Vector Control, School of Public Health, Tehran University of Medical Sciences, Tehran, Islamic Republic of Iran

^cInstitute for Environmental Research (IER), Tehran University of Medical Sciences, Tehran, Islamic Republic of Iran

^dDepartment of Epidemiology, School of Public Health, Hamadan University of Medical Sciences, Hamadan, Islamic Republic of Iran

Background: Despite improved living conditions, individual health and health indicators in human societies, head lice infestation continues to be a major health problem. Therefore, studying the efficacy of drugs and identifying treatment - resistant cases in the prevention and elimination of this disease is of particular importance.

Objective: The study aimed to determine the efficacy of dimethicone and permethrin in infected pupils in elementary schools for girls.

Methods: This study was conducted on primary schoolgirls infected with head lice in Doroud County, Lorestan-Iran. The eighty infected girls were detected and randomized into two groups receiving any of the treatments listed. By the end of second week, the efficacy of the drugs was determined.

Results: The efficacy of dimethicone and permethrin after treatment were 82.6% and 54.7%, respectively, for removing head lice at the end of two weeks. In other words, the efficacy of dimethicone was 27.9% higher than permethrin. The Chi-square test at the end of the first week, there were no significant differences between the treatment groups ($P = 0.065$). The outcomes of dimethicone treatment at the end of the second week ($P = 0.025$) and generally at the end of two weeks ($P = 0.006$) were more effective than permethrin.

Conclusion: The results of this study suggest that dimethicone is useful in the treatment of head lice compared to permethrin.

Keywords: Dimethicone, head lice, *Pediculus capitis*, permethrin.

Among the arthropod-pests of medical importance or arthropod associated diseases ^(1 - 13), *Pediculus*

capitis (De Geer) (Anoplura: Pediculidae) is mentioned as one of the most important human obligatory ectoparasites that generally infest the human hair and scalp. *Pediculus capitis* (head lice) is a global parasitic disease that can be considered a public health threat that occurs in different parts of the world, mainly affecting school-aged children with a maximum incidence of under 15 years, including Iran. Today, as in the past, head lice infestation remains a major health issue ⁽¹⁴⁾ despite improvements in living conditions, individual health and health indicators in human

*Correspondence to: Hassan Nasirian, Department of Medical Entomology and Vector Control, School of Public Health, Tehran University of Medical Sciences, Tehran, Islamic Republic of Iran; Institute for Environmental Research (IER), Tehran University of Medical Sciences, Tehran, Islamic Republic of Iran
E-mail: hanasirian@yahoo.com

Received : January 22, 2023

Revised: February 28, 2023

Accepted: March 30, 2023

societies. All stages of *P. capitis* are associated with human hosts for their lifetime. The nymphs and adult insects nourish only human blood without having wings and jumping ability. These insects are mostly spread by physical contact. (15 - 18) Topical medications are the first choice for managing head lice. In the first line of head lice treatment, topical medications such as dimethicone and permethrin are used. Resistance to topical medicines, particularly pyrethroids, is a growing concern around the world, and certain genetic mutations have been introduced to justify it. (19) Dimethicone 4% lotion is a recent new product development in treatment of head louse infestation. Dimethicone [polydimethylsiloxane, $(\text{CH}_3)_3\text{SiO}[\text{SiO}(\text{CH}_3)_2]_n\text{Si}(\text{CH}_3)_3$] is a silicon-based polymer used as a lubricant and conditioning agent. The mode of action of the product was to kill the insects by physical mechanisms rather than by neurotoxicity. However, the dimethicone mode of action has been unclear. Other investigators have loosely referred to the mode of action of different molecular weight products, as well as straightforward occlusive creams, as “asphyxiation” or “suffocation” but without clear biological evidence in support of their claims. In Iran, head lice infestation is emerging as a health problem due to the increasing villager migration to the cities, marginalization, and the creation of satellite cities with minimal health and welfare facilities. The prevalence rate of head lice infestation in primary school students was reported at 7.4%, with 1.6% for boys and 8.8% for girls. (20) Therefore, studying the efficacy of drugs and identifying treatment-resistant cases in the prevention and elimination of this disease is of particular importance. This study was conducted to determine the efficacy of dimethicone and permethrin in infected pupils in elementary schools for girls in Doroud County, Lorestan-Iran.

Materials and methods

Detection of infected students

This cross-sectional and interventional study was conducted on infected primary school girls in Doroud County, Lorestan-Iran. The necessary permits were obtained from the Doroud County Education Office and the Health and Medical Network Office prior to field work. To explain the study objectives, a coordination and justification meeting was held with school administrators, health educators and other stakeholders. Then, in agreement with the school administrators and health educators, we went to

primary schools to identify the infected students on certain days and according to the previous agreement. The county was divided into five geographical areas, including the west, east, north, south and center. In each area, a few girls' schools were randomly selected and a sampling was conducted. It was estimated that there were 896 students needed in accordance with the below formula used to estimate sample size. Additional required information was gathered from student files or questions and included on the checklist. While meeting the students, ethical principles were observed.

How to estimate sample size

The sample size was estimated in accordance with the formula used to estimate the ratio and prevalence of infected cases in previous studies. Based on consideration of 95.0% confidence level and accuracy, “ $Z_{1-\alpha/2} = 1.96$ ” and “ $d = 0.05$ ”. In past studies, “ P ” was “0.8”, therefore “ q ” would be “0.2”. By putting the numbers into the following formula, the sample size was estimated at 682 students. To complete the study in five schools, it was expanded to 896 students.

To estimate the sample size to compare the effect of two topical medications, the Devore CD, *et al.* method (21) was used using the following formula. The number of students in each group was established as “6” and for greater certainty, “40 students” were considered for each drug.

$$n = \frac{(Z_{1-\alpha/2} + Z_{1-\beta})^2 * [P_1(1-P_1) + P_2(1-P_2)]}{(P_1 - P_2)^2}$$

Where “ $\alpha = 0.05$ ”, “ $\beta = 0.2$ ”, “ $P_1 = 100\%$ ” and “ $P_2 = 45.0\%$ ”.

Treatment

The eighty infected girls were detected and randomized into two groups receiving any of the treatments listed. The amount of permethrin 1% shampoo and dimethicone 4.0% lotion was 60 ml. Affected students and their parents learned proper drug-use skills. If failure occurs at the end of the first week, treatment is continued for an extra week. By the end of week two, the efficacy of the drugs was determined. The Chi-square test was used to compare the drug efficacy. $P < 0.05$ was considered statistically significant.

Table 1. The efficacy of dimethicone and permethrin for the treatment of head lice infestation in girl students.

Treatment groups	Infected students Individuals	Improved Individuals	%	Unimproved Individuals	%
Dimethicone					
At the end of the first week	40	34	85.0	6	15.0
At the end of the second week	6	4	66.7	2	33.3
At the end of two weeks	46	38	82.6	8	17.4
Permethrin					
At the end of the first week	40	27	67.5	13	32.5
At the end of the second week	13	2	15.4	11	84.6
At the end of two weeks	53	29	54.7	24	45.3

Results

Table 1 illustrates the efficacy of the treatment groups. The efficacy of dimethicone and permethrin after treatment were 82.6 and 54.7%, respectively, for removing head lice at the end of two weeks. In other words, the efficacy of dimethicone was 27.9% higher than permethrin (Table 1). At the end of the first week, there were no significant differences between the treatment groups ($P = 0.065$). The outcomes of dimethicone treatment at the end of the second week ($P = 0.025$) and generally at the end of two weeks ($P = 0.006$) were more effective than permethrin.

Discussion

Pediculus capitis is mentioned as one of the most important human obligatory ectoparasites among arthropod-pests of medical importance, such as ticks, mosquitoes and cockroaches or arthropod-associated diseases like leishmaniosis, myiasis and scabies.⁽²²⁻³³⁾ In recent decades, there has been a significant increase in head lice infestations around the world. One of the reasons is drug resistance to consumed drugs.^(34, 35) The efficacy of permethrin was 27.9% lower than that of dimethicone. In other words, for treatment failure, permethrin was found to be 27.9% greater than dimethicone. For students who continued treatment with permethrin for 14 days, the recovery rate (15.4%) was not the same as the non-recovery rate (84.6%) in the second week, and this issue can emphasize the development of drug resistance to permethrin (Table 1). In a study conducted by Rafinejad J, *et al.* treatment failure for permethrin was 20.5% among female students, after seven days⁽³⁶⁾, which is less than the present study.

However, in a study conducted by Soleimani-Ahmadi, treatment failure for permethrin was 49.0% among female students after seven days and diminished to 21.0% after 14 days.⁽³⁷⁾ Treatment outcomes during the first week did not reveal any significant differences between treatment groups. Dimethicone treatment within the second week and generally for two weeks was more effective at removing head lice than permethrin. Zahirnia A, *et al.* reported that the rate of treatment for permethrin was 88.0%.⁽³⁸⁾ In a study conducted by Izri A, *et al.*, the lice killing effect of dimethicone was determined to be 83.2%.⁽³⁹⁾ However, in another study conducted by Ihde ES, *et al.* the results showed that the efficacy of dimethicone and permethrin were 80.7% and 45.3%, respectively, after 14 days.⁽⁴⁰⁾ Tashakori G, *et al.* reported the rate of treatment with permethrin 45.3% after four weeks.⁽⁴¹⁾ One reason for diminishing the therapeutic effect of permethrin is the resistance of *P. capitis* to permethrin in comparison to dimethicone. It appears that the intensity of resistance of *P. capitis* to permethrin has progressively increased over the past decade. The dimethicone mechanism is oil-based, which physically kills the insects by blocking the breathing holes *P. capitis*. Consequently, there is less potential for resistance to dimethicone.⁽²⁰⁾ The difference between dimethicone and permethrin treatment was clearly observed by Burgess IF, *et al.* who suggested the removal of permethrin from the treatment of head lice.⁽⁴²⁾ The results of this study like one conducted by Kalari H, *et al.* confirm the usefulness of dimethicone for the treatment of head lice compared to permethrin.⁽⁴³⁾

Conclusion

The results of this study suggest that dimethicone is useful in the treatment of head lice compared to permethrin.

Acknowledgements

This study forms part of Somayeh Barik-Abi M.Sc. thesis in Medical entomology and vector control in the Department of Medical Entomology, School of Medicine, which was supported financially by the Research Deputy of Hamadan University of Medical Sciences (Grant No: 9710186255). This study was also approved by the Ethical Committee of Hamadan University of Medical Sciences (IR.UMSHA.REC.1397.578).

Conflicts of interest statement

Each of the authors has completed an ICMJE disclosure form. None of the authors declare any potential or actual relationship, activity, or interest related to the content of this article.

Data sharing statement

The present review is based on the references cited. Further details, opinions, and interpretation are available from the corresponding authors on reasonable request.

References

- Nasirian H, Vazirianzadeh B, Taghi Sadeghi SM, Nazmara S. *Culiseta subochrea* as a bioindicator of metal contamination in Shadegan International Wetland, Iran (Diptera: Culicidae). *J Insect Sci* 2014;14:258.
- Nasirian H. Contamination of cockroaches (Insecta: Blattaria) to medically fungi: A systematic review and meta-analysis. *J Mycol Med* 2017; 27:427-48.
- Davari B, Alam FN, Nasirian H, Nazari M, Abdigoudarzi M, Salehzadeh A. Seasonal distribution and faunistic of ticks in the Alashtar county (Lorestan Province), Iran. *Pan Afr Med J* 2017;27:284.
- Nasirian H. Infestation of cockroaches (Insecta: Blattaria) in the human dwelling environments: A systematic review and meta-analysis. *Acta Trop* 2017;167:86-98.
- Nasirian H. Crimean-Congo hemorrhagic fever (CCHF) seroprevalence: A systematic review and meta-analysis. *Acta Trop* 2019;196:102-20.
- Nasirian H. Recent cockroach bacterial contamination trend in the human dwelling environments: A systematic review and meta-analysis. *Bangladesh J Med Sci* 2019;18:540-5.
- Nasirian H. Contamination of cockroaches (Insecta: Blattaria) by medically important bacteria: A systematic review and meta-analysis. *J Med Entomol* 2019;56:1534-54.
- Nasirian H, Salehzadeh A. Control of cockroaches (Blattaria) in sewers: A practical approach systematic review. *J Med Entomol* 2019;56:181-91.
- Nasirian H. New aspects about Crimean-Congo hemorrhagic fever (CCHF) cases and associated fatality trends: A global systematic review and meta-analysis. *Comp Immunol Microbiol Infect Dis* 2020;69:101429.
- Kassiri H, Nasirian H. New insights about human tick infestation features: A systematic review and meta-analysis. *Environ Sci Pollut Res Int* 2021; 28:17000-28.
- Salavati B, Zahirnia AH, Nasirian H, Azari-Hamidian S. Trend of mosquito (Diptera: Culicidae) monthly distribution in Sanandaj County of Iran. *J Biol Diver* 2021;22 11:4705-15.
- Nasirian H, Saghafipour A. Efficacy of several insecticide formulations against *Periplaneta americana* (L.) (Blattaria: Blattidae) in sewers. *Bangladesh J Med Sci* 2021;20:569-85.
- Nasirian H. Ticks infected with Crimean-Congo hemorrhagic fever virus (CCHFV): A decision approach systematic review and meta-analysis regarding their role as vectors. *Travel Med Infect Dis* 2022;47:102309.
- Hajiloie T, Zahirnia A H, Nasirian H, Davari B. Prevalence of head lice infestation and its associated factors among female primary school students in Gahavand County and compared with the previous studies. *Qom Univ Med Sci J* 2022;15:684-95.
- Zahirnia A, Aminpoor MA, Nasirian H. The impact and trend of factors affecting the prevalence of head lice (*Pediculus capitis*) infestation in primary school students. *Chula Med J* 2021;65: 359-68.
- Eroglu F, Basaran Ü, Kürklü CG, Yüceer M, Yalcıntürk RG, Tanrıverdi M, et al. Pediculosis capitis is a growing neglected infestation due to migration in southeast Turkey. *Parasitol Res* 2016;115:2397-401.
- Barik-Abi S, Davari B, Nasirian H. Investigation of factors affecting head lice (Pediculosis Humanus Capitis) infestation prevalence and evaluation of permethrin 1% and dimethicone 4% for its treatment in primary girl students in Doroud County. *J Zabol Med Sch* 2022;5:15-23.
- Tappeh KH, Chavshin A, Hajipirloo HM, Khashaveh S, Hanifian H, Bozorgomid A, et al. Pediculosis capitis among primary school children and related risk factors in Urmia, the main city of west Azarbaijan, Iran. *J Arthropod Borne Dis* 2012;6:79-85.

19. Durand R, Bouvresse S, Berdjane Z, Izri A, Chosidow O, Clark J. Insecticide resistance in head lice: clinical, parasitological and genetic aspects. *Clin Microbiol Infect* 2012;18:338-44.
20. Burgess IF. The mode of action of dimeticone 4% lotion against head lice, *Pediculus capitis*. *BMC Pharmacol* 2009;9:3.
21. Devore C D, Schutze G E, Council on school health, Committee on Infections Diseases. Head lice. *Pediatrics* 2015;135:e1355-e65.
22. Nasirian H. Detailed new insights about tick infestations in domestic ruminant groups: A global systematic review and meta-analysis. *J Parasitic Dis* 2022;46:526-601.
23. Nasirian H, Ladonni H. Artificial bloodfeeding of *Anopheles stephensi* on a membrane apparatus with human whole blood. *J Am Mosq Control Assoc* 2006;22:54-6.
24. Nasirian H, Ladonni H, Poudat A. Mass rearing of *Anopheles stephensi* on human blood by artificial feeding under laboratory conditions. *Bimonthly J Hormozgan Univ Med Sci* 2008; 12:137-42.
25. Nasirian H. An overview of German cockroach, *Blattella germanica*, studies conducted in Iran. *Pak J Biol Sci* 2010;13:1077-84.
26. Nasirian H. New aspects about *Supella longipalpa* (Blattaria: Blattellidae). *Asian Pac J Trop Biomed* 2016;6:1065-75.
27. Davari B, Kashani S, Nasirian H, Nazari M, Salehzadeh A. The Efficacy of MaxForce and Avion gel baits containing fipronil, clothianidin and indoxacarb against the German cockroach (*Blattella germanica*). *Entomol Res* 2018;48:459-65.
28. Nasirian H, Salehzadeh A. Effect of seasonality on the population density of wetland aquatic insects: A case study of the Hawr Al Azim and Shadegan wetlands, Iran. *Vet World* 2019;12:584-92.
29. Kakeh-Khani A, Nazari M, Nasirian H. Insecticide resistance studies on German cockroach (*Blattella germanica*) strains to malathion, propoxur and lambda-cyhalothrin. *Chula Med J* 2020;64:357-65.
30. Salehzadeh A, Darvish Z, Davari B, Nasirian H. The efficacy of baits containing abamectin, dinotefuran, imidacloprid and pyriproxyfen+ abamectin against *Blattella germanica* (L.) (Blattaria: Blattellidae), the German cockroach. *Afr Entomol* 2020;28:225-37.
31. Nasirian H, Zahirnia A. Detailed infestation spectrums about biological stages of hard ticks (Acari: Ixodida: Ixodidae) in humans: A systematic review and meta-analysis. *Acta Parasitol* 2021; 66:770-96.
32. Tavakoli MM, Davari B, Nasirian H, Salehzadeh A, Moradkhani S, Zahirnia AH. Investigation of insecticidal properties of *Rosmarinus officinalis* and *Lavandula angustifolia* essential oils against German cockroach in laboratory. *J Kashan Univ Med Sci* 2021;25:994-1002.
33. Nasirian H. Reply to comment on “ticks infected with Crimean-Congo hemorrhagic fever virus (CCHFV): A decision approach systematic review and meta-analysis regarding their role as vectors”. *Travel Med Infect Dis* 2023;51:102420.
34. Bohl B, Evetts J, McClain K, Rosenauer A, Stellitano E. Clinical practice update: pediculosis capitis. *Pediatr Nurs* 2015;41:227-34.
35. Koch E, Clark JM, Cohen B, Meinking TL, Ryan WG, Stevenson A, et al. Management of head louse infestations in the United States- a literature review. *Pediatr Dermatol* 2016;33:466-72.
36. Rafinejad J, Nourollahi A, Biglarian A, Javadian E, Kazemnejad A, Doosti S. The comparison of the effect of permethrin shampoo and lindane lotion on the treatment of head lice (*Pediculus humanus capitis*) in the primary school pupils. *J Maz Univ Med Sci* 2011;21:35-41.
37. Soleimani-Ahmadi M, Jaberhashemi SA, Zare M, Sanei-Dehkordi A. Prevalence of head lice infestation and pediculicidal effect of permethrine shampoo in primary school girls in a low-income area in southeast of Iran. *BMC Dermatol* 2017;17:1-6.
38. Zahirnia AH, Taherkhani H, Bathaie SJ. A comparative study on the effectiveness of three different shampoos in treatment of head lice (*Pediculus capitis*) in primary school-children in Hamadan province, IRAN 2000-2001. *J Maz Univ Med Sci* 2005;15:16-24.
39. Izri A, Uzzan B, Maigret M, Gordon MS, Bouges-Michel C. Clinical efficacy and safety in head lice infection by *Pediculus humanis capitis* De Geer (Anoplura: Pediculidae) of a capillary spray containing a silicon-oil complex. *Parasite* 2010;17: 329-35.
40. Ihde ES, Boscamp JR, Loh JM, Rosen L. Safety and efficacy of a 100% dimethicone pediculocide in school-age children. *BMC Pediatr* 2015;15:70.
41. Tashakori G, Dayer MS, Mashayekhi-Ghoyonlo V. Comparative efficacy of three control protocols of head lice (*Pediculus humanus capitis*) infesting schoolchildren in Mashhad city, Iran. *Int J Pediatr* 2018;6:7803-14.
42. Burgess IF, Brunton ER, Burgess NA. Single application of 4% dimeticone liquid gel versus two applications of 1% permethrin creme rinse for treatment of head louse infestation: A randomised controlled trial. *BMC Dermatol* 2013;13:5.
43. Kalari H, Soltani A, Azizi K, Faramarzi H, Moemenbellah-Fard MD. Comparative efficacy of three pediculicides to treat head lice infestation in primary school girls: A randomised controlled assessor blind trial in rural Iran. *BMC Dermatol* 2019;19:1-9.