

Spray-dried powder of *Bacillus amyloliquefaciens* strain C2-1 for control of rice diseases

Usmana Meehae¹, Jutarut Iewkittayakorn^{1*}, Aunkamol Kumngen¹, Wilaiwan Chotigeat¹, Sudarat Suwannarat¹, Sukhuman Whankaew² and Siwapong Leunram³

¹ Division of Biological Science, Faculty of Science, Prince of Songkla University, Hat Yai, Songkhla 90110, Thailand

² Department of Plant Science, Faculty of Technology and Community Development, Thaksin University, Phatthalung 93210, Thailand

³ BIOAXEL Co., Ltd. 6/9 Moo 3 Bophut Chaweng Beach Road, Koh Sa-mui, Surat Thani 84320, Thailand

*Corresponding author: jutarut.p@psu.ac.th

Received: December 10, 2022. Revised: March 7, 2023. Accepted: March 14, 2023.

ABSTRACT

Rice cultivation in Thailand faces challenges in achieving high yields due to diseases. This research aims to develop a spray-dried powder containing *Bacillus amyloliquefaciens* strain C2-1 for control economically damaging rice diseases caused by *Xanthomonas oryzae*, *Fusarium moniliforme*, *Pyricularia grisea*, and *Bipolaris oryzae*. The *B. amyloliquefaciens* strain C2-1 was cultured in 3 different media, including 1) medium contained nutrient broth with glucose (NB), 2) medium contained beef extract, molasses, K₂HPO₄ and KH₂PO₄ (BM), and 3) medium contained yeast extract, molasses, K₂HPO₄, and KH₂PO₄ (YM). The results showed that YM medium gave the highest concentration of bacteria at 1.30x10¹⁶ cfu/ml, followed by BM medium at 9.88x10¹² cfu/ml, and NB medium at 5.3x10¹²cfu/ml. Therefore, YM medium was selected as the optimal culture medium. Subsequently, the *B. amyloliquefaciens* strain C2-1 was produced into a spray-dried powder using the spray-drying technique. Tapioca starch and carboxymethyl cellulose were used as spray-drying carriers in a ratio of 10:1. The optimal spray-drying conditions included a hot air inlet temperature of 90°C, a hot air outlet temperature of 70°C, and a feed rate of 10-15 ml/min, resulting in the highest survival rate at 2.8x10¹² CFU/g⁻¹. The efficacy of *B. amyloliquefacien* strain C2-1 powder was then tested against rice diseases. The inhibition of *X. oryzae* was assessed using the paper disc diffusion method, while the inhibitions of *F. moniliforme*, *P. grisea*, and *B. oryzae* were evaluated using dual culture assay. The results demonstrated that *B.amyloliquefacien* strain C2-1) powder exhibited strong inhibitory effects against *X. oryzae*, with clear zone measuring 5.5, 5.7, and 5.7 mm at 24, 48, and 72 h, respectively. Regarding inhibiting *F. moniliforme*, *P. grisea*, and *B. oryzae*, *B. amyloliquefaciens* strain C2-1 demonstrated effective control of these rice pathogenic fungi, resulting in inhibitions of 73.0%, 68.72%, and 48.42% at 7 days, respectively. These findings strongly suggest that the *B.amyloliquefacien* strain C2-1 obtained from spray drying serves as a promising biocontrol agent against bacterial leaf blight disease, bacterial blight, bakanae disease, rice blast disease, and brown spot disease caused by *F. moniliforme*, *P. grisea*, and *B. oryzae*, respectively.

Keywords: Antagonist bacteria, *B.amyloliquefacien*, biological control, rice disease, spray drying

INTRODUCTION

Rice (*Oryza sativa* L.) is the crucial main food crop belonging to the family Poaceae and is widely cultivated in most tropical and subtropical regions (Ezuka and Kaku, 2000). Nowadays, rice cultivation in Thailand still faces problems that affect the growth and yield of rice caused by weeds, soil, water, and environmental changes. One primary disease of rice is the bacterial leaf blight disease caused by *Xanthomonas oryzae* which can infect rice at any growth stage causing production losses depending on climate, growing season, rice variety, and growth; if the disease is severe, the yield can be reduced by 50 percent (Chintaganon et al., 2022). *Pyricularia grisea* is a fungus that causes blast disease that damages rice. Rice blast disease is

favored by a number of factors such as high relative humidity (>80 percent), low temperature (15°C - 26°C), cloudy weather, more wet or rainy days, and excessive doses of nitrogen fertilizers (Saleh et al., 2020). *Bipolaris oryzae* causes brown spot disease that affects both quantity and quality of rice grains. This pathogen, under favorable epidemic conditions, can cause reduction in grain yield of up to 90 percent (Dorneles et al., 2020).

Strains of *B. amyloliquefaciens* strain C2-1 have been reported as a potential biological agent for controlling other crop diseases (Shrestha et al., 2016). This is accompanied by Srivastava et al. (2016), who reported that *B. amyloliquefaciens* (SN13) acts as a biocontrol agent and enhance the immune response against *R. solani*,

a necrotrophic fungus causing sheath blight in rice, by modulating various physiological, metabolic, and molecular functions. Moreover, Prabhukarthikeyan et al. (2019) also reported that *Bacillus amyloliquefaciens* strain BS5 effectively reduced the brown spot disease in rice under *in vitro*, glasshouse, and field conditions.

The spray-drying technique has been routinely used for the preservation and concentration of microorganisms due to its gentle protocols of drying by exposing substances to only a short burst of extreme temperature, then providing a cooling effect via the evaporation process, which is able to maintain a high survival rate of microorganisms and low production cost (Desmond et al., 2001). In order to prolong shelf life, and increase efficacy, it is important to keep the bacterium in a dormant state, while maintaining its viability. The efficacy of bio-product was tested in a greenhouse. The result showed that the disease incidence was reduced by 100 and 61.54% when treated the rhizome with bio-products MTR13 and PS6, respectively, before planting and application of their suspension was done one hour after the pathogen inoculation when compared with the control. (Thano and Akarapisan, 2018). The aim of this study was to develop bio-based products based on *B. amyloliquefaciens* strain C2-1 under optimal spray drying conditions for controlling rice disease. Therefore, the information on efficacy testing may be useful and able to apply or recommended to farmers or those who are interested in controlling diseases in rice fields.

MATERIALS AND METHODS

Microorganism and inoculum preparation

The antagonist bacterium *B. amyloliquefacien* strain C2-1 was isolated from food waste compost obtained from the Center for Genomics and Bioinformatics Research, Faculty of Science, Prince of Songkla University, Songkhla, Thailand. The rice pathogen, namely *X. oryzae*, *F. moniliforme*, *P. grisea*, *B. oryzae*. The mass losses may be attributed to the effect of the different parameters of spray drying, such as inlet temperature and airflow, as well as to the low Tg of carrying agents obtained from Phattalung Rice Research Center, Phattalung, Thailand. The *B. amyloliquefaciens* strain C2-1 and *X. oryzae* were cultured in nutrient broth (NB) (nutrient broth 13 g/L; glucose 10 g/L) and glucose broth medium (NGB) (13 g of nutrient broth and 12.5 g of glucose anhydrous per 1 L of distilled water), respectively at 37°C for 24 h on 200 rpm of the rotary shaker to prepare the inoculum. The *F. moniliforme*, *P. grisea*,

B. oryzae were grown on potato dextrose agar (PDA) at 28°C for 7 days to prepare the inoculum.

Substrates and chemicals

All chemicals were of analytical grade, including nutrient broth (NB) (Himedia, Nashik, India) and KH₂PO₄ (anhydrous, Loba Chemie, Mumbai, India). Yeast extract powder was obtained from TM Mida (Rajasthan, India), beef extract from Srichem (Mumbai, India); KH₂PO₄ was obtained from Cloisters (Cherrybrook, Australia); glucose was purchased from Utopian Co., Ltd., Samuthprakarn, Thailand; and molasses was purchased from local markets (Songkla, Thailand).

Effect of culture media on antagonist bacterium growth

The antagonist bacterium *B. amyloliquefaciens* strain C2-1 was cultured in different culture media, including nutrient broth (NB), beef extract molasses broth (BM) (beef extract 3 g/L; molasses 20 g/L; K₂HPO₄ 0.05 g/L; KH₂PO₄ 0.5 g/L) and yeast extract molasses broth (YM) (Yeast extract 0.5 g/L; Molasses 20 g/L; K₂HPO₄ 0.05 g/L; KH₂PO₄ 0.5 g/L) were prepared in each 250 mL flask with 100 mL medium containing. Each flask was cultured at 37°C for 24 h on 200 rpm of the rotary shaker before checking the colony forming unit (CFU/mL) using the serial dilution method on the NA agar medium. This experiment was maintained with three replications.

Preparation of antagonist bacterium in the spray-dried powder using spray-drying technique

The antagonist bacterium was cultured in YM, NB, and BM at 37 °C for 24 h on 200 rpm of the rotary shaker. The cell suspension was adjusted to 0.5 optical density (O.D. 600 nm) using the spectrophotometer (Thermo Fisher Scientific, USA) before mixing with tapioca starch and carboxymethyl cellulose (CMC). The mixing solution consisted of 30 mL cell suspension, 18.18 g tapioca starch, and 1.82 g CMC (Teera-Arunsiri et al., 2003) Spray drying was carried out in a Mini Spray Dryer B290 (BÜCHI, Labortechnik AG, Flawil, Switzerland) at the following operation conditions: feed pump of 5 ml/min °C, aspiration of 100% and pressure of 1.5 bar. The inlet and outlet temperatures were set as 90 and 70 °C, respectively. The obtained spray-dried powder was calculated as a percentage yield as following Eq. (1) that those reported by (Chumthong et al., 2016). Three replicates were performed for each formulation.

$$\% \text{ Yield} = [\text{outlet weight}/\text{inlet weight}] \times 100 \quad (1)$$

Viable bacterial counts in the formulations were tested against the bacterium before drying using the drop-on plate method (Zuberer, 1994) with bef. 0.1g/mL suspension of spray-dried powder was prepared in sterile distilled water and was serially diluted from 1×10^{-1} to 1×10^{-12} . The viable bacteria were cultured on NA medium at 37 °C for 24 h, after which the CFU was counted. The number of colony-forming unit per gram (CFU/g) was the average of six drops per dilution. The cell concentrations exhibited between 30 and 300 CFU (1×10^{-4} and 1×10^{-5}) were selected (Daniel et al., 2020). Equation (2) was employed for the quantification of culturability. All experiments were conducted in triplicate, and the reported values represented the average of the calculated values. Formulations with the concentrations of highest were chosen for further studies.

$$\text{CFU/g} = \frac{[\text{N}^\circ \text{plate colonies} \times \text{dilution factor}]}{\text{mL sample seeded}} \quad (2)$$

When N° is the initial population of *B. amyloliquefaciens* strain C2-1 (cfumL⁻¹)

Antagonistic activities of *B. amyloliquefaciens* strain C2-1 for pathogenic bacteria inhibition assays

Rice pathogenic bacteria, *X. oryzae* was cultivated in an NGB medium. 100 mL working volume of the NGB medium was conducted in each 250 ml Erlenmeyer flask for cultivation. The optical density of cell suspension was measured at 600 nm (OD600) with a UV-vis spectrophotometer (Thermo Fisher Scientific, USA). The culture flasks were inoculated with 10% (v/v) inoculums (initial cell suspension equal to 0.2 at OD600) at 25°C for 24 h on 200 rpm of the rotary shaker. The inhibition assays of spray-dried powder of *B. amyloliquefaciens* strain C2-1 testing were done using the agar well diffusion method (Yang et al., 2012). Glucose agar medium was prepared by pouring 10 mL glucose agar (NGA) medium into a petri dish, and let it set in a laminar flow hood as a basal layer. After that, use a cork borer, a diameter of 5 mm, drill 5 holes, and then using a cotton swab sterile dip the pathogens that have been prepared and make a swab over the entire surface of the culture medium. In this method, the agar plate was inoculated with a powder of *B. amyloliquefaciens* strain C2-1- dissolved in water by dropping each hole 10 ul of and placed on the four corners of the seed layer. At the same time, 10 uL of sterile water was dropped into one hole in the center of the same petri dish (control). The petri dishes were then incubated at 37 °C for 24, 48, 72 h, and 1 week, and then measured the diameters of inhibition growth zones. The inhibition growth zones can be calculated from Equation (3).

$$R_a = (D_c - D_s) / 2 \quad (3)$$

where R_a is the inhibition radius (mm); D_c is the diameter of the clear area (mm), and D_s is the diameter of the specimen (mm)

Antagonistic activities of *B. amyloliquefaciens* strain C2-1 for pathogen fungi inhibition assays

Three fungal pathogens, *F. moniliforme*, *P. grisea*, and *B. oryzae* were cultured on PDA at 28°C for 7 days to test the inhibition of pathogen mycelial growth. The pathogen fungi inhibition assays of spray-dried powder of *B. amyloliquefaciens* strain C2-1 spray-dried was dissolved in 1 mL of sterile water. Each treatment was conducted in 3 replications. Mycelial inhibition of pathogen fungi was assessed as a percentage of mycelial inhibition at 3, 5, and 7 days after culturing pathogen fungi. The inhibition growth zones can be calculated from Equation (4) (Morton and Stroube, 1995).

$$\begin{aligned} \text{Percentage of pathogen mycelial inhibition} \\ = [(R_1 - R_2) / R_1] \times 100 \end{aligned} \quad (4)$$

Where R_1 is the colony radius of pathogen fungi on PDA incorporated powder *B. amyloliquefaciens* strain C2-1 solution of bacterial formulation, and R_2 is the colony radius of pathogen fungi control on PDA.

RESULTS AND DISCUSSION

Cultural media basic suitable for growth

The antagonist bacterium *B. amyloliquefaciens* strain C2-1 grown in YM medium performs the highest cell viability, followed by BM and NB, respectively (Table 1). As a result, YM and BM media contained a higher nutrient composition than the NB medium. It also can be seen that YM used molasses as a major ingredient which is a cheap carbon source, and yeast extract is a good source of nitrogen for growth. The buffered K_2HPO_4 and KH_2PO_4 can also help the infection grow better. It may help to adjust the pH condition so that it does not change according to the metabolites produced by the germ during growth.

Table 1. Viability of *B. amyloliquefaciens* C2-1 grown in different media

Formulation	cfu/g*		
	1 st	2 nd	3 rd
NB	1.8±0.0 x 10 ¹²	2.8±0.1 x 10 ¹²	2.6±0.0 x 10 ¹²
BM	3.6±0.1 x 10 ¹²	3.8±0.2 x 10 ¹²	5.5±0.2 x 10 ¹²
YM	4.9± 0.0 x 10 ¹⁵	5.2± 0.0 x 10 ¹⁵	4.7±0.1 x 10 ¹⁵

Antagonist bacterium in the spray-dried powder using spray-drying

The initial amounts of powder dry matter of *B. amyloliquefaciens* strain C2-1 that grew in all three media after spray drying at a constant 90 °C inlet temperature were similar. From the initial amount of dry matter, 20 grams, after drying, the remaining 13 grams, thus the calculated average yield was 65%. The mass losses may be attributed to the effect of the different parameters of spray drying, such as inlet temperature and airflow, as well as to the low Tg of carrying agents (León-Martínez et al., 2010). Although the spray-dried product of *Bacillus* sp., especially *B. subtilis* (Yáñez et al., 2012), has been devised and tested to control insect pests and plant disease, in this study, a spray-dried product of *B. amyloliquefaciens* strain C2-1 had been formulated and tested for the control of rice disease in the laboratory. The characteristic of spray-dried powder can be seen in Fig.1. The powder is fine and slightly yellowish-white. The viability of spray-dried powder of *B. amyloliquefaciens* strain C2-1 grown in different media is shown in Table 2. The result found

that *B. amyloliquefaciens* strain C2-1 grown in YM medium performed the highest cell viability followed by BM and NB, respectively, which gave the same experiment result as the effect of culture media on antagonist bacterium growth. Therefore, the spray-dry powder of *B. amyloliquefaciens* strain C2-1 with YM medium was further studied for the pathogen inhibition assays.

**Figure 1.** Characteristic of powder of *B. amyloliquefaciens* strain C2-1 after spray drying at constant 90 °C inlet temperature.**Table 2.** Viability of spray-dried powder of *B. amyloliquefaciens* strain C2-1 grown in different media

Formulation	cfu/g*		
	1 st	2 nd	3 rd
NB	3.4±0.0 x 10 ⁸	1.8±0.1 x 10 ⁷	1.7±0.1 x 10 ⁷
BM	4.2±0.2 x 10 ¹⁰	5.7±0.1 x 10 ¹⁰	5.8±0.0 x 10 ¹⁰
YM	2.6±0.1 x 10 ¹²	3.4±0.1 x 10 ¹²	2.8±0.1 x 10 ¹²

Biocontrol activities of antagonistic *B. amyloliquefacien* strain C2-1

The test results of the efficacy in inhibiting the rice pathogen *X. oryzae* by the spray-dried powder of *B. amyloliquefaciens* strain C2-1 can be seen in Figure 2. The inhibition zone showed that the spray-dried powder was still effective against *X.*

oryzae rice pathogens at 24, having clear zones of 1.65. This result shows that *B. amyloliquefaciens* strain C2-1 gave the inhibition of the causative agent of rice leaf blight. This inhibition may be caused by volatile substances from *B. amyloliquefaciens* strain C2-1, according to Wu et al. (2015), who reported that *B. amyloliquefacien* FZB42 had antibacterial activity against *Xoo* rice pathogens.




Figure 2. Reaction of the *X. oryzae* in agar well diffusion assay after inoculation for 24 hr. at 37 °C temperature, when inhibited by Powder *B. amyloliquefaciens* strain C2-1.

The spray-dried powder of *B. amyloliquefaciens* strain C2-1 showed antifungal activity against all 3 rice pathogenic fungi in dual culture tests incubated for 7 days. The spray-dried powder showed the strongest inhibition, as seen in Fig. 3, mycelial growth was reduced compared to control plate 3, 5, and 7 days. The inhibition percentages *F. moniliforme* were 0.78, 52.51, and 73.0%, respectively. For *B. oryzae*, the inhibition percentages by powder *B. amyloliquefaciens* strain C2-1 were 0.65, 22.86, and 48.42%, respectively. The mycelial growth of the pathogen *P. grisea*, and the inhibition levels by powder *B. amyloliquefaciens* strain C2-1 were 0.76, 56.99, and 68.72 %, respectively. Thus, powder *B. amyloliquefaciens* strain C2-1 had comparatively high effectiveness in suppressing the pathogens, after incubation for 3, 5, and 7 days.




Figure 3. Effects of spray-dried powder of *B. amyloliquefaciens* strain C2-1 against *F. moniliforme*, *P. grisea*, and *B. oryzae* on dual culture plates incubated for 3, 5, and 7 days at room temperature.

CONCLUSIONS

In this study, *B. amyloliquefacien* strain C2-1 isolated from food waste compost, effectively inhibiting the growth of *X. oryzae* as well as the mycelial growth of *F. moniliforme*, *P. grisea*, and *B. oryzae* strain. Among the three general media tested, YM was selected as the optimum culture media for cell growth and viability after spray drying due to its cost-effectiveness and highest cell concentration. The inhibition of *Xoo B. amyloliquefaciens* strain C2-1 powder against *X. oryzae* gave the widest clear zones. Regarding inhibiting *F. moniliforme*, *P. grisea*, and *B. oryzae*, it was found that *B. amyloliquefacien* strain C2-1 was able to control the growth of all these pathogenic rice fungi with 73.0%, 68.72 and 48.42 % inhibitions, at 7 days respectively. Thus, our results highlight the potential of spray-dried powder of *B. amyloliquefaciens* strain C2-1 as a promising biocontrol agent against the prevalent rice diseases currently affecting crops.

ACKNOWLEDGMENTS

This research was supported by Higher Education for Industry Consortium (Hi-Fi) under the Experiential Learning Program, Office of The Permanent Secretary (OPS), Ministry of Higher Education, Science, Research and Innovation. The authors would like to thank the Division of Biological Science and the Center for Genomics and Bioinformatics Research, Faculty of Science, Prince of Songkla University (PSU), Hat Yai, Songkhla, Thailand.

REFERENCES

- Chintaganom, T., Watcharachaiyakup, J., Wonnapijiti, P., Tongyoo and Patarapuwadol, S. 2022. Draft genome sequence and population structure of *Xanthomonas oryzae* pv. *oryzae* causing bacterial blight disease on rice in Thailand. Thai Agriculture Research Journal. 40(1). <https://doi.org/10.14456/thaidoa-agres.2022.4>
- Chumthong, A. wiwattanapatapee, R. viernstein, H. pengnoo, A. and kanjanamaneesathian, M. 2016. Spray-dried powder of *Bacillus megaterium* for control of rice sheath blight disease: Formulation protocol and efficacy testing in laboratory and greenhouse. Cereal Research Communications. 44(1): 131–140. <https://doi.org/10.1556/0806.43.2015.034>.
- Desmond, C., Stanton, C., Fitzgerald, F.G., Collins, K., Ross, P.R. 2001. Environmental adaptation of probiotic lactobacilli towards improvement of performance during spray drying. International Dairy Journal. 11(10): 801-808. [https://doi.org/10.1016/S0958-6946\(01\)00121-2](https://doi.org/10.1016/S0958-6946(01)00121-2).
- Daniel, V.M., Vicente, E.S., César, L.P., Patricia, A.B., Fidel, M.G., Manuel, R.A. and María, Z.S.L. 2020. Preparation of spray-dried functional food: Effect of adding *Bacillus clausii* bacteria as a co-microencapsulating agent on the conservation of Resveratrol. Processes. 8(7): 849. <https://doi.org/10.3390/pr8070849>.
- Dorneles, R.K., Refatti, P.J., Pazdiora, C.P., Avila, A.L., Deuner, S. and Dallagnol, J.L. 2020. Biochemical defenses of rice against *Bipolaris oryzae* increase with high atmospheric concentration of CO₂. Physiological and Molecular Plant Pathology. 110: 101484 <https://doi.org/10.1016/j.pmpp.2020.101484>.
- Ezuka, A. and Kaku, H. 2000. A historical review of bacterial blight of rice. Bulletin of the National Institute of Agrobiological Resource. 15:1-207.
- León-Martínez, F.M., Méndez-Lagunas, L.L., Rodríguez-Ramírez, J. 2010. Spray drying of nopal mucilage (*Opuntia ficus-indica*): Effects on powder properties and characterization. Carbohydrate Polymer. 81: 864-870. <https://doi.org/10.1016/j.carbpol.2010.03.061>.
- Morton, D.J., and Stroube, W.H. 1955. Antagonistic and stimulatory effects of soil microorganisms upon *Sclerotium rolfsii*. Phytopathol. 45: 417-420.
- Prabhukarthikeyan, S.R., Yadav, M.K., Anandan, A., Aravindan, S., Keerthana, U., Raghu, S., Baite, M.S., Parameswaran, C., Panneerselvam, P., and Rath, P.C. 2019. Bio-protection of brown spot disease of rice and insight into the molecular basis of interaction between *Oryza sativa*, *Bipolaris oryzae* and *Bacillus amyloliquefaciens*, Biological Control. 137: 104018. <https://doi.org/10.1016/j.biocontrol.2019.104018>.
- Saleh, A.S., Imtiaz, A.A., Hossian Md.B., Husna, A., and khatun Eaty, Most.N. 2020. Review: Rice blast disease. Annual Research & Review in Biology. 35(1): 50-64 <https://doi.org/10.9734/ARRB/2020/v35i130180>.
- Shrestha, B.K., Karki, H.S., Groth, D.E., Jungkhun, N., and Ham, J.H. 2016. Biological control activities of rice-associated *Bacillus* sp. strains against sheath blight and bacterial panicle blight of rice. PLoS One. 11(1): e0146764. <https://doi.org/10.1371/journal.pone.0146764>.
- Srivastava, S., Bist, V., Srivastava, S., Singh, P.C., Trivedi, P.K., Asif, M.H., Chauhan, P.S., and Nautiyal, C.S. 2016. Unraveling aspects of *Bacillus amyloliquefaciens* mediated enhanced production of rice under biotic stress of *Rhizoctonia solani*. Front. Plant Sci., 06 May 2016. <https://doi.org/10.3389/fpls.2016.00587>.
- Teera-Arumsiri, A., Suphantharika, M., and Ketunuti, U. 2003. Preparation of spray-dried wettable powder formulations of *Bacillus thuringiensis*-based biopesticides. Journal of Economic Entomology. 96: 292–299. <https://doi.org/10.1603/0022-0493-96.2.292>.
- Thano P., and Akarapisan A. 2018. Development of bio-product from *Bacillus amyloliquefaciens* for controlling bacterial wilt of curcuma (*Curcuma lismatifolia*). Agricultural Sci. J. 49: 1 (Suppl.): 650-652.
- Wu, L., Wu, H., Chen, L., Yu, X.F., Borriss, R., and Gao, X. 2015. Difficidin and bacilysin from *Bacillus amyloliquefaciens* FZB42 have antibacterial activity against *Xanthomonas oryzae* rice pathogens. Sci. Rep.5:12975. <https://doi.org/10.1038/srep12975>.
- Yang, E., Fan, L., Jiang, Y., Doucette, C., and Filmore S. 2012. Antimicrobial Activity of Bacteriocin-Producing Lactic Acid Bacteria Isolated from Cheeses and Yogurts. AMB express a Springer Open journal. 2(48).
- Yáñez, V., Vinas, I., Usall, J., and Torres, R. 2012. Formulation development of the biocontrol agent *Bacillus subtilis* strain CPA-8 by spray-drying. Journal of Applied Microbiology 112(5):954-65 <https://doi.org/10.1111/j.1365-2672.2012.05258.x>.
- Zuberer, D.A. 1994. Recovery and enumeration of viable bacteria. In: Weaver, R.W., Angle, S., Bottomley, P., Bezdicek, D., Smith, S., Tabatabai, A., Wollum, A. (eds). Methods of Soil Analysis: Part 2, Microbiological and Biochemical Properties, Number 5. Soil Science Society of America Book Series. Soil Science Soc. of America Inc. Wisconsin. p. 119–143.