



# A Note on Regularity of Certain Semigroups of Transformations Preserving a Reflexive and Transitive Relation

Kitsanachai Sripon<sup>1</sup>, Ekkachai Laysirikul<sup>1,\*</sup>, Yanisa Chaiya<sup>2,3</sup>

<sup>1</sup>Department of Mathematics, Faculty of Science, Naresuan University, Phitsanulok 65000, Thailand

<sup>2</sup>Department of Mathematics and Statistics, Faculty of Science and Technology, Thammasat University, Pathum Thani 12120, Thailand

<sup>3</sup>Thammasat University Research Unit in Algebra and Its Applications, Thammasat University, Pathum Thani 12120, Thailand

Received 23 March 2023; Received in revised form 12 June 2023

Accepted 28 June 2023; Available online 26 September 2023

## ABSTRACT

In this paper, we introduce semigroups of transformations that preserve a reflexive and transitive relation. These semigroups are generalizations of well-known classes of transformation semigroups, namely semigroups of transformations preserving an equivalence relation and semigroups of order-preserving transformations. Additionally, we investigate the necessary and sufficient conditions for elements within such a semigroup, in certain cases, to be regular. These results extend the existing knowledge in the aforementioned semigroups.

**Keywords:** Reflexive relation; Regular element; Transformation semigroup; Transitive relation

## 1. Introduction

Let  $X$  be a nonempty set and  $T(X)$  the set of all functions from  $X$  into  $X$ . Then  $T(X)$  is a semigroup under the operation of composition. It is well-known that each  $\alpha \in T(X)$ ,  $\alpha = \alpha\beta\alpha$  for some  $\beta \in T(X)$ , that is,  $T(X)$  is a regular semigroup (see [1], for details). Since every semigroup can be embedded in  $T(X)$ , for some appropriate set  $X$ , the structural properties on such type

of semigroups have been researched extensively. For extending results, generalized transformation semigroups have been continually constructed.

For an equivalence relation  $E$  on a set  $X$ , let

$$T_E(X) = \{\alpha \in T(X) : \forall x, y \in X, (x, y) \in E \text{ implies } (x\alpha, y\alpha) \in E\}.$$

Then  $T_E(X)$  is a subsemigroup of  $T(X)$ . In

particular, if  $E = \{(x, x) : x \in X\}$  or  $E = X \times X$ , then  $T_E(X) = T(X)$ . However, in general,  $T_E(X)$  is not regular. All regular elements within such semigroups have been completely described in [2]. Several properties of  $T_E(X)$  have been identified, including those in [3–8].

Let  $(X, \leq)$  be a partially ordered set. A map  $\alpha : X \rightarrow X$  is said to be order-preserving if  $x \leq y$  implies  $x\alpha \leq y\alpha$  for all  $x, y \in X$ . We denoted by  $OT(X)$  the semigroup of all order-preserving transformations on a partially ordered set  $X$ . Such a semigroup is a subsemigroup of  $T(X)$  and plays an important role in the study of algebraic systems. Necessary and sufficient conditions for any elements of  $OT(X)$  to be regular were discovered in [9]. Many others properties of  $OT(X)$  have also been explored, see, for example [10–14].

In this paper, we introduce a subsemigroup of  $T(X)$ , which is a generalization of  $T_E(X)$  and  $OT(X)$ , by letting  $\sigma_{RT}$  be a reflexive and transitive relation on  $X$  and

$$T(X, \sigma_{RT}) = \{\alpha \in T(X) : \forall x, y \in X, (x, y) \in \sigma_{RT} \text{ implies } (x\alpha, y\alpha) \in \sigma_{RT}\}.$$

$T(X, \sigma_{RT})$  is called the semigroup of full transformations on  $X$  which preserve  $\sigma_{RT}$ . Particularly,  $T(X, \sigma_{RT}) = T_{\sigma_{RT}}(X)$  whenever  $\sigma_{RT}$  is symmetric. Moreover, if  $(X, \sigma_{RT})$  is a partially ordered set, then  $T(X, \sigma_{RT}) = OT(X)$ .

## 2. Main Results

Here, we present necessary and sufficient conditions for elements in  $T(X, \sigma_{RT})$  to be regular, where  $\sigma_{RT}$  satisfies the condition,  $\forall x, y \in X, (x, y) \in \sigma_{RT}$  or  $(y, x) \in \sigma_{RT}$ . Moreover, we prove that if  $T(X, \sigma_{RT})$  is a regular semigroup, then  $\sigma_{RT} = X \times X$  or  $\sigma_{RT}$  is a totally order relation on  $X$ . Let  $X$  be a nonempty set and  $\sigma_{RT}$  an arbitrary

reflexive and transitive relation on  $X$ . For each  $x \in X$ , let

$$\bar{x} = \{y \in X : (x, y), (y, x) \in \sigma_{RT}\} \text{ and } \overline{X} = \{\bar{x} : x \in X\}.$$

Obviously,  $\bigcup \overline{X} = X$ . Especially, for the case  $\sigma_{RT}$  is an equivalence relation,  $\bar{x}$  is an equivalence class of  $\sigma_{RT}$  containing  $x$  and  $\overline{X}$  is readily a partition of  $X$ . Although  $\sigma_{RT}$  is not an equivalence relation,  $\overline{X}$  is a partition of  $X$ , as shown in the following lemma.

**Lemma 2.1.**  $\overline{X}$  is a partition of  $X$ .

*Proof.* Let  $\bar{x}, \bar{y} \in \overline{X}$  be such that  $\bar{x} \cap \bar{y} \neq \emptyset$ . Then there exists  $z \in \bar{x} \cap \bar{y}$ . Hence,  $(x, z), (z, x), (y, z), (z, y) \in \sigma_{RT}$ . By the transitivity of  $\sigma_{RT}$ , we get  $(x, y), (y, x) \in \sigma_{RT}$ . Let  $a \in \bar{x}$ . Then  $(x, a), (a, x) \in \sigma_{RT}$ , which implies  $(a, y), (y, a) \in \sigma_{RT}$ . Thus,  $a \in \bar{y}$  and so  $\bar{x} \subseteq \bar{y}$ . Similarly, we can show that  $\bar{y} \subseteq \bar{x}$ . Therefore,  $\overline{X}$  is a partition of  $X$ .  $\square$

Note, for  $\alpha \in T(X, \sigma_{RT})$ , that  $\bar{x}\alpha \subseteq \overline{x\alpha}$ , since  $(x, z), (z, x) \in \sigma_{RT}$  implies  $(x\alpha, z\alpha), (z\alpha, x\alpha) \in \sigma_{RT}$  for all  $z \in \sigma_{RT}$ . Hereafter, we denote by  $\text{Reg}(T(X, \sigma_{RT}))$  the set of all regular elements of  $T(X, \sigma_{RT})$ .

**Lemma 2.2.** Let  $\alpha \in T(X, \sigma_{RT})$  and  $\bar{x} \in \overline{X}$ , such that  $\bar{x} \cap X\alpha \neq \emptyset$ . If  $\alpha \in \text{Reg}(T(X, \sigma_{RT}))$ , then there exists  $\bar{y} \in \overline{X}$  such that  $\bar{x} \cap X\alpha = \bar{y}\alpha$ .

*Proof.* Assume  $\alpha = \alpha\beta\alpha$  for some  $\beta \in T(X, \sigma_{RT})$ . Choose  $\bar{y} = \overline{x\beta} \in \overline{X}$ . By the above note, we have  $\bar{x}\beta \subseteq \overline{x\beta} = \bar{y}$ . Let  $z \in \bar{x} \cap X\alpha$ . Then there exists  $x' \in X$ , such that  $z = x'\alpha$ .  $z = x'\alpha = x'\alpha\beta\alpha = z\beta\alpha \in \overline{x\beta}\alpha \subseteq \overline{x\beta\alpha} = \bar{y}\alpha$ . This implies  $\bar{x} \cap X\alpha \subseteq \bar{y}\alpha$ . Since  $\overline{X}$  is a partition of  $X$  and  $\bar{x} \cap \bar{y}\alpha \neq \emptyset$ , we have  $\bar{y}\alpha \subseteq \bar{x}$ . Therefore,  $\bar{x} \cap X\alpha = \bar{y}\alpha$ .  $\square$

For each  $\emptyset \neq A \subseteq X$  and  $x \in X$ , let

$$U(A, x) = \{y \in A : (x, y) \in \sigma_{RT}\} \text{ and}$$

$$L(A, x) = \{y \in A : (y, x) \in \sigma_{RT}\}.$$

Evidently,  $U(A, x), L(A, x) \subseteq A$ . In particular, if  $\alpha \in \text{Reg}(T(X, \sigma_{RT}))$ , we have the following properties.

**Lemma 2.3.** *Let  $\alpha \in \text{Reg}(T(X, \sigma_{RT}))$  and let  $x \in X$ .*

1. *If  $L(X\alpha, x) = X\alpha$ , then there exists  $m \in X\alpha$  such that  $L(X\alpha, m) = X\alpha$ .*
2. *If  $U(X\alpha, x) = X\alpha$ , then there exists  $n \in X\alpha$  such that  $U(X\alpha, n) = X\alpha$ .*

*Proof.* (1) Assume that  $\alpha = \alpha\beta\alpha$  for some  $\beta \in T(X, \sigma_{RT})$  and  $L(X\alpha, x) = X\alpha$ . Choose  $m = x\beta\alpha \in X\alpha$ . To show that  $L(X\alpha, m) = X\alpha$ , we let  $y \in X\alpha = L(X\alpha, x)$ . Then  $y = x'\alpha$  for some  $x' \in X$  and  $(y, x) \in \sigma_{RT}$ . Hence,  $y = x'\alpha = x'\alpha\beta\alpha = y\beta\alpha \subseteq X\alpha = L(X\alpha, x)$ . Thus,  $(y, m) = (y\beta\alpha, x\beta\alpha) \in \sigma_{RT}$ , and so  $y \in L(X\alpha, m)$ . Therefore,  $L(X\alpha, m) = X\alpha$ , as required.

(2) It can be proved similar to (1).  $\square$

We now characterize all regular elements in  $T(X, \sigma_{RT})$ , where  $\sigma_{RT}$  is an arbitrary reflexive and transitive relation satisfies the following condition:

$$\forall x, y \in X, (x, y) \in \sigma_{RT} \text{ or } (y, x) \in \sigma_{RT}. \quad (*)$$

Especially, Lemmas 2.3 and 2.4 give the necessary conditions for this case. The following lemma shows all remaining such conditions.

**Lemma 2.4.** *Let  $\alpha \in \text{Reg}(T(X, \sigma_{RT}))$  and  $x \in X$ . If both  $L(X\alpha, x)$  and  $U(X\alpha, x)$  are nonempty, then one of the following holds:*

1. *there exists  $u \in U(X\alpha, x)$  such that  $U(X\alpha, x) \subseteq U(X\alpha, u)$ ;*
2. *there exists  $l \in L(X\alpha, x)$  such that  $L(X\alpha, x) \subseteq L(X\alpha, l)$ .*

*Proof.* Assume that  $\alpha = \alpha\beta\alpha$  for some  $\beta \in T(X, \sigma_{RT})$  and  $L(X\alpha, x) \neq \emptyset \neq U(X\alpha, x)$ . Since  $x\beta\alpha \in X\alpha \subseteq X$ , by the property of  $\sigma_{RT}$ , we get  $(x, x\beta\alpha) \in \sigma_{RT}$  or  $(x\beta\alpha, x) \in \sigma_{RT}$ .

**Case 1:**  $(x, x\beta\alpha) \in \sigma_{RT}$ . Then we set  $u = x\beta\alpha \in U(X\alpha, x)$ . Let  $y \in U(X\alpha, x)$ . Thus  $y \in X\alpha$  and  $(x, y) \in \sigma_{RT}$ . Hence,  $y = x'\alpha$  for some  $x' \in X$  and  $(u, y) = (x\beta\alpha, x'\alpha) = (x\beta\alpha, x'\alpha\beta\alpha) = (x\beta\alpha, y\beta\alpha) \in \sigma_{RT}$ . This implies  $y \in U(X\alpha, u)$  and so  $U(X\alpha, x) \subseteq U(X\alpha, u)$ .

**Case 2:**  $(x\beta\alpha, x) \in \sigma_{RT}$ . Then we set  $l = x\beta\alpha \in L(X\alpha, x)$ . Let  $y \in L(X\alpha, x)$ . Thus  $y \in X\alpha$  and  $(y, x) \in \sigma_{RT}$ . Hence,  $y = x'\alpha$  for some  $x' \in X$  and  $(y, l) = (x'\alpha, x\beta\alpha) = (x'\alpha\beta\alpha, x\beta\alpha) = (y\beta\alpha, x\beta\alpha) \in \sigma_{RT}$ . This implies  $y \in L(X\alpha, l)$  and so  $L(X\alpha, x) \subseteq L(X\alpha, l)$ .  $\square$

**Theorem 2.5.** *Let  $\alpha \in T(X, \sigma_{RT})$ . Then  $\alpha \in \text{Reg}(T(X, \sigma_{RT}))$  if and only if, for each  $x \in X$ , the following four conditions hold.*

1. *If  $\bar{x} \cap X\alpha \neq \emptyset$ , then there exists  $\bar{y} \in \bar{X}$  such that  $\bar{x} \cap X\alpha = \bar{y}\alpha$ .*
2. *If  $L(X\alpha, x) = X\alpha$ , then there exists  $m \in X\alpha$  such that  $L(X\alpha, m) = X\alpha$ .*
3. *If  $U(X\alpha, x) = X\alpha$ , then there exists  $n \in X\alpha$  such that  $U(X\alpha, n) = X\alpha$ .*
4. *If  $L(X\alpha, x) \neq \emptyset$  and  $U(X\alpha, x) \neq \emptyset$ , then there exists  $u \in U(X\alpha, x)$  such that  $U(X\alpha, u) = U(X\alpha, x)$  or there exists  $l \in L(X\alpha, x)$  such that  $L(X\alpha, l) = L(X\alpha, x)$ .*

*Proof.* If  $\alpha \in \text{Reg}(T(X, \sigma_{RT}))$ , then (1) – (3) hold by Lemmas 2.2 – 2.3. From Lemma 2.4, there exists  $u \in U(X\alpha, x)$  such that  $U(X\alpha, x) \subseteq U(X\alpha, u)$ . Let  $y \in U(X\alpha, u)$ . Then  $y \in X\alpha$  and  $(u, y) \in \sigma_{RT}$ . Since  $(x, u) \in \sigma_{RT}$  and by transitivity of  $\sigma_{RT}$ , we have  $y \in U(X\alpha, x)$ . Hence  $U(X\alpha, u) = U(X\alpha, x)$ . Similarly, we get  $L(X\alpha, l) = L(X\alpha, x)$ .

On the other hand, suppose that all four aforementioned conditions hold. Let  $\bar{x} \in \bar{X}$ . Then, either  $\bar{x} \cap X\alpha \neq \emptyset$  or  $\bar{x} \cap X\alpha = \emptyset$ . If  $\bar{x} \cap X\alpha \neq \emptyset$ , from (1), we choose  $\bar{y} \in \bar{X}$  such that  $\bar{x} \cap X\alpha = \bar{y}\alpha$ . For each  $a \in \bar{x} \cap X\alpha$ , we choose  $a' \in \bar{y}$  such that  $a = a'\alpha$ . Define  $\beta_{\bar{x}} : \bar{x} \rightarrow \bar{y}$  by

$$a\beta_{\bar{x}} = \begin{cases} a' & \text{if } x \in X\alpha, \\ y & \text{otherwise.} \end{cases}$$

Clearly,  $\bar{x}\beta_{\bar{x}} \subseteq \bar{y}$ . Suppose that  $\bar{x} \cap X\alpha = \emptyset$ .

- If for all  $y \in X\alpha$ ,  $(y, x) \in \sigma_{RT}$ , then  $L(X\alpha, x) = X\alpha$ , from (2), there exists  $n \in X\alpha$  such that  $L(X\alpha, n) = X\alpha$ . Define  $\beta_{\bar{x}} : \bar{x} \rightarrow X$  by  $a\beta_{\bar{x}} = n'$  for all  $a \in \bar{x}$ .
- If for all  $y \in X\alpha$ ,  $(x, y) \in \sigma_{RT}$ , then  $U(X\alpha, x) = X\alpha$ , from (3), there exists  $m \in X\alpha$  such that  $U(X\alpha, m) = X\alpha$ . Define  $\beta_{\bar{x}} : \bar{x} \rightarrow X$  by  $a\beta_{\bar{x}} = m'$  for all  $a \in \bar{x}$ .
- If there exist  $m, n \in X\alpha$  such that  $(n, x), (x, m) \in \sigma_{RT}$ , then  $U(X\alpha, x) \neq \emptyset$  and  $L(X\alpha, x) \neq \emptyset$ , we first consider the case of  $u \in U(X\alpha, x)$ , such that  $U(X\alpha, u) = U(X\alpha, x)$  exists. In this case, we define  $\beta_{\bar{x}} : \bar{x} \rightarrow X$  by  $a\beta_{\bar{x}} = u'$  for all  $a \in \bar{x}$ . For the case  $U(X\alpha, u) \neq U(X\alpha, x)$  for all  $u \in U(X\alpha, x)$ , from (4), there exists  $l \in L(X\alpha, x)$  such that  $L(X\alpha, l) = L(X\alpha, x)$ . Define  $\beta_{\bar{x}} : \bar{x} \rightarrow X$  by  $a\beta_{\bar{x}} = l'$  for all  $a \in \bar{x}$ .

We notice from three cases that  $\beta_{\bar{x}}$  is a constant mapping. Now, we define  $\beta : X \rightarrow X$  by  $x\beta = x\beta_{\bar{x}}$  for all  $x \in X$ . Since  $\bar{X}$  is a partition of  $X$ , we have  $\beta$  is well-defined. Let  $(x, y) \in \sigma_{RT}$ . If  $(y, x) \in \sigma_{RT}$ , then  $\bar{x} = \bar{y}$ . It follows from the definition of  $\beta_{\bar{x}}$  that  $(x\beta, y\beta) = (x\beta_{\bar{x}}, y\beta_{\bar{x}}) \in \sigma_{RT}$ . Suppose that  $(y, x) \notin \sigma_{RT}$ .

**Case 1:**  $\bar{x} \cap X\alpha \neq \emptyset$  and  $\bar{y} \cap X\alpha \neq \emptyset$ .

Let  $x_1 \in \bar{x} \cap X\alpha$  and  $y_1 \in \bar{y} \cap X\alpha$ . By the definition of  $\beta$ , we obtain  $x_1\beta = x'_1$  where  $\bar{x}\beta \subseteq \bar{x}'_1$  and  $y_1\beta = y'_1$  where  $\bar{y}\beta \subseteq \bar{y}'_1$ . Note that  $(x'_1, y'_1) \in \sigma_{RT}$  or  $(y'_1, x'_1) \in \sigma_{RT}$ . If  $(x'_1, y'_1) \notin \sigma_{RT}$ , then  $(y_1, x_1) = (y'_1\alpha, x'_1\alpha) \in \sigma_{RT}$ . By the transitivity of  $\sigma_{RT}$ , we have  $(y, x) \in \sigma_{RT}$  which is a contradiction. Therefore  $(x_1\beta_{\bar{x}}, y_1\beta_{\bar{x}}) = (x'_1, y'_1) \in \sigma_{RT}$ . Since  $x\beta \in \bar{x}\beta_{\bar{x}} \subseteq \bar{x}'_1$  and  $y\beta \in \bar{y}\beta_{\bar{y}} \subseteq \bar{y}'_1$ , we have  $(x\beta, x'_1), (y'_1, y\beta) \in \sigma_{RT}$ . By the transitivity of  $\sigma_{RT}$ , we obtain  $(x\beta, y\beta) \in \sigma_{RT}$ .

**Case 2:**  $\bar{x} \cap X\alpha \neq \emptyset$  and  $\bar{y} \cap X\alpha = \emptyset$ .

Let  $x_1 \in \bar{x} \cap X\alpha$ . By the definition of  $\sigma_{RT}$ , we have  $(x_1, y) \in \sigma_{RT}$ . Then  $L(X\alpha, y) \neq \emptyset$ .

**Subcase 2.1:**  $L(X\alpha, y) = X\alpha$ . From the definition of  $\beta$ ,  $y\beta = m'$  where  $m \in X\alpha$  and  $L(X\alpha, m) = X\alpha$ . Since  $x_1 \in X\alpha$ , we have  $(x_1, m) \in \sigma_{RT}$ . From the transitivity of  $\sigma_{RT}$ , we get  $(x, m) \in \sigma_{RT}$ . It follows from Case 1 that  $(x\beta, y\beta) = (x\beta, m\beta) \in \sigma_{RT}$ .

**Subcase 2.2:**  $L(X\alpha, y) \neq X\alpha$ . Then  $U(X\alpha, y) \neq \emptyset$ . If  $y\beta = u'$  where  $u \in U(X\alpha, y)$  and  $U(X\alpha, u) = U(X\alpha, y)$ , then  $(y, u) \in \sigma_{RT}$ . From the transitivity of  $\sigma_{RT}$  and  $(x, y), (y, u) \in \sigma_{RT}$ , we get  $(x, u) \in \sigma_{RT}$ . It follows from Case 1 that  $(x\beta, y\beta) = (x\beta, u\beta) \in \sigma_{RT}$ . If  $y\beta = l'$  where  $l \in L(X\alpha, y)$  and  $L(X\alpha, l) = L(X\alpha, y)$ , then  $(x_1, l) \in \sigma_{RT}$  since  $x_1 \in L(X\alpha, y)$ . From the transitivity of  $\sigma_{RT}$ , we have  $(x, l) \in$

$\sigma_{RT}$ . It follows from Case 1 that  $(x\beta, y\beta) = (x\beta, l\beta) \in \sigma_{RT}$ .

**Case 3:**  $\bar{x} \cap X\alpha = \emptyset$  and  $\bar{y} \cap X\alpha \neq \emptyset$ . Let  $y_1 \in \bar{y} \cap X\alpha$ . Then  $(x, y_1) \in \sigma_{RT}$  and so  $U(X\alpha, x) \neq \emptyset$ .

**Subcase 3.1:**  $U(X\alpha, x) = X\alpha$ . From the definition of  $\beta$ ,  $x\beta = n'$  where  $n \in X\alpha$  and  $U(X\alpha, n) = X\alpha$ . Since  $y_1 \in X\alpha$ , we have  $(n, y_1) \in \sigma_{RT}$ . From the transitivity of  $\sigma_{RT}$ , we get  $(n, y) \in \sigma_{RT}$ . It follows from Case 1 that  $(x\beta, y\beta) = (n\beta, y\beta) \in \sigma_{RT}$ .

**Subcase 3.2:**  $U(X\alpha, x) \neq X\alpha$ . Then  $L(X\alpha, x) \neq \emptyset$ . If  $x\beta = u'$  where  $u \in U(X\alpha, x)$  and  $U(X\alpha, u) = U(X\alpha, x)$ , then  $(u, y_1) \in \sigma_{RT}$  since  $y_1 \in U(X\alpha, x)$ . From the transitivity of  $\sigma_{RT}$  and  $(u, y_1), (y_1, y) \in \sigma_{RT}$ , we get  $(u, y) \in \sigma_{RT}$ . It follows from Case 1 that  $(x\beta, y\beta) = (u\beta, y\beta) \in \sigma_{RT}$ . If  $x\beta = l'$  where  $l \in L(X\alpha, x)$  and  $L(X\alpha, l) = L(X\alpha, x)$ , then  $(l, x) \in \sigma_{RT}$ . From the transitivity of  $\sigma_{RT}$ , we get  $(l, y) \in \sigma_{RT}$ . It follows from Case 1 that  $(x\beta, y\beta) = (l\beta, y\beta) \in \sigma_{RT}$ .

**Case 4:**  $\bar{x} \cap X\alpha = \emptyset$  and  $\bar{y} \cap X\alpha = \emptyset$ .

**Subcase 4.1:**  $L(X\alpha, x) = X\alpha$ . Then  $L(X\alpha, y) = X\alpha$ . From the definition of  $\beta$ ,  $x\beta = m'_1$  where  $m_1 \in X\alpha$ ,  $L(X\alpha, m_1) = X\alpha$  and  $y\beta = m'_2$  where  $m_2 \in X\alpha$ ,  $L(X\alpha, m_2) = X\alpha$ . Since  $m_1 \in X\alpha$ , we have  $(m_1, m_2) \in \sigma_{RT}$ . It follows from Case 1 that  $(x\beta, y\beta) = (m_1\beta, m_2\beta) \in \sigma_{RT}$ .

**Subcase 4.2:**  $\emptyset \neq L(X\alpha, x) \neq X\alpha$  and  $L(X\alpha, y) = X\alpha$ . Then  $U(X\alpha, x) \neq \emptyset$ . From the definition of  $\beta$ ,  $y\beta = m'$  where  $m \in X\alpha$  and  $L(X\alpha, m) = X\alpha$  and  $x\beta = k'$  where  $k \in X\alpha$ . It follows that  $(k, m) \in \sigma_{RT}$ . It follows from Case 1 that  $(x\beta, y\beta) = (k\beta, m\beta) \in \sigma_{RT}$ .

**Subcase 4.3:**  $\emptyset \neq L(X\alpha, y) \neq X\alpha$  and  $\emptyset \neq L(X\alpha, x) \neq X\alpha$ . Then  $U(X\alpha, x) \neq \emptyset$  and  $U(X\alpha, y) \neq \emptyset$ . Then  $U(X\alpha, y) \subseteq U(X\alpha, x)$  and  $L(X\alpha, x) \subseteq L(X\alpha, y)$ .

If  $x\beta = u'_1$  where  $u_1 \in U(X\alpha, x)$ ,  $U(X\alpha, u_1) = U(X\alpha, x)$  and  $y\beta = u'_2$  where  $u_2 \in U(X\alpha, y)$ ,  $U(X\alpha, u_2) = U(X\alpha, y)$ , then  $(u_1, u_2) \in \sigma_{RT}$  since  $u_2 \in U(X\alpha, y) \subseteq U(X\alpha, x)$ . It follows from Case 1 that  $(x\beta, y\beta) = (u_1\beta, u_2\beta) \in \sigma_{RT}$ . If  $x\beta = l'_1$  where  $l_1 \in L(X\alpha, x)$ ,  $L(X\alpha, l_1) = L(X\alpha, x)$  and  $y\beta = l'_2$  where  $l_2 \in L(X\alpha, y)$ ,  $L(X\alpha, l_2) = L(X\alpha, y)$ , then  $(l_1, l_2) \in \sigma_{RT}$  since  $l_1 \in L(X\alpha, x) \subseteq L(X\alpha, y)$ . It follows from Case 1 that  $(x\beta, y\beta) = (l_1\beta, l_2\beta) \in \sigma_{RT}$ . If  $x\beta = u'_1$  where  $u_1 \in U(X\alpha, x)$ ,  $U(X\alpha, u_1) = U(X\alpha, x)$  and  $y\beta = l'_2$  where  $l_2 \in L(X\alpha, y)$ ,  $L(X\alpha, l_2) = L(X\alpha, y)$ . By definition of  $\beta$ , we get  $U(X\alpha, y) \neq U(X\alpha, u)$  for all  $u \in U(X\alpha, y)$ . If  $U(X\alpha, x) = U(X\alpha, y)$ , then  $u_1 \in U(X\alpha, x) = U(X\alpha, y)$  and  $U(X\alpha, u_1) = U(X\alpha, x) = U(X\alpha, y)$ . It is a contradiction. Hence  $U(X\alpha, x) \neq U(X\alpha, y)$ . There exists  $w \in U(X\alpha, x)$  and  $w \notin U(X\alpha, y)$ . Thus  $w \in X\alpha$  such that  $(x, w) \in \sigma_{RT}$  and  $(w, y) \in \sigma_{RT}$ . So  $w \in L(X\alpha, y)$ . Hence  $(w, l_2) \in \sigma_{RT}$ . By the transitivity of  $\sigma_{RT}$ ,  $(x, l_2) \in \sigma_{RT}$ . Therefore  $l_2 \in U(X\alpha, x)$  and then  $(u_1, l_2) \in \sigma_{RT}$ . It follows from Case 1 that  $(x\beta, y\beta) = (u_1\beta, l_2\beta) \in \sigma_{RT}$ . If  $x\beta = l'_1$  where  $l_1 \in L(X\alpha, x)$ ,  $L(X\alpha, l_1) = L(X\alpha, x)$  and  $y\beta = u'_2$  where  $u_2 \in U(X\alpha, y)$ ,  $U(X\alpha, u_2) = U(X\alpha, y)$ , then  $(l_1, x) \in \sigma_{RT}$  since  $l_1 \in L(X\alpha, x)$  and  $(y, u_2) \in \sigma_{RT}$  since  $u_2 \in U(X\alpha, y)$ . By the transitivity of  $\sigma_{RT}$ , we have  $(l_1, u_2) \in \sigma_{RT}$ . It follows from Case 1 that  $(x\beta, y\beta) = (l_1\beta, u_2\beta) \in \sigma_{RT}$ .

**Subcase 4.4:**  $L(X\alpha, x) = \emptyset$  and  $\emptyset \neq L(X\alpha, y) \neq X\alpha$ . Then  $U(X\alpha, x) = X\alpha$ . From the definition of  $\beta$ ,  $x\beta = n'$  where  $n \in X\alpha$ ,  $U(X\alpha, n) = X\alpha$  and  $y\beta = k'$  where  $k \in X\alpha$ . It follows that  $(n, k) \in \sigma_{RT}$ . It follows from Case 1 that  $(x\beta, y\beta) = (n\beta, k\beta) \in \sigma_{RT}$ .

**Subcase 4.5:**  $L(X\alpha, x) = \emptyset$  and  $L(X\alpha, y) = \emptyset$ . Then  $U(X\alpha, x) =$

$U(X\alpha, y) = X\alpha$ . From the definition of  $\beta$ ,  $x\beta = n'_1$  where  $n_1 \in X\alpha$ ,  $U(X\alpha, n_1) = X\alpha$  and  $y\beta = n'_2$  where  $n_2 \in X\alpha$ ,  $U(X\alpha, n_2) = X\alpha$ . Since  $n_2 \in X\alpha$ , we have  $(n_1, n_2) \in \sigma_{RT}$ . It follows from Case 1 that  $(x\beta, y\beta) = (n_1\beta, n_2\beta) \in \sigma_{RT}$ .

From four cases, we obtain  $\beta \in T(X, \sigma_{RT})$ . Finally, for each  $x \in X$ , we have  $x\alpha\beta\alpha = (x\alpha)\beta\alpha = (x\alpha)\beta_{\bar{x}\alpha}\alpha = (x\alpha)'\alpha = x\alpha$ . We conclude that  $\alpha = \alpha\beta\alpha$ .  $\square$

**Corollary 2.6.** *Let  $\alpha \in T(X, \sigma_{RT})$ . Then  $\alpha \in \text{Reg}(T(X, \sigma_{RT}))$  if and only if, for each  $x \in X$ , the following two conditions hold.*

1. *If  $\bar{x} \cap X\alpha \neq \emptyset$ , then there exists  $\bar{y} \in \bar{X}$  such that  $\bar{x} \cap X\alpha = \bar{y}\alpha$ .*
2. *If  $L(X\alpha, x) \neq \emptyset$  or  $U(X\alpha, x) \neq \emptyset$ , then there exists  $u \in U(X\alpha, x)$  such that  $U(X\alpha, u) = U(X\alpha, x)$  or there exists  $l \in L(X\alpha, x)$  such that  $L(X\alpha, l) = L(X\alpha, x)$ .*

**Theorem 2.7.** *If  $T(X, \sigma_{RT})$  is a regular semigroup, then  $\sigma_{RT} = X \times X$  or  $\sigma_{RT}$  is a totally order relation on  $X$ .*

*Proof.* Assume that  $\sigma_{RT} \neq X \times X$  and  $\sigma_{RT}$  is not a totally ordered on  $X$ . Then there exists  $y, z \in X$  such that  $(z, y), (y, z) \in \sigma_{RT}$  and  $y \neq z$ . There exists  $a \in \bar{x}$  such that  $(a, z) \notin \sigma_{RT}$  or  $(z, a) \notin \sigma_{RT}$ . Define  $\alpha : X \rightarrow X$  by

$$x\alpha = \begin{cases} y & \text{if } x \in \bar{z}, \\ z & \text{otherwise.} \end{cases}$$

Clearly that  $X\alpha = \{y, z\}$ . Thus  $\alpha$  preserves relation  $\sigma_{RT}$ . Note that  $\bar{z} \cap X\alpha = \{y, z\} \neq \bar{x}\alpha$  for all  $\bar{x} \in \bar{X}$ . Hence  $\alpha$  is not regular.  $\square$

### 3. Conclusion

Our research yields crucial finding on the regularity of elements in set  $T(X, \sigma_{RT})$ , with the satisfaction of condition (\*). Specifically, we establish that a regular semigroup  $T(X, \sigma_{RT})$  implies  $\sigma_{RT} = X \times X$  or  $\sigma_{RT}$  is a totally ordered relation. Nevertheless, there is room for further advancement in this field by either eliminating condition (\*) or substituting it with weaker conditions, thus paving the way for future investigations.

### References

- [1] Howie JM. Fundamentals of Semigroup Theory, Oxford: Clarendon Press; 1995.
- [2] Pei H. Regularity and Green's relations for semigroups of transformations that preserve an equivalence, Communications in Algebra 2005;33(1):109-18.
- [3] Kaewnoi T, Petapirak M, Chinram R. Magnifying elements in a semigroup of transformations preserving equivalence relation, Korean Journal of Mathematics 2019;27:269-77.
- [4] Pei H. Equivalences,  $\alpha$ -semigroups and  $\alpha$ -congruences, Semigroup Forum 1994;49(1):49-58.
- [5] Pei H. A regular  $\alpha$ -semigroup inducing a certain lattice, Semigroup Forum 1996;53:98-113.
- [6] Pei H. On the rank of the semigroup  $T_E(X)$ , Semigroup Forum 2005;70:107-17.
- [7] Pei H, Zhou H. Abundant semigroups of transformations preserving an equivalence relation, Algebra Colloquium 2011;18(1):77-82.
- [8] Sun L, Pei H, Cheng Z. Naturally ordered transformation semigroups preserving an equivalence, Bulletin of the Australian Mathematical Society 2008;78:117-28.

- [9] Mora W, Kemprasit Y. Regular elements of some order-preserving transformation semigroups, *International journal of Algebra* 2010;4(13):631-41.
- [10] Fernandes VH, Honyam P, Quinteiro TM, Singha B. On semigroups of endomorphisms of a chain with restricted range, *Semigroup Forum* 2014;89(1):77-104.
- [11] Gluskin LM. Semi-groups of isotone transformations, *Uspekhi Matematicheskikh Nauk* 1961;16(5):157-62.
- [12] Higgins PM. *Techniques of Semigroup Theory*, New York: Oxford University Press; 1992.
- [13] Kemprasit Y, Changphas T. Regular order-preserving transformation semigroups, *Bulletin of the Australian Mathematical Society* 2000;62:511-24.
- [14] Repnitskiĭ VB, Vernitskiĭ AS. Semigroups of order-preserving mappings, *Communications in Algebra* 2000;28:3635-41.