

Patraporn Luksirikul 2006: Hydroboration Reaction: The Effect of B and N Atoms Doping onto the (5,5) Armchair Sidewalls of Single-Wall Carbon Nanotubes (SWNTs). Master of Science (Chemistry),
Major Field: Chemistry, Department of Chemistry.
Thesis Advisor: Assistant Professor Piboon Pantu, Ph.D. 52 pages.
ISBN 974-16-2786-6

Hydroboration of the perfect structure and the defective structure which doped B,N atoms onto the sidewall of (5,5) armchair single-wall carbon nanotubes with borane (BH_3) has been investigated within the framework of our-Own-N-layered-Integrated molecular Orbital and molecular Mechanics (ONIOM) approaches utilizing the two layered ONIOM scheme. Three different systems of hydroboration reaction: (1) perfect sidewall of SWNT; (2) BN-doped SWNT; (3) NB-doped SWNT, respectively were studied and found to occur through a concerted mechanism of the electrophilic attack of borane to carbon nanotubes. The transition states and products of doped systems onto hydroboration were stabilized by electron transfer from B and N atoms substituted nearby the reactive site to C reactive center (demonstrated by NPA analysis). The BN-doped and NB-doped activation energies of hydroboration with could enhance reactivity of (5,5) armchair single-wall carbon nanotubes decreased by 4.43 and 2.32 kcal/mol, respectively such their reaction energies

Student's signature

Thesis Advisor's signature

ACKNOWLEDGEMENTS

First, I would like to express my sincere gratitude to Professor Dr. Jumras Limtrakul for being my advisor. Under his guidance, encouragement and excellent advice during to complete this thesis, I successfully overcame many difficulties and gained much knowledge. I am grateful to my graduate committees, Assistant Professor Piboon Pantu, Ajarn Pensri Bunsawansong, and Associate Professor Penchantr Singh, a representative of the Graduate School of Kasetsart University for their kind advice and willpower that they instilled in me. I want to thank Dr. Pipat Kongpracha, Supawadee Namuangruk and Bundet Boekfa for their invaluable help on this thesis. Thanks also to the Laboratory for Computational and Applied Chemistry (LCAC) and the Pirun Cluster, Kasetsart University for a good computational environment. I acknowledge the financial support by the Development and Promotion of Science and Technology Talent Projects (DPST), Kasetsart University Research and Development Institute (KURDI), the Ministry of University Affairs under the Science and Technology Higher Education Development Project (Petroleum and Petrochemical Consortium funded by MUA-ADB), the Thailand Research Fund, and the National Nanotechnology Center.

Last but certainly not least, I would like to show my gratitude to my parents, Houyhuang, Damrong and my lovely sisters Luckkana Luksirikul who have been generous with their encouragement throughout, for their continuous and unconditional support of all my undertakings, scholastic and otherwise. This thesis is dedicated to all of them whom I love deeply.

Patraporn Luksirikul

September, 2006