

## TABLE OF CONTENTS

	<b>Page</b>
TABLE OF CONTENTS	i
LIST OF TABLES	iii
LIST OF FIGURES	iv
INTRODUCTION	1
LITERATURE REVIEWS	3
Biology and Characteristic of <i>Ralstonia solanacearum</i>	3
Pathogenicity and Host Range of <i>Ralstonia solanacearum</i>	5
Molecular Characterization and Genetic Diversity of <i>Ralstonia solanacearum</i>	6
A Model System for the Study of Bacterial Pathogenicity	7
Insights from the <i>R. solanacearum</i> Genome Sequence	10
Possible Roles for Polyols in Plant Pathogen	12
MATERIALS AND METHODS	14
Bacterial Strain and Growth Condition	14
Pathogenicity Tests and Biovar Classification	16
DNA Amplification	16
Enzyme Activity Assay	18
Chromosomal DNA and Plasmid Preparation	19
DNA Manipulation	20
Construction of Expression <i>polS</i> in Plasmid pGEX-2T	22
Expression of Recombinant SDH	23
Protein Analysis on SDS-PAGE analysis	23
Construction for IDM Plasmid for Gene Disruption by Homologous Recombination	25
Southern Blot Hybridization	27
Quantification of EPS	29

## TABLE OF CONTENTS (cont'd)

	<b>Page</b>
RESULTS	31
Amplification of <i>polS</i> Gene by Specific Primers and	
Hybridization of <i>polS</i> Gene	31
Nucleotide Sequencing and Analysis of <i>polS</i> Gene	35
SDS-PAGE of Expression Sorbitol Dehydrogenase	36
Construction of Plasmid for Insertion-Duplication Mutagenesis (IDM)	44
Conjugation pKS into <i>R. solanacearum</i> and Detection for IDM	
and AE by PCR and Hybridization Analysis	44
Biovar Test and Enzyme Activity Assay for Mutation	53
Pathogenicity, HR Test and EPS Quantification of	
Non-oxidizing Sorbitol Mutant	57
DISCUSSION	62
CONCLUSION	72
LITERATURE CITED	74
APPENDIX	91

## LIST OF TABLES

Table		Page
1	Differentiation of <i>Ralstonia solanacearum</i> biovars based on utilization and oxidation of sugar as positive (+) and negative (-)	4
2	Bacterial strains and plasmids used in this study	15
3	List of oligonucleotides used to amplify internal and whole <i>polS</i> gene regions for identification and mutagenesis in this study	17
4	List of oligonucleotides used for PCR analysis of mutants	18
5	Sources of sorbitol dehydrogenase protein sequences from Different strains of <i>Ralstonia solanacearum</i> and other bacteria	40
6	Enzyme activity of crude extract from <i>Ralstonia solanacearum</i> sorbitol dehydrogenase	56
7	Pathogenicity Tests of <i>Ralstonia solanacearum</i> wild type (To-Ud3-WT), nalidixic marker (To-Ud3-N) and non-oxidizing sorbitol (MT5) strains on tomato host (L390)	59
8	The amount of EPS produced by <i>Ralstonia solanacearum</i> wild type (To-Ud3-WT), nalidixic marker (To-Ud3-N) and mutant strains in rich medium	60
 <b>Appendix Table</b>		
1	Dilution series of BSA standards	94
2	The correlation between final concentration of standard BSA and their absorbance value at 595 nm	95
3	Dilution series of <i>N</i> -acetyl-D-glucosamine standards	96
4	The correlation between final concentration of standard <i>N</i> -acetyl-D-glucosamine and their absorbance value at 530 nm	97

## LIST OF FIGURES

Figure	Page
1 Agarose gel electrophoresis of 600 bp fragment amplified with sorF and sorR primers and 800 bp fragment amplified by polSF and polSR primers using DNA extracted from <i>Ralstonia solanacearum</i> as a template	33
2 Southern blot analysis between <i>sorS</i> probe and DNA extracted from different biovars of <i>Ralstonia solanacearum</i> and partially digested with <i>Cla</i> I showing specific 1.8 kb DNA product in biovars 3 and 4	34
3 Partial nucleotide and deduced amino acid sequence of sorbitol dehydrogenase ( <i>polS</i> ) of <i>R. solanacearum</i> strain To 264	37
4 Multiple sequence alignment of the SDR proteins	38
5 Phylogenetic tree of sorbitol dehydrogenase protein sequences from seven bacteria. Neighbor joining method of ClustalW program was used to construct phylogenetic trees	41
6 Nucleotide and amino acid sequence similarity of sorbitol dehydrogenase (SDH) and its homologues among seven bacterial species and percent nucleotide similarity is denoted above the diagonal and percent amino acid similarity is below	42
7 SDS-PAGE of crude extract from <i>E. coli</i> harboring pGEX-2T and pGEX- <i>polS</i> in which cells were harvested at 2, 4, and 6 hr. after induction	43
8 The creation of deletion constructs and the nucleotide sequence of endogenous <i>polS</i> gene was deleted by overlapping PCR method	47
9 Southern blot hybridization between <i>kan</i> and <i>sorS</i> probe with <i>Bam</i> HI- <i>Eco</i> RI or <i>Cla</i> I partially digested DNA extracted from wild type strain and transconjugant clones	49
10 Electroporesis gel of PCR product amplified by various primers located upstream and downstream of endogenous <i>polS</i> gene ( <i>cla</i> F and <i>cla</i> R), located on plasmid ( <i>mob</i> F- <i>mob</i> R and <i>km</i> F- <i>km</i> R) and located on <i>polS</i> gene ( <i>sor</i> F and <i>sor</i> R)	50

## LIST OF FIGURES (cont'd)

Figure	Page
11 Generation of a non-polar insertion mutation.	52
12 Comparison of biovar test between <i>Ralstonia solanacearum</i> wild type strain (To-Ud3) and mutant	54
13 Enzyme activity assay for conversion of sorbitol by the sorbitol dehydrogenase from <i>Ralstonia solanacearum</i> wild type (To-Ud3-WT), nalidixic marker (To-Ud3-N) and mutants (MT5, MT6, MT7). The absorbance of crude extract from biovar 1 (FC328) and 2 (PO1155) were measured as a control	55
14 Bacterial wilt symptom and hypersensitive reaction (HR) test caused by <i>Ralstonia solanacearum</i> strain To-Ud3 and non-oxidizing sorbitol mutant	61
 <b>Appendix Figure</b>	
1 The standard curve of BSA ( $\mu\text{g/ml}$ ) and their absorbance value at OD <sub>595</sub>	95
2 The standard curve of <i>N</i> -acetyl-D-glucosamine ( $\mu\text{g/ml}$ ) and their absorbance value at OD <sub>530</sub>	97
3 pGEM <sup>®</sup> -T Easy Vector circular map and sequence reference points	100
4 The promoter and multiple sequence of the pGEM <sup>®</sup> -T Easy Vector	101
5 pGEX-2T Expression Vector circular map and multiple cloning site	102
6 pK18mob Integrating Vector circular map and multiple cloning sites	104