

เอกสารอ้างอิง

- [1] A.I. Gasco-Lopez, R. Izquierdo-Hornillos, A. Jimenez, **“Development and validation of a high performance liquid chromatography method for the determination of cold relief ingredients in chewing gum”**, Journal of Chromatography A. 775 (1997) 179-185
- [2] A. Marin, E. Garcia, A. Garcia, C. Barbas, **“Validation of a HPLC quantification of acetaminophen, phenylephrine and chlorpheniramine in pharmaceutical formulations : capsules and sachets”**, Journal of Pharmaceutacal and Biomedical Analysis 29 (2002) 701-714
- [3] Susumu Yamato, Masaharu Nakajima, Kenji Shimada, **“Simultaneous determination of chlorpheniramine and maleate by High performance liquid chromatography using tetra-*n*-butylammonium phosphate as an ion-pair reagent”**, Journal of Chromatography A, 731 (1996) 346-350
- [4] Darryl J. Hood, H.Y. Cheung, **“A chromatographic method for rapid and simultaneous analysis of codeine phosphate, ephedrine HCl and chlorpheniramine maleate in cough-cold syrup formulation.”**, Journal of Pharmaceutical and Biomedical Analysis 30(2003) 1595-1601
- [5] Thomas A. Biemer., **“Simultaneous analysis of acetaminophen, pseudoephedrine HCl and chlorpheniramine maleate in a cold tablet using an isocratic, mixed micellar high performance liquid chromatography mobile phase”**., Journal of Chromatography, 410(1987) 206-210
- [6] Leena Suntornsuk, Ongart Pipitharome, Prapin Wilairat., **“Simultaneous determination of paracetamol and chlorpheniramine maleate by micellar electrokinetic chromatography”**, Journal of Pharmaceutical and Biomedical Analysis 33(2003)441-449
- [7] Krishna K. Verma, Sunil K. Sanghi, Archana Jain and Dayashanker Gupta., **“Determination of aspirin by pre-column transacetylation reaction of 3-aminophenol and reversed phase high performance liquid chromatography : Simultaneous determination of aspirin, acetaminophen and caffeine”**, Journal of Pharmaceutical Sciences Vol.76, No.7, July (1987) 551-553
- [8] <http://www.Science.manager200online.htm>.

- [9] Lijun He, Wenzhu Zhang, Liang Zhao, Xia Liu, Shengxiang Jiang., “ **Effect of 1-alkyl-3-methylimidazolium-based ionic liquids as the eluent on the separation of ephrines by liquid chromatography**”, Journal of Chromatography A, 1007(2003)39-45
- [10] สมิงเก่า เจริญ “สารพิษ” หมอชาวบ้าน, 2541, 298-302
- [11] สัมมน โฉมฉาย, สุรจิต สุนทรธรรม “Non-Narcotic Analgesics and NSAIDs Overdose” เวชศาสตร์ฉุกเฉิน, 2542; 847-854
- [12] Richard J. Lewis, Sr. Hawley’s Condensed Chemical Dictionary; 13th edition
- [13] <http://www.derlarrubiz.com>
- [14] <http://www.keepkidshealthy.com/adolescent/adolescent/adolescent.html>
- [15] พวงแก้ว ลัคณาทิพนธ์ “ความรู้พื้นฐานทางลึ่วิตโครมาโทกราฟี” คณะเภสัชศาสตร์ มหาวิทยาลัยมหิดล, 2540
- [16] คณิตา ดังคนานุรักษ์ “เทคนิคการแยกสารทางเคมี” โครงการตำราคณะวิทยาศาสตร์ สถาบันเทคโนโลยีพระจอมเกล้าเจ้าคุณทหารลาดกระบัง, พฤษภาคม, 2542
- [17] Trevor M. Letcher, Urszula Domanska, Malgorzata Marciniak, Andrzej Marciniak, “ **Activity coefficients at infinite dilution measurements for organic solutes in the ionic liquid 1-butyl-3-methyl-imidazolium 2-(2-methoxyethoxy) ethyl sulfate using g.l.c. at T = (298.15, 303.15, and 308.15) K,**” J. Chem. Thermodynamics 37 (2005) 587– 593
- [18] Fabrice Mutelet, Jean-Noel Jaubert, “**Accurate measurements of thermodynamic properties of solutes in ionic liquids using inverse gas chromatography,**” Journal of Chromatography A, 1102 (2006) 256–267
- [19] Zsombor Miskolczy, Krisztina Sebok-Nagy, Laszlo Biczok, Sinem Gokturk, “**Aggregation and micelle formation of ionic liquids in aqueous solution,**” Chemical Physics Letters 400 (2004) 296–300
- [20] A. Vicent Orchilles, Vicenta Gonzalez-Alfaro, Pablo J. Miguel, Ernesto Vercher, Antoni Martinez-Andreu, “**Volumetric properties of binary mixtures of ionic liquid 1-butyl-3-methylimidazolium octylsulfate with water or propanol in the temperature range of 278.15 K to 328.15 K,**” J. Chem. Thermodynamics 38 (2006) 1124–1129
- [21] Enrique G. Yanes, Samuel R. Gratz, Michael J. Baldwin, Sara E. Robison, and Apryll M. Stalcup, “ **Capillary Electrophoretic Application of 1-Alkyl-3-methylimidazolium-Based Ionic Liquids,**” Anal. Chem.(2001)73 3838-3844

- [22] A. Garcia , F.J. Ruperez , A. Marin , A. de la Maza , C. Barbas, **“Poly(ethyleneglycol) column for the determination of acetaminophen, phenylephrine and chlorpheniramine in pharmaceutical formulations”** Journal of Chromatography B, 785 (2003) 237–243
- [23] A. Marin, C. Barbas, **“CE versus HPLC for the dissolution test in a pharmaceutical formulation containing acetaminophen, phenylephrine and chlorpheniramine”**, Journal of Pharmaceutical and Biomedical Analysis 35 (2004) 769–777
- [24] B. Olmo, A. Garcia, A. Marin, C. Barbas, **“New approaches with two cyano columns to the separation of acetaminophen, phenylephrine, chlorpheniramine and related compounds”**, Journal of Chromatography B, 817 (2005) 159–165
- [25] J.C. Miller and J.N miller, “ statistics for analytical chemistry” 3rd edition, 1993, 58, 104 – 117
- [26] Werner Funk, Vera Dammann Gerhild Donnevert, “ Quality Assurance in Analytical Chemistry” , 1995, 29