

บรรณานุกรม

- กรมปศุสัตว์. 2553. ข้อมูลจำนวนเกษตรกรและปศุสัตว์ในประเทศไทยประจำปี 2553. ระบบฐานข้อมูลเกษตรกรผู้เลี้ยงสัตว์ (TH-LIFDS) ศูนย์สารสนเทศกรมปศุสัตว์ กระทรวงเกษตรและสหกรณ์ [Online] available: http://www.dld.go.th/ict/th/index.php?option=com_content&view=article&id=371:-2553&catid=207:-2553&Itemid=123. Citation: 30 พฤศจิกายน 2553.
- จรัญ จันทลักษณ์. 2515. การผลิตโคเนื้อ. ภาควิชาสัตวบาล. คณะเกษตรศาสตร์. มหาวิทยาลัยเกษตรศาสตร์.
- จุฑารัตน์ เศรษฐกุล และญาณิน โอภาสพัฒนกิจ. 2548. คุณภาพเนื้อโคภายใต้ระบบการผลิตและการตลาดของประเทศไทย. กรุงเทพมหานคร: บริษัทสุพีเรียพรีนดิ้งแฮ็ส จำกัด. 85 หน้า.
- สุรัชย์ ชาศรีรัตน์. 2545. การสืบพันธุ์และการผสมเทียมโค-กระบือ. กรุงเทพฯ: สำนักส่งเสริมและฝึกอบรม มหาวิทยาลัยเกษตรศาสตร์. 439 หน้า
- ศรเทพ ธรรมวาสร. 2548. การเลี้ยงโคเนื้อ: แนวทางการพัฒนาอาชีพของเกษตรกรไทย. พิมพ์ครั้งที่ 3 (ปรับปรุงใหม่). กรุงเทพฯ: อักษรสยามการพิมพ์. 452 หน้า.
- Blakely, J., and D.H. Bade. 1994. The science of Animal Husbandry. 6th Edition. Prentice Hall Career and Technology, Prentice-Hall, Inc., New Jersey, USA.
- Becker, P.L., Singer J.J., Walsh J.V.J.R. and Fay F.S. 1989. Regulation of calcium concentration in voltage clamped smooth muscle cells. Science. 244:211–214.
- Berlin J.R. and Konishi M. 1993. Ca²⁺ transients in cardiac myocytes measured with high and low affinity Ca²⁺ indicators. Biophys.J. 65:1632–1647.
- Betts, R., S. Weinsheimer, G.E. Blouse and J. Anagli. 2003. Structural determinants of the calpain inhibitory activity of calpastatin peptide B27-WT. J. Biol. Chem. 278(10):7800-7809. Epub 2002 Dec 24.
- Boccard, R., I. Buchter, M. Casteels, E. Cosentino, E. Dransfield and D.E. Hood. 1981. Procedures for measuring meat quality characteristics in beef production experiments. Lives. Prod. Sci. 8: 385-397.
- Boehm, M.L., T.L. Kendall, V.F. Thompson and D.E. Goll. 1998. Changes in the calpains and calpastatin during postmortem storage of bovine muscle. J. Anim. Sci. 76:2415-2434.

- Calkins C.R. and G. Sullivan. 2007. Ranking of beef muscles for tenderness. In Beef facts product enhancement. National Cattlemen's Beef Association. CO: Centennial.
- Cong, M., V.F.Thompson, G.E. Goll and P.B. Antin. 1998. The bovine calpastatin gene promoter and a new N-terminal region of the protein are targets for cAMP-dependent protein kinase activity. *J. Biol. Chem.* 273(1):660-666.
- Crawford, C. 1990. Protein and peptide inhibitors of calpains. In: Intracellular Calcium-Dependent Proteolysis. Boca Raton. FL:CRC. p.75-89.
- Cundiff, L.V., K.E. Gregory and J.E. Gosey. 1993. Cattle germplasm evaluation and germplasm utilization programs at MARC. *Proc. Recip. Meat Conf.* 46:45.
- Dargelos, E., S. Poussard, C. Brule, L. Daury, and P. Cottin. 2008. Calcium-dependent proteolytic system and muscle dysfunctions: a possible role of calpains in sarcopenia. *Biochimie.* 90: 359-368.
- Dransfield, E. 1994. Tenderness of meat, poultry and fish. in Pearson, A. M. and Dutson, T. R. *Quality Attributes and their Measurement in Meat, Poultry and Fish Product.* London: Blackie Academic and Professional. p. 289-315.
- Delgado, E.F., G.H. Geesink, J.A. Marchello, D.E. Goll and M. Koochmaraie. 2001. The calpain system in three muscles of normal and callipyge sheep. *J. Anim. Sci.* 79: 398-412.
- De Tullio, R., Aversa M., Stifanese R., Parr T., Bardsley R.G., Pontremoli S., Melloni E. 2007. Multiple rat brain calpastatin forms are produced by distinct starting points and alternative splicing of the N-terminal exons. *Arch. Biochem. Biophys.* 465(1):148-156.
- Ensminger, M.E. 1993. *Dairy Cattle Science.* 5th Edition. The Interstate Printer and Publisher, Inc. Danville, Illinois, USA. p. 550.
- Emori, Y., H. Kawasaki, S. Imajoh, Y. Minami and K. Suzuki. 1988. All four repeating domains of the endogenous inhibitor for calcium-dependent protease independently retain inhibitory activity. Expression of the cDNA fragments in *Escherichia coli*. *J. Biol. Chem.* 263(5):2364-2370.
- Ernst, C.W., A. Robic, M. Yerle, L. Wang and M.F. Rothschild. 1998. Mapping of calpastatin and three microsatellites to porcine chromosome 2q2.1-q2.4. *Anim. Genet.* 29(3):212-215.

- Fries, R. and A. Ruvinsky. 1999. *The Genetics of Cattle*. CABI Publishing, New York.
- Geesink, G.H. and M. Koohmaraie. 1999. Technical note: a rapid method for quantification of calpain and calpastatin activities. *J. Anim. Sci.* 77:3225-3229.
- Geesink, G.H., D. Nonneman and M. Koohmaraie. 1998. An improved purification protocol for heart and skeletal muscle calpastatin reveals two isoforms resulting from alternative splicing. *Arch. Biochem. Biophys.* 356:19–24.
- Geesink, G.H., A. Ouali and F.J.M. Smulders. 1992. Tenderisation, calpain/ calpastatin activities and osmolality of 6 different beef muscles. In: *Proc. 38th Int. Cong. of Meat Sci. and Technol.*, Clermont Ferrand. pp 363-366.
- Geesink, G.H., F.J. Smulders, H.L. van Laack, J.H. van der Kolk, T. Wensing and H.J. Breukink. 1993. Effect on meat quality of the use of clenbuterol in veal calves. *J. Anim. Sci.* 71:1161-1170.
- Goll, D.E., Kleese W.C., Okitani A., Kumamoto T., Cong J., and Kapprell H.-P. 1990. Historical background and current status of the Ca²⁺-dependent proteinase system. In: *Intracellular Calcium-Dependent Pro-teolysis*, edited by Mellgren RL and Murachi T. Boca Raton, FL:CRC. p.3–24.
- Goll, D.E., V.F. Thompson, H. Li, W. Wei and J. Cong. 2003. The calpain system. *Physiol Rev.* 83(3):731-801.
- Hall, T. 2007. BioEdit: Biological Sequence Alignment Editor for Windows 95/98/NT/XP. [Online]. Available: <http://www.mbio.ncsu.edu/BioEdit/bioedit.html>.
- Hoppe, B.L., Conti-Tronconi, B.M., and Horton, R.M. 1992. Gel loading dyes compatible with PCR. *BioTechniques* 12:679-680.
- Huang, X. 1992. A contig assembly program based on sensitive detection of fragment Overlaps. *Genomics.* 14(1): 18-25.
- Imajoh S., H. Kawasaki, Y. Emori and K. Suzuki. 1987. Calcium-activated neutral protease inhibitor from rabbit erythrocytes lacks the N-terminal region of the liver inhibitor but retains three inhibitory units. *Biochem. Biophys. Res. Commun.* 146:630–632.
- Inazawa, J., H. Nakagawa, S. Misawa, T. Abe, S. Minoshima, R. Fukuyama, M. Maki, T. Murachi, M. Hatanaka and N. Shimizu. 1990. Assignment of the human calpastatin

- gene (CAST) to chromosome 5 at region q14-q22. *Cytogenet Cell Genet.* 54(3-4):156-158.
- Inomata M., Nakamura M., Imajoh-ohmi S, and Kawashima S. 1993. A variety of calpain/calpastatin systems in mammalian erythrocytes. *Biochim. Biophys. Acta.* 1178:207–214.
- Inoue H., Nojima H., and Okayama H. 1990. High efficiency transformation of *Escherichia coli* with plasmids. *Gene* 96: 23-28.
- Iwamoto, H., Miura, T., Okamura, T., Shirakawa, K., Iwatate, M., Kawamura, S., Tatsuno, H., Ikeda, Y., Matsuzaki, M. 1999. Calpain inhibitor-1 reduces infarct size and DNA fragmentation of myocardium in ischemic/reperfused rat heart. *J. Cardiovasc. Pharmacol.* 33(4):580-586.
- Jiao W., McDonald D.Q., Coxon J.M. and Parker E.J. 2010. Molecular modeling studies of peptide inhibitors highlight the importance of conformational prearrangement for inhibition of calpain. *Biochem.* 49(26):5533-5539.
- Kapprell, H.P. and D.E. Goll. 1989. Effect of Ca^{2+} on binding of the calpains to calpastatin. *J. Biol. Chem.* 264:17888-17896.
- Kawasaki H., Emori Y. and Suzuki K. 1993. Calpastatin has two distinct sites for interaction with calpain – effect of calpastatin fragments on the binding of calpain to membranes. *Arch.Biochem.Biophys.* 305:467–472.
- Koohmaraie, M. 1988. The role of endogenous proteases in meat tenderness. *Proc. Recip. Meat Conf.* 41:89.
- Koohmaraie, M., G. Whipple, D.H. Kretchmar, J.D. Crouse, and H.J. Mersmann. 1991. Postmortem proteolysis in longissimus muscle from beef, lamb and pork carcasses. *J. Anim. Sci.* 69:617-624.
- Koohmaraie, M. 1992. Role of the neutral proteinases in postmortem muscle protein degradation and meat tenderness. *Proc. Recip. Meat Conf.* 45:63.
- Koohmaraie, M. 1994. Muscle proteinases and meat ageing. *Meat Sci.* 36: 93-104.
- Koohmaraie, M. 1996. Biochemical factors regulating the toughening and tenderization process of meat. *Meat Sci.* 43:193-201.

- Kunz, S., Niederberger, E., Ehnert, C., Coste, O., Pfenninger, A., Kruij, J., Wendrich, T.M., Schmidtko, A., Tegeder, I., Geisslinger, G. 2004. The calpain inhibitor MDL 28170 prevents inflammation-induced neurofilament light chain breakdown in the spinal cord and reduces thermal hyperalgesia. *Pain*. 110(1):409-418.
- Laemmli, U.K. 1970. Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature*. 227:680 - 685.
- Larkin, M.A., Blackshields, G., Brown, N.P., Chenna, R., McGettigan, P.A., McWilliam, H., Valentin, F., Wallace, I.M., Wilm, A., Lopez, R., Thompson, J.D., Gibson, T.J. and Higgins, D.G. 2007. ClustalW and ClustalX version 2. *Bioinformatics*. 23(21):2947-2948.
- Lee, W. J., Hatanaka, M. and Maki, M. 1992a. Multiple forms of rat calpastatin cDNA in the coding region of functionally unknown amino-terminal domain. *Biochim. Biophys. Acta*. 1129:251-253.
- Lee, W. J., Ma, H., Takano, E., Yang, H. Q., Hatanaka, M., and Maki, M. 1992b. Molecular diversity in amino-terminal domains of human calpastatin by exon skipping. *J. Biol. Chem*. 267:8437–8442.
- Limaye, P.B., Bhave, V.S., Palkar, P.S., Apte, U.M., Sawant, S.P., Yu, S., Latendresse, J.R., Reddy, J.K., Mehendale, H.M. 2006. Upregulation of calpastatin in regenerating and developing rat liver: role in resistance against hepatotoxicity. *Hepatology*. 44(2):379–388.
- Ma H., Yang H.Q., Takano E., Hatanaka M. and Maki M. 1994. Amino terminal conserved region in proteinase inhibitor domain of calpastatin potentiates its calpain inhibitory activity by interacting with the calmodulin-like domain of the proteinase. *J. Biol. Chem*. 269:24430–24436.
- Maki M., Hatanaka M., Takano E. and Murachi T. 1990. Structure-function relationship of the calpastatins. In: *Intracellular Calcium-Dependent Proteolysis*, edited by Mellgren R.L. and Murachi T. Boca Raton, FL:CRC. p 37–51.
- Melloni E., de Tullio R., Aversa M., Tedesco I., Salamino F., Sparatore B. and Pontremoli S. 1998. Properties of calpastatin isoforms in rat brain. *FEBS Lett*. 431:55–58.
- Murachi T. 1989. Intracellular regulatory system involving calpain and calpastatin. *Biochem. Int*. 18:263–294.

- Nakamura Y., Fukiage C., Ma H., Shih M., Azuma M. and Shearer T.R. 1999. Decreased sensitivity of lens-specific calpain Lp82 to calpastatin inhibitor. *Experimental Eye Research*. 69(2):155-162.
- Nishimura T. and Goll D.E. 1991. Binding of calpain fragments to calpastatin. *J. Biol. Chem.* 266:11842–11850.
- O'Connor, S.F., J.D. Tatum, D.M. Wulf, R.D. Green and G.C. Smith. 1997. Genetic Effects on Beef Tenderness in *Bos indicus* Composite and *Bos taurus* Cattle. *J. Anim. Sci.* 75:1822-1830.
- Otsuka Y. and Goll D.E. 1987. Purification of the Ca^{2+} -dependent proteinase inhibitor from bovine cardiac muscle and its interaction with the millimolar Ca^{2+} -dependent proteinase. *J. Biol. Chem.* 262:5839–5851.
- Ouali, A., and A. Talmant. 1990. Calpains and calpastatin distribution in bovine, porcine and ovine skeletal muscle. *Meat Sci.* 28:331-348.
- Parr, T., P.L. Sensky, R.G. Bardsley and P.J. Buttery. 2001. Calpastatin expression in porcine cardiac and skeletal muscle and partial gene structure. *Arch. Biochem. Biophys.* 395(1):1-13.
- Pfaffl, M.W. 2004. Quantification strategies in real-time PCR. In: Bustin SA (ed), *A–Z of Quantitative PCR*. pp. 87–120. La Jolla, CA: IUL Biotechnology Series, International University Line.
- Pringle, T.D., S.E. Williams, B.S. Lamb, D.D. Johnson and R.L. West. 1997. Carcass characteristics, the calpain proteinase system, and aged tenderness of Angus and Brahman crossbred steers. *J. Anim. Sci.* 75(11):2955-2961.
- Posmantur, R., Kampfl, A., Siman, R., Liu, J., Zhao, X., Clifton, G.L., Hayes, R.L. 1997. A calpain inhibitor attenuates cortical cytoskeletal protein loss after experimental traumatic brain injury in the rat. *Neurosci.* 77(3):875-888.
- Rao, M.V., Mohan, P.S., Peterhoff, C.M., Yang, D-S., Schmidt, S.D., Stavrides, P.H., Campbell, J., Chen, Y., Jiang, Y., Paskevich, P.A., Cataldo, A.M., Haroutunian, V., and Nixon R.A. 2008. Marked calpastatin (CAST) depletion in Alzheimer's disease accelerates cytoskeleton disruption and neurodegeneration: Neuroprotection by CAST overexpression. *J. Neurosci.* 28(47):12241–12254.

- Raynaud, P., M. Gillard, T. Parr, R. Bardsley, V. Amarger and H. Leveziel. 2005a. Correlation between bovine calpastatin mRNA transcripts and protein isoforms. *Arch. Biochem. Biophys.* 440:46-53.
- Raynaud, P., C. Jayat-Vignoles, M.P. Laforet, H. Leveziel, and V. Amarger. 2005b. Four promoters direct expression of the calpastatin gene. *Arch. Biochem. Biophys.* 437:69-77.
- Sambrook, J., Fritsch, E.F. and Maniatis, T. 1989. *Molecular Cloning: A Laboratory Cold Spring Harbor, New York: Cold Spring Harbor Press.*
- Sazili, A.Q., Parr, T., Sensky, P.L., Jones, S.W., Bardsley, R.G. and Buttery, P.J. 2005. The relationship between slow and fast myosin heavy chain content, calpastatin and meat tenderness in different ovine skeletal muscles. *Meat Sci.* 69(1):17-25.
- Shackelford, S.D., M. Koohmaraie, G. Whipple, T.L. Wheeler, M.F. Miller, J.D. Crouse, and J.O. Reagan. 1991. Predictors of beef tenderness: Development and verification. *J. Food Sci.* 56:1130-1140.
- Shackelford, S.D., M. Koohmaraie, L.V. Cundiff, K.E. Gregory, G.A. Rohrer and J.W. Savell. 1994. Heritabilities and phenotypic and genetic correlations for bovine postrigor calpastatin activity, intramuscular fat content, Warner-Bratzler shear force, retail product yield, and growth rate. *J. Anim. Sci.* 72:857.
- Shackelford, S.D., T.L. Wheeler, and M. Koohmaraie. 1995. Relationship between shear force and trained sensory panel tenderness ratings of 10 major muscles from *bos-indicus* and *bos-taurus* cattle. *J. Anim. Sci.* 73:3333-3340.
- Spencer, M.J. and Mellgren R.L. 2002. Overexpression of a calpastatin transgene in mdx muscle reduces dystrophic pathology. *Hum. Mol. Genet.* 11(21):2645-2655.
- Souaze F, Ntodou-Thome A, Tran CY, Rostene W, Forgez P. 1996. Quantitative RT-PCR: limits and accuracy. *Biotechniques.* 21(2): 280–285.
- Takano E., Ma H., Yang H.Q., Maki M. and Hatanaka M. 1995. Preference of calcium-dependent interactions between calmodulin-like domains of calpain and alpastatin subdomains. *FEBS. Lett.* 362:93–97.
- Takano, E., M. Maki, H. Mori, M. Hatanaka, T. Marti, K. Titani, R. Kannagi, T. Ooi and T. Murachi. 1988. Pig heart calpastatin: identification of repetitive domain structures

and anomalous behavior in polyacrylamide gel electrophoresis. *Biochemistry*. 27(6):1964-1972.

- Takano, E. and Murachi T. 1982. Purification and some properties of human erythrocyte calpastatin. *J. Biochem.* 92:2021–2028.
- Takano, J., M. Watanabe, K. Hitomi and M. Maki. 2000. Four types of calpastatin isoforms with distinct amino-terminal sequences are specified by alternative first exons and differentially expressed in mouse tissues. *J. Biochem.* 128(1):83-92.
- Tradewell, M.L. and Durham H.D. 2010. Calpastatin reduces toxicity of SOD1G93A in a culture model of amyotrophic lateral sclerosis. *Neuroreport*. 21(15): 976-979.
- Wang L.F., Wei S.G., Miao S.Y., Liu Q.Y. and Koide S.S. Calpastatin gene in human testis. *Biochem. Mol. BiolInt.* 33:245–252.
- Warriss, P.D. 2000. *Meat Science*. UK: CAB International.
- Wheeler, T.L., Savell, J.W., Cross, H.R., Lunt, D.K. and Smith, S.B. 1990. Mechanism associated with the variation in tenderness of meat from Brahman and Hereford cattle. *J. Anim. Sci.* 68:4206-4220.
- Whipple, G., M. Koohmaraie, M.E. Dikeman, J.D. Crouse, M.C. Hunt and R.D. Klemm. 1990. Evaluation of attributes that affect longissimus muscle tenderness in *Bos taurus* and *Bos indicus* cattle. *J. Anim. Sci.* 68:2716.
- Wulf, D.M., J.D. Tatum, R.D. Green, J.B. Morgan, B.L. Golden and G.C. Smith. 1996. Genetic influences on beef longissimus palatability in Charolais- and Limousin-sired steers and heifers. *J. Anim. Sci.* 74:2394.

ภาคผนวก

ภาคผนวก ก

ลำดับนิวคลีโอไทด์ของยีนคาลปาสเตติน

ลำดับนิวคลีโอไทด์ของยีนคาลปาสเตตินจากโคสายพันธุ์ชาโรเลส BT-CAST1 (NM_001030318), BT_CAST2 (NM_174003), BT_CAST3 (NM_001030319) และ BT_CAST4 (NM_001030320) เปรียบเทียบกับโคพื้นเมืองไทยและโคกำแพงแสน

จุดไข่ปลา (...) แสดงนิวคลีโอไทด์ที่มีความอนุรักษ์เมื่อเทียบกับลำดับนิวคลีโอไทด์ของยีน BT_CAST1, กรอบสี่เหลี่ยม แสดงขอบเขตของโดเมน I - IV

แถบสีเหลือง แสดงตำแหน่ง start codon (ATG), **แถบสีเทา** แสดงตำแหน่ง stop codon (TAA), **แถบสีฟ้า** แสดงลำดับนิวคลีโอไทด์ที่จำแนกได้จากงานวิจัยนี้

BT_CAST1	GGAGCTCTCGCGGGTTCGGGGCTGGGGTCGGAAAGCTGCCACAGCGCGCCCGCCAGCCCTCCGCGCCCTCGCTCCCTCCAGCGCTCCCGGGCTCCA	100
BT_CAST2	-----G..TTAG...AGT.AG.TCTG.GG.AG.T.GAGTG..AAC..G	44
BT_CAST3	-----	1
BT_CAST4	-----	1
Thai_Native	-----	1
Kampaengsaen	-----	1

BT_CAST1	GCCITCCCITCCAGGCTCCGGCCGGCCCGGGGAGGCGGTCGACCCGGCCTCGCCATGTCACAGCCGGCCCGAAGCCCGCCGCTCCCGGGCCCC	200
BT_CAST2	-----TGGC..T...G...T..A....TGTCCTC..C..G..C.T.G..T...T.GC..CGT.TCT...AA...CAT..AT...G..AT...AT	135
BT_CAST3	-----	1
BT_CAST4	-----	1
Thai_Native	-----	1
Kampaengsaen	-----	1

BT_CAST1	CGGGCGCTGCCCGCCACCCAGGAGCATGTAGAAAAACAGTGGATCGCCCTCCAAATCAGGAGAAAAAGGATCAGATGAGAAAAAGCAA	300
BT_CAST2	-----TT..AA...GTG.T...AC.....	231
BT_CAST3	-----	1
BT_CAST4	-----	1
Thai_Native	-----	1
Kampaengsaen	-----	1



BT_CAST1 AAAATTGTTAAACATCGGCCGGTACCAGCTGAACTATAAACAAGTAAACCACTCTGGAAAGTCAGACA TGGACA CTGCCTCGGATGACTTAATAGACACCTTTAG 900
 BT_CAST2 831
 BT_CAST3 499
 BT_CAST4 1
 Thai_Native 1
 Kampaengsaen 1

BT_CAST1 GAGAACCTGAAGAGATGAAAGAAGATAACAACAATAACCGGACCGGAAGTGTCCGGATCCAA TGAGTTCTACCTACATAGAGAACTGGGTAAAAGAGA 1000
 BT_CAST2 931
 BT_CAST3 599
 BT_CAST4 1
 Thai_Native 1
 Kampaengsaen 1

BT_CAST1 ATCCACACTCTCCAAAATAAAGGAACTCTTGAAATAAAGAAGAAGGGATCGCAGGGCCCTCCTCCAGACTCCTTGAAAACCCCTGGGGCCCCAAATGATGCC 1100
 BT_CAST2 1031
 BT_CAST3 699
 BT_CAST4 1
 Thai_Native 1
 Kampaengsaen 1

BT_CAST1 ATCGATGCCCTTGTCATCCGACTTCACCTGCAGTCCCTTACAGCTGATGCAAAAGAAAAC TGAAAAGAGAAA TCTACAGAAGAGGCCTTAAAAGCTCAGT 1200
 BT_CAST2 1131
 BT_CAST3 799
 BT_CAST4 58
 Thai_Native 1
 Kampaengsaen 1

BT_CAST1 CAGCTGGGGTGA TCAGAAGTGTCTCCACCCCAAGAGAAAAAAGGAAAAGTGGAAAAGGATGCCATGACTGAGCAGCCCTGGAGGCCCTGT CAGCCCTC 1300
 BT_CAST2 1231
 BT_CAST3 899
 BT_CAST4 147
 Thai_Native 1
 Kampaengsaen 1

BT_CAST1 1400
 BT_CAST2 1331
 BT_CAST3 999
 BT_CAST4 240
 Thai_Native 1
 Kampaengsaen 1

```

CC TTGGGCACCCGGAAAGCCGGAGCCGGAGCTCGACCCAGCTCCA TTAAGGAGGTCGA TGAGGCAAAAGCCAAAGAGAGAAAGTAAAGAAA TGTGGTGAA
.....
TAC.AAGG.GAA.GCAAG...G...AG.GC..TTG.CCTCC...CC.A.....
  
```

BT_CAST1 1500
 BT_CAST2 1431
 BT_CAST3 1099
 BT_CAST4 340
 Thai_Native 1
 Kampaengsaen 1

```

GATGAGGAAACAGTCCCA TCGGAGTACAGATTAATAA CCGGCCACAGATAAAGATGGAAA CCACTTTGCCAGAGGC TGAAGAAAGACCCCAAGCCCC TGA
.....
.....A.....
  
```

BT_CAST1 1600
 BT_CAST2 1531
 BT_CAST3 1199
 BT_CAST4 440
 Thai_Native 1
 Kampaengsaen 1

```

GTGAA TCAGAACTCA TCGATGAACTCTCAGAAGA TTTTGACCAGTCTAAGTGTAAAGAAAA CAACTAAAGCCAA CTGAAAAAACAAGAGGCATCCCCGGCC
.....
.....
  
```

BT_CAST1 1700
 BT_CAST2 1631
 BT_CAST3 1299
 BT_CAST4 540
 Thai_Native 85
 Kampaengsaen 85

```

CGCTGCCCCCTGTACCCCGTGGCAGAGGACGTGCC TCAGAGGCTATGTGTTCCGTCAGTCGGCTCCGCCCCACAGCAGCTCCACGTGAAGGGCATGGTGCCCA
.....
.....G..CC.....C.....
  
```

BT_CAST1 1800
 BT_CAST2 1731
 BT_CAST3 1399
 BT_CAST4 640
 Thai_Native 185
 Kampaengsaen 185

```

GACGATGCTGTGAAGCCTTGGCTGGAAAGCC TGGCCAAAAGGAAGCAGATCCAGAAGACGGAAAAGCC TGGAGGATAAAGTCAAGGAAAAAGCCCAAAG
.....
.....C.....G.....
  
```

BT_CAST1
BT_CAST2
BT_CAST3
BT_CAST4
Thai_Native
Kampaengsaen

```
AAGAGGATCGTGAGAAACTTGGTGAAAAAGAAGAAACGATTCCCTCTGATACAGATTTAGAGAAGACCAAAGGATAAAAGACGGGAAAACCACTGCTGCCCAAA  
1900  
1831  
1499  
740  
285  
285
```

BT_CAST1
BT_CAST2
BT_CAST3
BT_CAST4
Thai_Native
Kampaengsaen

```
AGAGGTCAAGGACCGCTCCCACCCCTTGAGTGAAGCGTCCCTCTCGATGCTGTCCAAGGACTTCACTGTCCCCTCAGACACATCATCGCCTCAATTT  
2000  
1931  
1599  
840  
385  
385
```

BT_CAST1
BT_CAST2
BT_CAST3
BT_CAST4
Thai_Native
Kampaengsaen

```
GAAGATGCTAAACTTTCAGCTGTCGTCCTGAAGTGGTTTCCCAAACCCCAGCTCCAACCCCAGGCAGCCGGTCCACCCCCCCAGCACTGGCAGCGTG  
2100  
2031  
1699  
940  
485  
485
```


BT_CAST1
BT_CAST2
BT_CAST3
BT_CAST4
Thai_Native
Kampaengsaen

```
ACACAAAGAACTTGACCGATGCCCTGGATCAACTTTCTGACAGCTCGGGCAAGACAGCCCTGATCCAGATGAGAATAAACCCGTAGAGGATAAAGTCAA  
2200  
2131  
1799  
1040  
585  
585
```

BT_CAST1
BT_CAST2
BT_CAST3
BT_CAST4
Thai_Native
Kampaengsaen

```
GGAAAAGCCAAAAGCTGAACACAGAGACAAGCTGGGAGAAAGAGATGACCCATCCCACCTAAATACCAACACTTTTTGGATGACACACAAGGGGCCACA  
2300  
2231  
1899  
1140  
685  
685
```

BT_CAST1 C C C G G G A A G C C C A A A G G C A T C A G A G A A G C C C C A A G G C A T C A G A G A A C C T G C A G G T G C C C A G G A C C C C C A T T G A T G C C C C T C T C A G G G A C T T T G A C A G C T G T C 2400
BT_CAST2 2331
BT_CAST3 1999
BT_CAST4 1240
Thai_Native 785
Kampaengsaen 785



BT_CAST1 C C T C G A C T A C A G A A A C C T C G A C A G A C A C C A A A G G A C A A A G A C C T G C T T C C A G T G C C G A A G C A C C T A G G A A T G G C G G A A A G C A A A G G A T T C 2500
BT_CAST2 2431
BT_CAST3 2099
BT_CAST4 1340
Thai_Native 885
Kampaengsaen 885

BT_CAST1 C A C A A G G C A A A G G A A A C T T C C A A G C C A A A A G C T G A T G G A A A A G T A C A A G T T A A G T T C A C A C T A T T T G G T A T C T G C A T A T A A A A T C T T C A G C G G G 2600
BT_CAST2 2531
BT_CAST3 2199
BT_CAST4 1440
Thai_Native 985
Kampaengsaen 985

BT_CAST1 T A G A T G G T G A C T T C T G A A G A G A A A G G C T T T G A T A A C A G A A A C A A T T T C T G G G T G G C T T G G A G A C A G T G G T A T T T T G C T G A G T C T T T T G A C C T C C T A A A C 2700
BT_CAST2 2631
BT_CAST3 2299
BT_CAST4 1540
Thai_Native 1002
Kampaengsaen 1002

BT_CAST1 A T T G T C T G T A T C T T T T C C T G A A A G A A A C T G A A T T T G T C T G G T T C A C C T G T G T T A T T C T A C T G A G T A T T G A T A A A C T T T T A A A A T T T T A A A A T T G C C T T 2800
BT_CAST2 2731
BT_CAST3 2399
BT_CAST4 1640
Thai_Native 1002
Kampaengsaen 1002

BT_CAST1 CAGTTGGGAGAGAAAAGGAACTTTATAATTTCTAAGAGATACATTTGATAGTTTCTTAAAGCAGCACACAAAAAGGAAAAACCCTTTGCAAACTTTTGCACA 2900
BT_CAST2 2831
BT_CAST3 2499
BT_CAST4 1740
Thai_Native 1100
Kampaengsaen CTT 1100

BT_CAST1 TTCTCCCCACAGTGCCCTGTAAACTCTCATTAGTATTTTCAAATTTGGCACTATTTTTGTTGTTAGCATTTGGAAAAAGATGCCCTCACGGTTCITTCAGTGTT 3000
BT_CAST2 2931
BT_CAST3 2599
BT_CAST4 1840
Thai_Native 1200
Kampaengsaen 1200

BT_CAST1 CTGATTTCTCATGACCCCTTTCCCTTATAGACTTGGACTGTGTTTGAATGTTTCTTTGGGTTGTTTATAAGTCAGTCAFAAAAATCTGTGCAATTGGG 3100
BT_CAST2 3031
BT_CAST3 2699
BT_CAST4 1940
Thai_Native G 1300
Kampaengsaen G C 1300

BT_CAST1 CACATGTCCTCTTGAGCTGCTAACTGTAGAGACCCCTGGACAGACCAGGAAGCCGCCACCCCTTTTTCAGGTTGAACCTTCTTTGCCAGAGGTTAGG 3200
BT_CAST2 3131
BT_CAST3 2799
BT_CAST4 2040
Thai_Native 1325
Kampaengsaen 1325

ภาคผนวก ข
ข้อมูลและผลการวิเคราะห์ทางสถิติ

ค่าแรงตัดผ่านเนื้อในกล้ามเนื้อสันนอก

ตารางที่ ข1 ค่าแรงตัดผ่านเนื้อในกล้ามเนื้อสันนอกของโคพื้นเมืองและโคกำแพงแสน

ค่าแรงตัดผ่านเนื้อ (กก.ต่อตร.ซม.)		
ตัวอย่างที่	โคพื้นเมือง (NT)	โคกำแพงแสน (KU)
1	10.32	7.26
2	12.37	7.58
3	12.45	10.27
4	10.17	11.26
5	12.49	11.10
6	18.34	11.38
7	17.51	10.31
8	19.86	11.15
9	15.21	11.36
10	15.30	9.99
11	19.33	12.16
12	15.81	12.06
13	17.50	13.47
14	16.71	11.68
15	16.71	10.97
Mean	15.34	10.80
SD	3.12	1.62

การแสดงผลของยีนคาลิปาสดัตินในกล้ามเนื้อและอวัยวะโคพื้นเมือง

ตารางที่ ข2 ผลการประเมินประสิทธิภาพการเพิ่มปริมาณยีนคาลิปาสดัตินโดยเทคนิค Realtime PCR

สัดส่วนการเจือจาง	ซ้ำที่	ค่า E	ค่า CT	ค่า Tm
1:100	1	1.62	9.42	80.8
	2	1.70	9.62	80.6
1:1000	1	1.72	12.00	80.6
	2	1.58	12.11	80.8
1:10000	1	1.60	15.35	80.4
	2	1.54	15.14	80.6
1:100000	1	1.60	18.19	80.4
	2	1.55	18.07	80.4
ค่าเฉลี่ย		1.61		80.6

ตารางที่ ข3 ค่าเฉลี่ยปริมาณการแสดงออกของยีนคาลปาสเตตินในกล้ามเนื้อของโคพื้นเมือง*

ชนิดกล้ามเนื้อ	ค่า $E^{\Delta_{CP}(\text{actin-cast})}$	relative expression to LD
LD	168.54	1.00
TA	202.61	1.202
LT	179.86	1.067
SM	57.96	0.344
ST	169.23	1.004
RA	392.86	2.331

* n = 3

LD = Longissimus dorsi, LT= Longissimus troracis, TA = Transversus abdominis,
SM = Semimembranosus, ST = Semitendinosus and RA = Rectus abdominis

ตารางที่ ข4 ค่าเฉลี่ยปริมาณการแสดงออกของยีนคาลปาสเตตินในอวัยวะของโคพื้นเมือง*

อวัยวะ	ค่า $E^{\Delta_{CP}(\text{actin-cast})}$	relative expression to LD
อ้นทะ (T)	6.51	0.039
ลำไส้ (I)	28.40	0.169
ไต (K)	6.11	0.036
หัวใจ (H)	28.62	0.170
ม้าม (S)	5.60	0.033
สมอง (B)	0.39	0.002
ตับ (L)	1.18	0.007
กล้ามเนื้อสันนอก (LD)	168.54	1.000

* n = 3

ตารางที่ ๗5 ค่า $E^{\Delta CP(\text{actin-cast})}$ ในกล้ามเนื้อสันนอกของโคพื้นเมืองและโคกำแพงแสน

ค่า $E^{\Delta CP(\text{actin-cast})}$		
ตัวอย่างที่	โคพื้นเมือง (NT)	โคกำแพงแสน (KU)
1	142.43	116.29
2	150.44	116.29
3	140.71	71.78
4	72.36	98.88
5	124.09	193.05
6	216.26	98.08
7	205.15	86.67
8	220.86	141.28
9	162.16	70.62
10	113.50	71.49
11	161.50	114.42
12	201.85	90.81
13	264.86	108.55
14	189.17	152.59
15	162.82	92.30
Mean	168.54	108.21
SD	48.793	33.518

ค่ากิจกรรมการทำงานของเอนไซม์คาลปาสเตติน

ตารางที่ ข6 ค่าการดูดกลืนแสงของตัวอย่างที่ชะออกจากคอลัมน์ DEAE-Sephacel™

fraction no.	ค่าการดูดกลืนแสงที่ 280 nm
1	0.000
2	0.000
3	0.000
4	0.000
5	0.056
6	0.133
7	0.767
8	2.217
9	3.353
10	1.403
11	0.925
12	0.546
13	0.383
14	0.297
15	0.137
16	0.077
17	0.043
18	0.017
19	0.000
20	0.000

ตารางที่ ๗ ค่ากิจกรรมการทำงานของเอนไซม์คาลปาสเตดินในกล้ามเนื้อสันนอกของโคพื้นเมือง

ค่ากิจกรรมการทำงานของเอนไซม์คาลปาสเตดิน			
โคพื้นเมือง (NT)	Unit/100 μ l	Unit/17ml	Unit/g of meat
1	0.292	49.58	24.79
2	0.300	51.06	25.53
3	0.304	51.68	25.84
4	0.268	45.50	22.75
5	0.267	45.45	22.72
6	0.295	50.21	25.10
7	0.282	47.88	23.94
8	0.299	50.77	25.39
9	0.309	52.47	26.24
10	0.291	49.47	24.74
Mean	0.291	49.41	24.70
SD	0.014	2.428	1.216

ตารางที่ ข8 ค่ากิจกรรมการทำงานของเอนไซม์คาลปาสเตตินในกล้ามเนื้อสันนอกของโคก่าแพงแสน

ค่ากิจกรรมการทำงานของเอนไซม์คาลปาสเตติน			
โคก่าแพงแสน (KU)	Unit/100 μ l	Unit/17ml	Unit/g of meat
1	0.193	32.81	16.41
2	0.272	46.18	23.09
3	0.265	44.99	22.50
4	0.284	48.34	24.17
5	0.264	44.88	22.44
6	0.292	49.58	24.79
7	0.289	49.13	24.57
8	0.286	48.68	24.34
9	0.262	44.60	22.30
10	0.303	51.51	25.76
11	0.297	50.43	25.22
12	0.285	48.51	24.25
13	0.296	50.26	25.13
Mean	0.276	46.92	23.46
SD	0.028	4.805	2.402

การวิเคราะห์ค่าความแตกต่างทางสถิติของโคพื้นเมืองและโคกำแพงแสนโดย T-test

1) ค่าแรงตัดผ่านเนื้อในกล้ามเนื้อสันนอก

Variable	breed	N	Lower CL		Upper CL		Lower CL		Upper CL	
			Mean	Mean	Mean	Std Dev	Std Dev	Std Dev	Std Err	
sf	KU	15	9.9056	10.8	11.694	1.1825	1.6151	2.5472	0.417	
sf	NT	15	13.613	15.339	17.064	2.2814	3.1161	4.9144	0.8046	
sf	Diff (1-2)		-6.395	-4.539	-2.682	1.9695	2.4818	3.3565	0.9062	

T-Tests						
Variable	Method	Variances		DF	t Value	Pr > t
sf	Pooled	Equal		28	-5.01	<.0001
sf	Satterthwaite	Unequal		21	-5.01	<.0001

Equality of Variances						
Variable	Method	Num DF	Den DF	F Value	Pr > F	
sf	Folded F	14	14	3.72	0.0194	

2) ปริมาณการแสดงออกของยีนคาลปาสเตตินในกล้ามเนื้อสันนอก [ค่า $E^{\Delta_{CP} \text{actin-cast}}$]

Variable	breed	N	Lower CL		Upper CL		Lower CL		Upper CL	
			Mean	Mean	Mean	Std Dev	Std Dev	Std Dev	Std Err	
exp	KU	15	89.645	108.21	126.77	24.539	33.518	52.861	8.6543	
exp	NT	15	141.52	168.54	195.56	35.722	48.793	76.951	12.598	
exp	Diff (1-2)		-91.65	-60.34	-29.03	33.218	41.858	56.611	15.284	

T-Tests						
Variable	Method	Variances		DF	t Value	Pr > t
exp	Pooled	Equal		28	-3.95	0.0005
exp	Satterthwaite	Unequal		24.8	-3.95	0.0006

Equality of Variances						
Variable	Method	Num DF	Den DF	F Value	Pr > F	
exp	Folded F	14	14	2.12	0.1723	

3) ค่ากิจกรรมการทำงานของเอนไซม์คาลปาสเตตินในกล้ามเนื้อสันนอก

Variable	breed	N	Lower CL		Upper CL		Lower CL		Upper CL	
			Mean	Mean	Mean	Std Dev	Std Dev	Std Dev	Std Err	
acti	KU	13	22.008	23.459	24.911	1.7223	2.4019	3.9648	0.6662	
acti	NT	10	23.834	24.704	25.574	0.8364	1.2159	2.2198	0.3845	
acti	Diff (1-2)		-2.979	-1.245	0.4894	1.5252	1.9825	2.8331	0.8339	

T-Tests						
Variable	Method	Variances		DF	t Value	Pr > t
acti	Pooled	Equal		21	-1.49	0.1504
acti	Satterthwaite	Unequal		18.6	-1.62	0.1224

Equality of Variances						
Variable	Method	Num DF	Den DF	F Value	Pr > F	
acti	Folded F	12	9	3.90	0.0487	

ผลการวิเคราะห์ค่าสหสัมพันธ์ Pearson Correlation Coefficient

1) The CORR Procedure โคพื้นเมือง

3 Variables: sf exp acti

Simple Statistics

Variable	N	Mean	Std Dev	Sum	Minimum	Maximum
sf	15	15.33867	3.11612	230.08000	10.17000	19.86000
exp	15	168.54400	48.79271	2528	72.36000	264.86000
acti	10	24.70400	1.21592	247.04000	22.72000	26.24000

Pearson Correlation Coefficients

Prob > |r| under H0: Rho=0

Number of Observations

	sf	exp	acti
sf	1.00000	0.73633 0.0017 15	-0.28037 0.4327 10
exp	0.73633 0.0017 15	1.00000	-0.35621 0.3124 10
acti	-0.28037 0.4327 10	-0.35621 0.3124 10	1.00000 10

2) The CORR Procedure โคกำแพงแสน

3 Variables: sf exp acti

Simple Statistics

Variable	N	Mean	Std Dev	Sum	Minimum	Maximum
sf	15	10.80000	1.61512	162.00000	7.26000	13.47000
exp	15	108.20667	33.51793	1623	70.62000	193.05000
acti	13	23.45923	2.40186	304.97000	16.41000	25.76000

Pearson Correlation Coefficients

Prob > |r| under H0: Rho=0

Number of Observations

	sf	exp	acti
sf	1.00000	0.04562 0.8718 15	0.65533 0.0150 13
exp	0.04562 0.8718 15	1.00000	-0.17575 0.5657 13
acti	0.65533 0.0150 13	-0.17575 0.5657 13	1.00000 13

ภาคผนวก ค
ผลงานวิจัยที่เผยแพร่

การประชุมนักวิจัยรุ่นใหม่พบเมธีวิจัยอาวุโส สกว. ในปี พ.ศ. 2550



Calpastatin Gene Study in Thai Native Cattle (*B. indicus*) and Kampaengsaen Cattle (*B. indicus* x *B. taurus*)

Jirajaroenrat K.*, Opatpatanakit Y., Sethakul J.

Department of Animal Production, Faculty of Agricultural Technology,
King Mongkut's Institute of Technology Ladkrabang, Bangkok, Thailand 10520
*e-mail: kjkanya@kmitl.ac.th

Introduction

Calpastatin is a specific protein inhibitor of the calcium dependent proteinases μ - and m -calpain^{1,2}. This enzyme system is known to be involved in meat tenderization, which is especially important in beef quality³. Four alternatively spliced mRNA transcripts of calpastatin gene (Type I, II, III and IV) have been reported in *Bos taurus* cattle⁴. The difference between them resides in their 5'-termini. The corresponding data for *B. indicus* has not been reported yet.

Objective

To study the calpastatin mRNA transcription in Thai native cattle (*B. indicus*) and Kampaengsaen cattle, the Thai developed *B. indicus* x *B. taurus* crossbred, in order to understand the relationship between calpastatin regulation and meat quality.

Method

The calpastatin cDNAs were cloned from skeletal muscle (Longissimus dorsi) by RT-PCR. Specific primers were designed based on the four *B. taurus* calpastatin mRNAs (Figure 1). Methodology flow chart is shown in Figure 2.




Figure 1 Primer design based on *B. taurus* mRNA transcripts (type I-IV)




Figure 2 Research methodology flowchart

Conclusion

The 3'-termini of calpastatin mRNA transcripts in *B. taurus* and *B. indicus* share highly conserved sequence. Therefore this gene region may be important for enzyme function. The 5'-termini of the mRNAs in both Thai breeds and their expression in different muscles and tissues are under further investigation.

Acknowledgement

- Centre of Agricultural Biotechnology Consortium for lab equipments
- Thailand Research Fund for financial support

Result

Two cDNA fragments of Thai native and Kampaengsaen breeds were amplified (Figure 3) and sequenced (Figure 4), labelled **CONS1** and **CONS2** fragments, corresponding to the 3'-end of coding region and the 3'-untranslated region of calpastatin gene respectively. The CONS1 fragments of both breeds (1,002 bp) shared 100% of the nucleotide identity with the *B. taurus* transcripts and the CONS2 fragments (323 bp) shared 98% and 99% of nucleotide identity with the *B. taurus* transcripts respectively.




Figure 3 RT-PCR products of Thai native (a) and Kampaengsaen (b) cattle no. 1, 2 and 3 with primer sets: CONS-F1/CONS-R1 (A), CONS-F2/CONS-R2 (B) and CAST-F5/CAST-R1-4 (C).




Figure 4 Calpastatin sequence alignment of *Bos taurus*: BT-CAST1 (NM_001030318), BT-CAST2 (NM_174003), BT-CAST3 (NM_001030319) and BT-CAST4 (NM_001030320), with Thai native and Kampaengsaen breeds

Reference

1. Koohmaraie M. 1988. *Proc. Recip. Meat Conf.* 41:89.
2. Koohmaraie M. 1992. *Proc. Recip. Meat Conf.* 45:63.
3. Whipple G., et al. 1990. *J. Anim. Sci.* 68:2716.
4. Raynaud P., et al. 2005. *Arch. Biochem. Biophys.* 437:69-77.



การประชุมนักวิจัยรุ่นใหม่พบเมธีวิจัยอาวุโส สกว. ในปี พ.ศ. 2551



Calpastatin Expression in Different Muscles of Thai Native Cattle

Jirajaroenrat, K.^{1*}, Opatpatanakit, Y.² and Sethakul, J.¹

¹Department of Animal Production, Faculty of Agricultural Technology, KMITL, Bangkok, Thailand 10520

²Department of Animal Technology, Faculty of Agricultural Production, Maejo University, Chiangmai, Thailand 50290

*e-mail: kjkanya@kmitl.ac.th

Introduction

Calpastatin is a specific protein inhibitor of the calcium dependent proteinases μ - and m-calpain[1-2]. This enzyme system is known to be involved in meat tenderization, which is especially important in beef quality[3]. Four alternatively spliced mRNA transcripts of calpastatin gene (Type I-IV) have been reported in European breed (*Bos taurus*) cattle. Three calpastatin isoforms migrated at 145, 125 and 70 kDa were detected[4]. Thai native cattle were identified to be a member of *Bos indicus* specie. Little molecular information on this specie has been reported.

Objective

To characterize the pattern of calpastatin expression in different muscles of Thai native beef cattle (*B. indicus*) in order to understand the relationship between calpastatin regulation and meat quality.

Method

Muscle Extraction: The protein were extracted from six muscles; semimembranosus (SM), rectus abdominis (RA), semitendinosus (ST), longissimus traracis (LT), longissimus dorsi (LD) and transversus abdominis (TA) according to the method described by Delgado *et al.* [5].

SDS-PAGE: Protein concentration in the supernatant fraction was diluted to 3 mg/mL. Protein amount of 50 μ g were separated by SDS-PAGE using 7 x 10 cm minigels. After staining, the protein bands were visualized comparing to prestained molecular weight marker (Invitrogen).

Western blot analysis: The seperated proteins were transferred to Hybond-P PVDF membrane. The 1st antibody used was a monoclonal antibody (Anti-DomIV; Sigma) at 1:500 dilution The polypeptides labeled with the 1st antibody were detected by using an polyclonal anti-mouse IgG (Fab fragments; Roche) conjugated with horseradish peroxidase at a 50 mU/ml. The 2nd antibody was detected by using TMB solution (3,3',5,5'- tetramethyl-benzidine: Roche). The image of the membrane was captured using gel document system (Syngene).

Result

We previously isolated the calpastatin mRNA transcripts by RT-PCR from skeletal muscles of Thai native cattle. About 1,500 bp of 3'-termini cDNA was isolated. The sequences shared 98-99% to that of the European cattle.

The muscle extracts were separated by SDS-PAGE (Figure 1). The expression of calpastatin isoforms in six muscles analysed by Western blotting showed different patterns. The 125 kDa and 47 kDa isoforms corresponding to the translation of Type III transcript, the erythrocyte isoform, respectively, were detected in SM and RA muscles, whereas LT, LM and TA muscles contained only 47 kDa isoform. Interestingly, the ST muscle contained the 36 kDa isoform the C-terminal proteolytic product of the calpastatin catalyzed by calpain enzymes (Figure 2).




Figure 1. SDS-PAGE of protein extracts from 6 muscles of Thai native beef cattle: semimembranosus (SM), rectus abdominis (RA), semitendinosus (ST), longissimus traracis (LT), longissimus dorsi (LD) and transversus abdominis (TA) respectively. Lane M is a prestained protein molecular weight and lane A is bovine serum albumin 2 ug.





Figure 2. Western blotting of protein extracts from 6 muscles of Thai native beef cattle: semimembranosus (SM), rectus abdominis (RA), semitendinosus (ST), longissimus traracis (LT), longissimus dorsi (LD) and transversus abdominis (TA) respectively, against anti-calpastatin domain IV. Lane M is a standard protein molecular weight.

Conclusion

Different patterns of the calpastatin expression in different muscles were confirmed by Western blotting. Three calpastatin isoforms: 125, 47 and 36kDa bands were detected. Relationship of the calpastatin expression and its mRNA transcription will be further investigated.

Acknowledgement

We thank the Centre of Agricultural Biotechnology, Faculty of Agricultural Technology, KMITL for providing lab equipments and Thailand Research Fund for financial support.



References

- [1] Koohmaraie. 1988. Proc. Recip. Meat Conf. 41:89.
- [2] Koohmaraie. 1992. Proc. Recip. Meat Conf. 45:63.
- [3] Whipple *et al.* 1990. J. Anim. Sci. 68:2716.
- [4] Raynaud *et al.* 2005. Arch. Biochem. Biophys. 437:69-77.
- [5] Delgado *et al.* 2001. J. Anim. Sci. 79:398-412.

