


บรรณานุกรม



สำนักวิทยบริการ, สถาบันราชภัฏบ้านสมเด็จเจ้าพระยา. เอกสารสมุนไพรไทย. กรุงเทพมหานคร

- Arima, K.¹, Uchikoba, T., Yonezawa, H., Shimada, M., and Kaneda, M. (2000). Isolation and characterization of a serine protease from the sprouts of *Pleioblastus hindsii* Nakai. Phytochemistry. 54. 559-565.
- Arima, K.², Uchikoba, T., Yonezawa, H., Shimada, M., and Kaneda, M. (2000). Cucumisin-like protease from the latex of *Euphorbia supina*. Phytochemistry. 53. 639-644.
- Asif-Ullah, M., Kim, K.S., and Yu, Y.G. (2006). Purification and characterization of a serine protease from *Cucumis trigonus* Roxburghi. Phytochemistry. 67. 870-875.
- Chantaphakul, H., Tiyasatapon, S., Phanupak, P., Ruxrungtham, K. (2002). Jackfruit induced anaphylaxis and latex allergy. J Allergy Clin Immunol. 109. S219.
- Devaraja, S., Girish, K.S., Gowtham, Y.N., and Kemparaju, K. (2011). The Hag-protease-II is a fibrin(ogen)ase from *Hippasa agelenoides* spider venom gland extract: purification, characterization and its role in hemostasis. Toxicon. 57. 248-258.
- Huang, K. F., Hung, C. C., and Chiou, S. H. (1993). Characterization of three fibrinogenolytic proteases isolated from the venom of Taiwan habu (*Trimeresurus mucrosquamatus*). Biochem Mol Biol Int. 31. 1041-1050.
- Khan, M. R., Omoloso, A. D., and Kihara, M. (2003). Antibacterial activity of *Artocarpus heterophyllus*. Fitoterapia. 74. 501-505.
- Kobayashi, T., Kobayashi, E., Sato, S., Hotta, Y., Miyajima, N., Tanaka, A., and Tabata, S. (1994). Characterization of cDNAs induced in meiotic prophase in lily microsporocytes. DNA Res. 1. 15-26.
- Matsubara, K., Hori, K., Matsuura, Y., and Miyazawa, K. (2000). Purification and characterization of a fibrinolytic enzyme and identification of fibrinogen clotting enzyme in a marine green alga, *Codium divaricatum*. Comp Biochem Physiol B Biochem Mol Biol. 125. 137-143.
- Mekkriengkrai, D., Ute, K., Swiezewska, E., Chojnacki, T., Tanaka, Y., and Sakdapipanich, J. T. (2004). Structural characterization of rubber from jackfruit and euphorbia as a model of natural rubber. Biomacromolecules. 5. 2013-2019.
- Patel, G. K., Kawale, A. A., and Sharma, A. K. (2012). Purification and physicochemical characterization of a serine protease with fibrinolytic activity from latex of a medicinal herb *Euphorbia hirta*, Plant Physiol Biochem. 52. 104-111.

- Patiño, A. C., Pereañez, J. A., Núñez, V., Benjumea, D. M., Fernandez, M., Rucavado, A., Sanz, L., and Calvete, J.J. (2010). Isolation and biological characterization of Batx-I, a weak hemorrhagic and fibrinogenolytic PI metalloproteinase from Colombian *Bothrops atrox* venom. Toxicon. 56. 936-943
- Povovic, T., Puizdar, V., and Brzin, J. (2002). A novel subtilase from common bean leaves. FEBS Lett. 530. 163–168.
- Prasad, R. K. M., and Virupaksha, T. K. (1990). Purification and characterization of a protease from jackfruit latex. Phytochemistry. 29. 1763-1766.
- Rajesh, R., Nataraju, A., Gowda, C. D., Frey, B.M., Frey, F. J., and Vishwanath, B. S. (2006). Purification and characterization of a 34-kDa, heat stable glycoprotein from *Synadenium grantii* latex: action on human fibrinogen and fibrin clot. Biochimie. 88. 1313-1322.
- Satake, M., Murata, Y., and Suzuki, T. (1963). Studies on snake venom. XIII. Chromatographic separation and properties of three proteinases from *Agkistrodon halys blomhoffii* venom. J Biochem. 53. 438-447.
- Sharma, A., Kumari, M., and Jagannadham, M.V. (2009). Benghalensin, a highly stable serine protease from the latex of medicinal plant *Ficus benghalensis*. J Agric Food Chem. 57. 11120-11126.
- Siritapetawee, J., Thammasirirak, S., and Samosornsuk, W. (2012). Antimicrobial activity of a 48-kDa protease (AMP48) from *Artocarpus heterophyllus* latex, Eur Rev Med Pharmacol Sci. 16.132-137.
- Tornero, P., Conejero, V., and Vera, P. (1997). Identification of a new pathogen-induced member of the subtilisin-like processing protease family from plants. J Biol Chem. 272. 14412–14419.
- Uchikoba, T., Yonezawa, H., and Kaneda, M. (1998). Cucumisin like protease from the sarcocarp of *Benincase hispida* var, RyuKyu. Phytochemistry. 49. 2215–2219.
- Wei, Q., Jin, Y., Lu, Q. M., Wei, J. F., Wang, W. Y., and Xiong, Y. L. (2002). Purification and characterization of alpha-mucrofibrase, a novel serine protease with alpha-fibrinogenase activity from the venom of Chinese Habu (*Trimeresurus mucrosquamatus*). J Nat Toxins. 11. 337-343.
- Wetprasit, N., Threesangsri, W., Klamklai, N., and Chulavatnatol, M. (2000). Jackfruit lectin: properties of mitogenicity and the inhibition of herpesvirus infection. Jpn J Infect Dis. 53. 156-161.
- Yadav, S. C., Pande, M., and Jagannadham, M.V. (2006). Highly stable glycosylated serine protease from the medicinal plant *Euphorbia milii*. Phytochemistry. 67. 1414-1426.
- Yamagata, H., Masuzawa, T., Nagaoka, Y., Ohnishi, T., and Iwasaki, T. (1994). Cucumisin, a serine protease from melon fruit, shares structural homology with subtilisin and is generated from a large precursor. J Biol Chem. 269. 32725–32731.

Yamagata, H., Uesugi, M., Saka, K., Iwasaki, T. and Aizono, Y. (2000). Molecular cloning and characterization of a cDNA and a gene for subtilisin-like serine proteases from rice (*Oryza sativa* L.) and *Arabidopsis thaliana*. Biosci Biotechnol Biochem. 64 1947–1957.