

## LIST OF TABLES

<b>Table</b>		<b>Page</b>
1	Phenotypes of PGPR/PGPB, <i>Bacillus amyloliquefaciens</i> wild-type strain KPS46 and UV mutant strains	61
2	Identified proteins from the extracellular secretome of <i>Bacillus amyloliquefaciens</i> KPS46	66
3	Effect of plant growth promoting rhizobacteria/ bacteria, <i>Bacillus amyloliquefaciens</i> KPS46 compare with two reference PGPR and salicylic acid on reduction disease severity of soybean bacterial pustule caused by <i>Xanthomonas axonopodis</i> pv. <i>glycines</i> ( <i>Xag</i> )	71
4	Identified difference proteins from soybean after induced by <i>Bacillus amyloliquefaciens</i> KPS46 compared with negative control at 7 day after planting	80
5	Identified difference proteins from soybean after induced by <i>Bacillus amyloliquefaciens</i> KPS46 compared with negative control, and challenge inoculation with <i>Xanthomonas axonopodis</i> pv. <i>glycines</i> at 14 day after planting	84
6	Effect of KPS46 application on disease progression in Arabidopsis wild type and mutant line inoculated with virulent <i>Pseudomonas syringae</i> pv. <i>tomato</i> strain DC3000	90