

## LIST OF FIGURES

| <b>Figure</b> |   | <b>Page</b> |
|---------------|---|-------------|
| 1             | The Brønsted acid site of catalytic zeolite   | 2           |
| 2             | Various types of zeolite pores and channels   | 2           |
| 3             | The alkylation of benzene with ethylene to ethylbenzene and the dehydrogenation of ethylbenzene to styrene  | 4           |
| 4             | Dimerization of ethylene to butene  | 4           |
| 5             | The isomerization of propene oxide to propanal and propanone  | 5           |
| 6             | a) Single-walled carbon nanotube (SWNT) bundle and b) Multi-walled carbon nanotubes (MWNT)  | 7           |
| 7             | Zigzag (9,0) CNT with a) Perfect and b) Stone-Wales defect (indicated in yellow color)  | 8           |
| 8             | Decomposition of nitrous oxide (N <sub>2</sub> O) over nanotube to nitrogen molecule (N <sub>2</sub> ) and epoxy-adduct   | 9           |
| 9             | A self-consistent field procedure used to solve the wavefunctions of the Schrödinger equation   | 40          |
| 10            | ONIOM3 layer of 84T cluster models of faujasite. Atoms belonging to the 12T quantum cluster are drawn as bonds and sticks. (a) Front view, (b) side view  | 49          |
| 11            | The 5T and 46T clusters represent the active site of ZSM-5 zeolite in the zigzag channel view. The ball-and-stick graphics illustrates the relaxed geometry resulting from calculations at B3LYP/6-31G(d,p), the lines from the universal force field (UFF) | 51          |
| 12            | Representation of the fragment tube of (a) perfect and (b) defective armchair (5,5)-SWNTs calculated at B3LYP/6-31G(d)  | 53          |

## LIST OF FIGURES (continued)

| Figure |   | Page |
|--------|---|------|
| 13     | Presentation of models of faujasite and interacting with adsorbates:<br>a) full 3T cluster model interacting with ethylene; b) full 3T cluster model interacting with benzene; and c) full 3T cluster model interacting with ethylbenzene   | 56   |
| 14     | Presentation of models of faujasite and interacting with adsorbates:<br>a) ONIOM2 layer models of 20T cluster interacting with ethylene; b) ONIOM2 layer models of 20T cluster interacting with benzene; and c) ONIOM2 layer models of 20T cluster interacting with ethylbenzene. Atoms belonging to the high level regions are drawn as spheres  | 57   |
| 15     | Presentation of models of faujasite and interacting with adsorbates:<br>a) ONIOM2 layer models of 84T cluster interacting with ethylene; b) ONIOM2 layer models of 84T cluster interacting with benzene; and c) ONIOM2 layer models of 84T cluster interacting with ethylbenzene. Atoms belonging to the high level regions are drawn as spheres. | 58   |
| 16     | (a) Calculated energy profile for the stepwise reactions in steps (1) and (2) of benzene alkylation and dimerization of ethylene. (b) Vibrational movement corresponding to the imaginary frequency at the transition structure of step (2)   | 66   |
| 17     | (a) Calculated energy profile for the second step of stepwise reactions for benzene alkylation. (b) Vibrational movement corresponding to the imaginary frequency at the transition structure   | 71   |
| 18     | (a) Calculated energy profile for the second step of stepwise for dimerization of alkoxide intermediate and ethylene molecule. (b) Vibrational movement corresponding to the imaginary frequency at the transition structure  | 73   |

**LIST OF FIGURES (continued)**

| <b>Figure</b> |  | <b>Page</b> |
|---------------|--|-------------|
| 19            | (a) Calculated energy profile for the concerted of benzene dimerization. (b) Vibrational movement corresponding to the imaginary frequency at the transition structure   | 76          |
| 20            | Calculated energetic profiles for the stepwise (solid line) and concerted (dashed line) reaction mechanisms of benzene alkylation with ethylene  | 79          |
| 21            | a) Calculated energy profile for the concerted mechanism of ethylene dimerization. b) Vibrational movement corresponding to the imaginary frequency at the transition structure  | 81          |
| 22            | Calculated energy profiles for the stepwise (solid line) and concerted (dashed line) reaction mechanisms of ethylene dimerization  | 83          |
| 23            | Presentation of models of ZSM-5 and its interaction with adsorbates: ONIOM2 layer models of 46T cluster interacts with (a) propene oxide, (b) propanal, and (c) propanone  | 87          |
| 24            | a) The calculated energy profile for the isomerization of propene oxide to propanal over 46T and 5T (in parentheses) clusters; b) the vibrational movement corresponding to the imaginary frequency at the TS1_A; c) the vibrational movement corresponding to the imaginary frequency at the TS2_A  | 91          |
| 25            | a) The calculated energy profile for the isomerization of propene oxide to propanone over 46T and 5T (in parentheses) clusters; b) the vibrational movement corresponding to the imaginary frequency at the TS1_B; c) the vibrational movement corresponding to the imaginary frequency at the TS2_B | 95          |

**LIST OF FIGURES (continued)**

| <b>Figure</b> |  | <b>Page</b> |
|---------------|--|-------------|
| 26            | The energy profile of the isomerization of propene oxide to propanal on H-ZSM-5 zeolite (solid line), to propanone on H-ZSM-5 zeolite (dash line) calculated at ONIOM(B3LYP/6-31G(d,p):UFF), to propanal in gas phase (dash dot dot line), and to propanone in gas phase (dot line) calculated at B3LYP/6-31G(d,p)                               | 96          |
| 27            | The reaction complexes associated with the stepwise mechanism for the decomposition of nitrous oxide to nitrogen on the perfect site of the armchair (5,5)-SWNT. (a) TS1p, (b) LM1p, (c) TS2p, and (d) LM2p. The values in parenthesis are the relative energies (in kcal/mol) of each individual complex with respect to the isolated molecules | 99          |
| 28            | The reaction complexes associated with the concerted mechanism for the decomposition of nitrous oxide to nitrogen on the perfect site of the armchair (5,5)-SWNT. a) TSp, and (b) LM2p. The values in parenthesis are the relative energies (in kcal/mol) of each individual complex with respect to the isolated molecules                      | 101         |
| 29            | The reaction complexes associated with the stepwise mechanism for the decomposition of nitrous oxide to nitrogen on the Stone–Wales defective site in the armchair (5,5)-pSWNT. The values in parenthesis are the relative energies of each individual complex with respect to the isolated molecules  | 103         |
| 30            | The reaction complexes associated with the concerted mechanism for the decomposition of nitrous oxide to nitrogen on the Stone–Wales defective site in the armchair (5,5)-pSWNT. The values in parenthesis are the relative energies of each individual complex with respect to the isolated molecules   | 104         |

**LIST OF FIGURES (continued)**

| <b>Figure</b> |  | <b>Page</b> |
|---------------|--|-------------|
| 31            | Energy profiles for the nitrous oxide decomposition on pSWNT and dSWNT fully optimized at B3LYP/6-31G(d) | 106         |