THESIS

PHOTOCATALYTIC DEGRADATION OF TRYPAN BLUE USING GOLD/TITANIUM DIOXIDE

PIMPORN PHOACHARERN

A Thesis Submitted in Partial Fulfillment of the Requirements for the Degree of Master of Science (Chemistry) Graduate School, Kasetsart University 2006

ISBN 974-16-1885-9

Pimporn Phoacharern 2006: Photocatalytic Degradation of Trypan
Blue Using Gold/Titanium Dioxide. Master of Science (Chemistry),
Major Field: Chemistry, Department of Chemistry.
Thesis Advisor: Associate Professor Pakawadee Sutthivaiyakit, Ph.D.
76 pages.
ISBN 974-16-1885-9

An azo dyes are the largest group of the dyes used in dyeing industry, and the use of them is that they must be highly stable. Therefore, their remaining in effluents causes the environmental problem, moreover, azo dyes may be decomposed into carcinogenic compounds. Although photocatalytic degradation with titanium dioxide (TiO₂) is an efficient process, there are a few drawbacks. The objective of this research is to enhance the efficient of TiO₂ and utilize the solar light for treatment of Trypan Blue with gold deposition.

Gold deposited TiO₂ has been prepared by reducing $[AuCl_4]$ on the surface of preformed TiO₂ colloids using sodium borohydride in acetonitrile. The effect of gold contents, calcination temperature, initial pH of Trypan Blue, gold/TiO₂ dosage and dye concentration on the photocatalytic activity under a solar light simulating source (Xe-arc lamp, 150W) radiation using Trypan Blue were investigated. The change in dye concentration and formation of intermediates were monitored as a function of irradiation time. The rate of disappearance of dye follows first order kinetics. The degradation rate of 50 ppm Trypan Blue is complete within 50 min in the presence of 0.36% gold/TiO₂ that faster than in the presence of TiO_2 more seven folds, focusing at first 30 min of irradiation. In addition, this method has been applied to the wastewater. The Trypan Blue which spiked in wastewater disappears in 2 h. As a result, gold doped on TiO₂ could improve the photodegradation efficiency of Trypan Blue, although, it required longer irradiation time in the case of wastewater sample.

ACKNOWLEDGEMENTS

I wish to express my sincere gratitude to my advisor, Associate Professor. Dr. Pakawadee Sutthivaiyakit for her valuable guidance, inspiration, supervision and encouragement throughout the course of my postgraduate study. I am deeply grateful to Mrs.Saijai Charnsethikul, Assistant Professor Chongkolnee Kaewsriprakai and Associate Professor Saisuda Somchit for their valuable suggestions as thesis committee.

I would like to thank the Department of Chemistry and its staffs, Faculty of Science, Kasetsart University for making this research program possible, and to Postgraduate Education and Research Program in Chemistry (PERCH) for providing financial support.

Finally, I would also like to thank my family and friends for their encouragement and understanding.

Pimporn Phoacharern May 2006