

**THESIS**

**PHYSIOLOGY AND MOLECULAR BIOLOGY OF ETHYLENE  
SYNTHESIS IN DENDROBIUM FLOWERS FOLLOWING COMPATIBLE  
AND INCOMPATIBLE POLLINATION**

**KANJANA LUANGSUWALAI**

**A Thesis Submitted in Partial Fulfillment of  
the Requirements for the Degree of  
Doctor of Philosophy (Horticulture)  
Graduate School, Kasetsart University  
2007**

**ISBN 974-16-1039-4**

Kanjana Luangsuwalai 2007: Physiology and Molecular Biology of Ethylene Synthesis in *Dendrobium* Flowers Following Compatible and Incompatible Pollination. Doctor of Philosophy (Horticulture), Major Field: Horticulture, Department of Horticulture. Thesis Advisor: Professor Saichol Ketsa, Ph.D. 89 pages.  
ISBN 974-16-1039-4

Open flowers of *Dendrobium* 'Miss Teen' (or 'Kenny') were cross pollinated by placing pollinia from five cultivars of *Dendrobium* namely 'Miss Teen', 'Karen', 'Sakura', 'Willie', and 'Pompadour'. The pollinated flowers were held in distilled water at ambient temperature. The characteristics of ethylene production and post-pollination development were monitored. The pollinated flowers were divided into two groups. The first group was compatible pollination which consisted of flowers pollinated by pollinia from 'Sakura', 'Willie', and 'Pompadour'. In this group, the premature senescence of perianth took place within a day. The second group was incompatible pollination which consisted of flowers pollinated by 'Karen' and 'Miss Teen' pollinia. Incompatible pollinated flowers did not exhibit postpollination symptoms. Compatible pollination of 'Miss Teen' flowers resulted in an immediate burst of ethylene production and a rapid stimulation of ovary growth followed by an increase in pollen tube length. In contrast, when 'Miss Teen' flowers were self-pollinated or pollinated with 'Karen' pollinia, these changes did not occur. Pollinia of the cultivars 'Pompadour', 'Sakura' and 'Willie' had greater amounts of auxin and 1-aminocyclopropane -1- carboxylic acid (ACC) than that of the 'Miss Teen' and 'Karen' pollinia. Compatible pollination resulted an increment in higher ACC synthase and ACC oxidase activities than that of incompatible pollination. ACC oxidase activities of compatibly pollinated flowers rapidly increased within 12 h after pollination. Nonpollinated flowers showed a low ACC oxidase activity, while, column plus pedicel contained higher ACC oxidase activity than petal plus sepal and lip. *ACO* mRNAs was highly up-regulated by compatible pollination in the *Dendrobium* 'Miss Teen'.

Soaking pollinia in distilled water before pollination reduced ovary growth of pollinated flowers and lowered auxin content in soaked pollinia. Application of auxin transport inhibitors, 2, 3, 5-triiodobenzoic acid (TIBA) and  $\alpha$ -p-chlorophenoxy isobutyric acid (PCIB), delayed effect of pollination induced ovary growth. In addition, pretreatment with ethylene inhibitors, aminooxyacetic acid (AOA) and/or 1-methylcyclopropene (1-MCP), completely inhibited effect of pollination. This indicated that both auxin and ethylene are necessary for postpollination development. Application of octanoic acid to the stigma of *Dendrobium* 'Miss Teen' prior to pollination by 'Karen' pollinia or application of 1-aminocyclopropane -1- carboxylic acid and/or ethylene did not increase the sensitivity to ethylene of 'Miss Teen' flowers. All treated flowers exhibited indifferent symptoms from flowers without octanoic acid application. This observation indicated that octanoic acid was not the ethylene sensitivity factor of pollinated *Dendrobium* 'Miss Teen' flowers

---

Student's signature

---

Thesis Advisor's signature

## ACKNOWLEDGMENTS

I would like to thank many people without their contributions this thesis would not have been done. I am deeply indebted to them. First, I would like to express my deepest appreciation to my advisor Professor Dr. Saichol Ketsa for his great patient, guidance, knowledge, and spiritual support throughout my program of study and completion of this research.

I extend my special thanks to Assistant Professor Dr. Lop Phavaphutanon, Dr. Orawan Chatchawankanphanich and Professor Dr. Jean Cluade Pech for their intellectual support, encouragement, comments, and scholarly reading of this research. I greatly appreciated their advice insight and perspective.

I am indebted to the Royal Golden Jubilee Program under the Thailand Research Fund and the Postgraduate Education and Research Development Project in Postharvest Technology for awarding me a scholarship to undertake this study. The staff of Postharvest Research Unit, Central Laboratory and Greenhouse Complex, Kamphang Saen Campus, Nakorn Pathom deserves special thanks for their help.

I would like to thank Dr. Parichart Burns for her kind advice and help. Lastly, I would like to thank all current and past members of our laboratory for their help and friendship throughout the study.

Kanjana Luangsuwalai

December 2006